

Evolution of thermal tolerance in multifarious environments

Cuenca Cambroneró, Maria; Beasley, Jordan; Kissane, Stephen; Orsini, Luisa

DOI:

[10.1111/mec.14890](https://doi.org/10.1111/mec.14890)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Cuenca Cambroneró, M, Beasley, J, Kissane, S & Orsini, L 2018, 'Evolution of thermal tolerance in multifarious environments', *Molecular Ecology*, vol. 27, no. 22, pp. 4529-4541. <https://doi.org/10.1111/mec.14890>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Evolution of thermal tolerance in multifarious environments

Maria Cuenca Cambroner¹ | Jordan Beasley^{1,2} | Stephen Kissane¹ | Luisa Orsini¹ 

¹Environmental Genomics Group, School of Biosciences, The University of Birmingham, Birmingham, UK

²Department of Genetics, University of Leicester, Leicester, UK

***Correspondence**

Luisa Orsini, Environmental Genomics Group, School of Biosciences, University of Birmingham, Birmingham, UK.
Email: l.orsini@bham.ac.uk

Funding information

Natural Environment Research Council, Grant/Award Number: NE/N016777/1

Abstract

Species extinction rates are many times greater than the direst predictions made two decades ago by environmentalists, largely because of human impact. Major concerns are associated with the predicted higher recurrence and severity of extreme events, such as heat waves. Although tolerance to these extreme events is instrumental to species survival, little is known whether and how it evolves in natural populations, and to what extent it is affected by other environmental stressors. Here, we study physiological and molecular mechanisms of thermal tolerance over evolutionary times in multifarious environments. Using the practice of “resurrection ecology” on the keystone grazer *Daphnia magna*, we quantified genetic and plastic differences in physiological and molecular traits linked to thermal tolerance in historical and modern genotypes of the same population. This population experienced an increase in average temperature and occurrence of heat waves, in addition to dramatic changes in water chemistry, over five decades. On genotypes resurrected across the five decades, we measured plastic and genetic differences in CT_{max} , body size, Hb content and differential expression of four heat shock proteins after exposure to temperature as single stress and in combination with food levels and insecticide loads. We observed evolution of the critical thermal maximum and plastic response in body size, HSP expression and Hb content over time in a warming only scenario. Molecular and physiological responses to extreme temperature in multifarious environments were not predictable from the response to warming alone. Underestimating the effect of multiple stressors on thermal tolerance can lead to wrong estimates of species evolvability and persistence.

KEYWORDS

body size, CT_{max} , *Daphnia magna*, global change, haemoglobin, heat shock proteins

1 | INTRODUCTION

Twenty-five years ago, the Convention on Biological Diversity predicted that over 30% of multicellular species would become extinct by 2,100, largely because of human activities. These predictions correspond to more than 6,000 species going extinct and many more being impacted by altered synchrony with food and/or habitat requirements (Bellard, Bertelsmeier, Leadley, Thuiller, &

Courchamp, 2012; Easterling et al., 2000; Parmesan, Root, & Willig, 2000). Current evidence shows that species extinction rates are many times greater than the direst predictions made two decades ago by environmentalists (Hallmann et al., 2017; Isbell et al., 2011). One of the major concerns associated with changing climate is the recurrence of extreme events, such as heat waves (Easterling et al., 2000; Parmesan et al., 2000). These events, co-occurring with average temperature increase, are predicted to be more severe in Europe

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Molecular Ecology* Published by John Wiley & Sons Ltd

and North America due to atmospheric circulation patterns intensified by greenhouse gases (Meehl & Tebaldi, 2004), and to become more intense, longer lasting and/or more frequent (Karl & Trenberth, 2003).

Physiological and/or molecular plasticity has been shown to play an important role in organismal response to extreme temperatures. A typical physiological short-term plastic response to warming is reduction in body size (Atkinson, 1994; Daufresne, Lengfellner, & Sommer, 2009), a fundamental biological characteristic linked to important ecological properties such as fecundity, population growth rate and competitive abilities (Gianuca, Pantel, & De Meester, 2016; Millien et al., 2006). Short-term responses to thermal stress can also be mediated by increase in the critical thermal maximum (CT_{max}), the upper temperature at which animals lose motor function (Angilletta, 2009). Higher CT_{max} has been shown to evolve with temperature at different altitudes (Garcia-Robledo, Kuprewicz, Staines, Erwin, & Kress, 2016; Oyen, Giri, & Dillon, 2016) and over evolutionary time in response to warmer climates in some species (Daufresne et al., 2009; Geerts et al., 2015; Jansen et al., 2017; Kellermann et al., 2012) but to be phylogenetically constrained in others (Kellermann et al., 2012). Because of the link between temperature and oxygen solubility, hypoxia or low oxygen availability is generally associated with high temperature (Kobayashi, Fujiki, & Suzuki, 1988; Lamkemeyer, Zeis, & Paul, 2003). Commonly linked to high oxygen affinity in ectotherms is the regulation of the haemoglobin gene family (Hb), which guarantees oxygen supply to tissues (Gerke, Börding, Zeis, & Paul, 2011). Another typical response to sudden and extreme stress, including temperature stress, is the regulation of heat shock proteins (HSP) (Sorensen, Kristensen, & Loeschcke, 2003). HSPs function as molecular chaperones protecting cells against accumulation of damaged proteins (Sorensen et al., 2003) and playing a vital role in stress tolerance and survival under adverse conditions (Mayer & Bukau, 2005). The HSP20 protein family plays an important role in modulating cellular defence under environmental stress conditions (Seo, Lee, Park, & Lee, 2006); the HSP60 family controls the modification of newly synthesized proteins, repairs damaged ones and prevents peptides from accumulating (Pockley, 2003); the HSP70 family is responsible for cellular homeostasis under nonstress conditions and plays a vital role in stress tolerance and survival under adverse conditions (Mayer & Bukau, 2005; Schumpert, Handy, Dudycha, & Patel, 2014); the HSP90 family plays a major role in stress tolerance by removing proteins with incorrect structure and by mediating proper folding under stress conditions (Schnaider, Somogyi, Csermely, & Szamel, 2000). Both plastic responses in the expression of HSP proteins (Jansen et al., 2017; Mikulski, Bernatowicz, Grzesiuk, Kloc, & Pijanowska, 2011; Mikulski, Grzesiuk, Kloc, & Pijanowska, 2009) and evolution of HSPs over microevolutionary time scales have been associated with thermal stress (Bettencourt, Feder, & Cavicchi, 1999; Ketola, Laakso, Kaitala, & Airaksinen, 2004; Riehle, Bennett, Lenski, & Long, 2003). Yet, it is unclear how plasticity impacts long-term evolutionary responses to thermal stress as plasticity can either help (Ghalambor, McKay, Carroll, & Reznick, 2007; Mitchell, Sgro, & Hoffmann, 2011) or hinder (Hendry, 2016) evolutionary responses.

Here, we study molecular and physiological mechanisms of thermal tolerance in an invertebrate ectotherm common to European lotic freshwater ecosystems, *Daphnia magna* (Miner, De Meester, Pfrender, Lampert, & Hairston, 2012). *Daphnia* spp are keystone species in lakes and ponds worldwide, and drivers of ecosystem dynamics (Altshuler et al., 2011; Miner et al., 2012). In the natural environment, *D. magna* is exposed to severe spatial and temporal environmental changes, including temperature, low oxygen and eutrophication, to which it responds via an ecoresponsive genome (Colbourne et al., 2011; Orsini, Gilbert et al., 2016 Orsini et al., 2018). *Daphnia magna* has a parthenogenetic life cycle that allows the rearing of populations of genetically identical individuals (clones) from a single genotype. In unfavourable environmental conditions, *D. magna* switches from asexual to sexual reproduction, producing dormant embryos that arrest their development entering dormancy (Ebert, 2005). These dormant embryos sink into lake sediment and build up archives of living fossils that remain viable for decades or even centuries (Cousyn et al., 2001; Decaestecker et al., 2007; Frisch et al., 2014; Orsini, Spanier, & De Meester, 2012; Orsini, Marshall, et al., 2016). With the practice of “resurrection ecology” (Kerfoot & Weider, 2004), the dormant embryos can be resuscitated and historical and modern populations used in the same experimental settings. These properties provide us with the unique opportunity to study mechanisms of response to environmental changes through evolutionary time and to disentangle the relative contribution of plastic and genetic response to environmental change.

Previous studies have shown adaptive response of *D. magna* to temperature changes via physiological and molecular mechanisms (Cambroner, Zeis, & Orsini, 2017; Geerts et al., 2015; Jansen et al., 2017). Geerts and coworkers provided evidence of evolution of the temperature of maximum tolerance (CT_{max}) across few decades in response to warmer climates (Geerts et al., 2015). In a follow-up study, Jansen et al. showed that evolution in CT_{max} was mediated by both plastic and evolutionary changes in gene expression at a number of candidate genes, including heat shock proteins (Jansen et al., 2017). Cambroner et al. provided evidence that haemoglobin (Hb)-rich *D. magna* genotypes have superior competitive abilities as compared to Hb-poor genotypes under hyperthermal stress, showing that Hb has an important role in thermal stress response (Cambroner et al., 2017).

Three populations of *D. magna* separated in time were previously sampled from a well-characterized lake in Denmark, Lake Ring, which experienced an increase in average temperature and occurrence of heat waves across five decades (1960–2005) (Cuenca Cambroner & Orsini, 2018; Orsini, Marshall, et al., 2016). The lake also experienced changes in water chemistry transitioning from a severe event of eutrophication (1960–1970) to a partial recovery in modern times (>1999); the lake also suffered from increased agricultural land use between 1975 and 1985 (Cambroner et al., 2017; Orsini, Marshall, et al., 2016). In a previous study, Cambroner et al. (in review) used these three populations in common garden experiments to measure genetic and plastic response in life history traits to temperature as single stress and in combination with insecticide

loads and food levels, mimicking major stressors that occurred in the lake (Cambronero et al., in review). Here, we assessed the impact of multifarious environments on thermal tolerance by measuring the critical thermal maximum (CT_{max}) and by quantifying the differential expression of four heat shock proteins (HSP20, HSP60, HSP70 and HSP90) at the upper thermal limit on the genotypes used in the common garden experiments. Furthermore, we reanalysed previous data generated on Hb content and body size collected on the same genotypes (Cambronero et al., 2017, in review).

We expected the physiological and molecular traits to evolve over time in response to an increase in average ambient temperature and occurrence of heat waves. Our results show that coping mechanisms to thermal stress over evolutionary time in the presence of multiple environmental stressors are not predictable from the response to temperature as a single stressor.

2 | MATERIALS AND METHODS

2.1 | Sampling site and Lake Ring paleolimnological profile

The study site is Lake Ring, a well-characterized lake in Jutland, Denmark (55°57'51.83" N, 9°35'46.87" E). In 2004, a sedimentary archive was sampled from the lake and stored in dark and cold (4°C) conditions. In 2015, a radiometric chronology of this archive was completed by ENSIS Ltd (UCL London) following standard protocols (Appleby, 2001). Environmental changes in the Lake were reconstructed from the paleolimnological analysis of sediment and historical records as described in Ref. (Cambronero et al., 2017; Cuenca Cambronero & Orsini, 2018). The paleolimnological analysis consisted of quantifying the organic matter in the sediment, measured as loss on ignition (LOI) (Heiri, Lotter, & Lemcke, 2001), and the invertebrate community (*Cladocera*) assemblage over time. Furthermore, total phosphorus, nitrogen and water transparency were measured in Lake Ring by the County of Vejle in the Jutland peninsula for the years 1971–1999 as part of a monitoring programme following standard protocols (Søndergaard et al., 1990). Temperature records were collected by the Danish Meteorological Institute at a weather station located 80 km from the Lake over the past century (<http://www.dmi.dk/laer-om/generelt/dmi-publikationer/2013/>). Because air and water surface temperature have a positive correlation for shallow streams and lakes (Preudhomme & Stefan, 1992), especially for the summer months (e.g., (Livingstone & Lotter, 1998), the data from the weather station were previously used as estimates of the monthly water temperature in the lake (Cambronero et al., in review). Historical records of pesticides used in Denmark were collected from the Danish national archives (www.middeldatabasen.dk). According to these archives, carbamate pesticides (e.g., Carbaryl) were commonly used in the 1980s. The paleolimnological and the historical data revealed that temperature, high primary production and Carbaryl levels were key environmental factors impacting Lake Ring in the five decades analysed (Cambronero et al., in

review; Michels, 2007). Based on the occurrence of these stressors over time, three main phases were identified as follows: (a) eutrophication phase (EP, 1960–1970) characterized by high primary production (high LOI) due to sewage inflow; (b) pesticides phase (PP, 1975–1985) driven by increase in agricultural land use. This phase is associated with heavy usage of carbamate insecticides among others; and (c) clear water or recovery phase (CWP, >1999), associated with a decrease in primary production, phosphorus and nitrogen, and increase in water transparency. This phase coincided with the diversion of sewage inflow and reduction in agricultural land use in modern times.

From each lake phase, Cambronero et al. (in review) resurrected *D. magna* populations separated in time following established protocols (Cuenca Cambronero & Orsini, 2018). Among the total hatched embryos from the sediment core ($N = 262$), 10 random genotypes from each lake phase ($N = 30$) were used in common garden experiments to assess genetic and plastic response in life history traits (size at maturity, age at maturity, fecundity and mortality) to temperature as single stress and in combination with loads of the insecticide Carbaryl or food levels. Prior to starting the CGEs, the genotypes were acclimated and synchronized for two generations in common garden conditions (16:8 light: dark regime, $16 \pm 1^\circ\text{C}$ and fed ad libitum 0.8 mg Carbon/L of *C. vulgaris* daily) to reduce interference from maternal effect. After two generations in these conditions, 24–48 hr old juveniles from the second or following broods were randomly assigned to the experimental conditions in which life history traits were measured in the time spanning an individual life cycle (until release of the second brood). In CGE1, the populations were exposed to $24 \pm 1^\circ\text{C}$ and $18 \pm 1^\circ\text{C}$; the experimental animals were fed ad libitum with *Chlorella vulgaris* (0.8 mg C/L). In CGE2, the two experimental temperatures ($18 \pm 1^\circ\text{C}$ and $24 \pm 1^\circ\text{C}$) were crossed with two nutrient levels: 0.2 mg C/L and 2.4 mg C/L. In CGE3, the two experimental temperatures ($18 \pm 1^\circ\text{C}$ and $24 \pm 1^\circ\text{C}$) were crossed with two concentrations of the insecticide Carbaryl: 4 $\mu\text{g/L}$ and 10 $\mu\text{g/L}$; the animals were fed ad libitum with *C. vulgaris*. Concentrations of food and insecticide imposing sublethal effects on *D. magna* life history traits were estimated in a pilot experiment prior to the common garden experiments (Cambronero et al., in review).

The genotypes are an unbiased representation of the local population genetic diversity as hatching success fluctuated along the sedimentary archive but did not systematically decrease with the age of the sediment (Cuenca Cambronero & Orsini, 2018). Moreover, previous results on the genetic composition of *D. magna* in Lake Ring showed that genetic drift and selection did not have a detectable impact on the neutral genetic diversity over time, measured both on the hatched and unhatched local population of *D. magna* (Orsini, Marshall, et al., 2016). Negligible impact of drift and selection on neutral genetic diversity in the presence of strong environmental selection provide an ideal system to study evolutionary responses in physiological and molecular traits. The sample size per population was chosen based on previous results showing that 10 genotypes are representative of the genetic diversity of local *D. magna* populations (Orsini, Marshall, et al., 2016).

2.2 | Physiological and molecular response to thermal stress

To assess the impact of multifarious environments on *D. magna* thermal tolerance, we studied physiological and molecular responses of the 30 genotypes resurrected from Lake Ring after exposure to temperature as single stress and in combination with either biotic stress (two food levels) or abiotic stress (two concentrations of the insecticide Carbaryl) in common garden experiments. At completion of the common garden experiments described above, we measured the critical thermal maximum (CT_{max}) on the 30 genotypes across all exposures. Furthermore, we quantified the expression of four heat shock proteins (HSP20, HSP60, HSP70 and HSP90) at the upper thermal limit. Differential expression was measured between genotypes at their upper thermal limit and the same genotypes reared in nonstress conditions. HSP expression was measured on a subset of the genotypes from the common garden experiments exposed to temperature as a single stress and in combination with food limitation or one insecticide concentration. These specific conditions were previously identified as imposing severe sublethal effects on life history traits (Cambronero et al., in review). Methods for measuring CT_{max} and HSP expression are described below. In addition to data on CT_{max} and HSP expression, we reanalysed data on body size from Cambronero et al. (in review) and data on haemoglobin induction available on the same genotypes (Cambronero et al., 2017).

2.2.1 | Critical thermal maximum

Critical thermal maximum was measured on the experimental animals from the common garden experiments conducted by Cambronero et al. (in review) (Table S1). These included exposure to temperature as single stress and in combination with biotic or abiotic stress. Following the protocol optimized by Geerts et al. (2015), we exposed individual adult *Daphnia* after they released their second brood to temperature ramping of 1°C increments every 20 ± 5 seconds until the animals stopped swimming (Geerts et al., 2015). Ten genotypes (biological replicates) per population and treatment, in a single clonal replicate, were used. The temperature ramping essay reflects a rapid increase in temperature associated with heat waves. In shallow lakes and ponds, air and water temperature have a positive correlation (Livingstone & Lotter, 1998; Schneider & Hook, 2010); hence, the simulated rapid increase in temperature is a realistic approximation of severe heat waves occurring in the natural environment.

The individuals exposed to temperature ramping were placed in Eppendorf tubes (1.5 ml) within their own medium. The temperature ramping, starting at 16°C, was performed in a thermal block (Eppendorf ThermoMixer C) where the swimming activity was continuously monitored by an operator, who recorded the temperature of ceased swimming activity using a voice recording device. The temperature was monitored both on the display of the thermal block and with a thermometer. A second operator assisted by flash freezing animals in liquid nitrogen immediately after they stopped swimming. On these animals, we quantified the expression of four heat

shock proteins using as reference the same genotypes not exposed to temperature ramping.

2.2.2 | Heat shock proteins

Differential expression (DE) of four heat shock proteins [HSP20, HSP60, HSP70 and HSP90, (Jansen et al., 2017)] was quantified in a subset of genotypes from the common garden experiments. HSPs differential expression was measured in three genotypes (biological replicates) per population, in three technical replicates. These genotypes were from the common garden experiments assessing the impact of temperature as single stress and in combination with food limitation (0.2 mg C/L) and one concentration of Carbaryl (4µg/L) ($N = 1,440$, Table S2).

Total RNA was extracted from single individuals using Agencourt® RNAdvance™ Tissue kit, following the manufacturer's instructions. RNA was quantified using Nanodrop® Technologies (USA) and cDNA synthesis was performed using the AffinityScript cDNA synthesis kit, following the manufacturer's instructions. qPCR was performed using a Roche LC96 qPCR machine, following the SYBR® Premix Ex Taq (Tli, RNaseH Plus, Takara) protocol in 20 µL final volume. The cycling was as follows: initial denaturation for 1 min at 95°C, followed by 40 cycles consisting of 5" at 95°C, 30" at 60°C and 1" minute at 72°C. A final extension of 5 min at 72°C was used. We calculated the mean CT (cycle threshold) value per sample averaging among replicates before rescaling the value. CT values per sample and per protein were rescaled using an interplate calibrator consisting of a pool of RNA extracted from different genotypes at different developmental stages and including genotypes from the three populations studied here following (Jansen et al., 2017).

2.2.3 | Mechanisms of heat tolerance

We performed an analysis of variance on the physiological (CT_{max} and body size) and molecular traits (HSP and Hb) to assess evolutionary mechanisms—plastic, genetic or a combination thereof—underpinning thermal tolerance. Haemoglobin content of each genotype from the three populations separated in time was quantified in a previous study at $20 \pm 1^\circ\text{C}$ and $30 \pm 1^\circ\text{C}$ (Cambronero et al., 2017). The body size of adult *Daphnia* was quantified in the common garden experiments described above after exposure to temperature as single stressors and in combination with biotic and abiotic stress (Cambronero et al., in review)

Results of the variance analysis are interpreted as follows: a) a significant population term indicates genetic differences among populations in the molecular or physiological trait; b) a significant response to treatment(s) indicates plasticity in the trait; c) a significant interaction between the population and the treatment(s) indicates evolution of plasticity in the trait. We analysed physiological traits via ANOVAs using linear mixed models (LMMs) in R v.3.3.3 (Team RC, 2017) and including genotypes as a random effect. We visualized the main effects of population and treatment(s) as well as population \times treatment interactions on CT_{max} and body size through

univariate reaction norms, which describe the pattern of phenotypic expression of each genotype across treatments (Roff, 1997).

We analysed the expression of HSP proteins at the upper thermal limit as compared to nonstressful conditions via multivariate statistics (MANOVA) followed by a univariate analysis per single protein (ANOVA). Both analyses were performed using linear mixed models (LMMs) in R v.3.3.3 (Team 2017). For these analyses, we included a random error structure in each model to account for genotype-specific differences in gene expression within each population. We fitted one model per gene. In this analysis, the term "population" represents the constitutive differences in gene expression among populations. The term "temperature ramping (TR)" assessed whether the single proteins (ANOVA) or the four proteins at once (MANOVA) displayed a significant change in expression in response to the treatment. This term measures the short-term plastic response of gene expression when genotypes are exposed to sudden and severe temperature increase. The interaction term "TR" \times "population" quantifies the difference in gene expression among populations due to the treatment, and it reflects the evolution of plasticity in the expression of individual proteins or all HSPs considered at once.

We visualized the multivariate analysis results using phenotypic trajectories (PTA) (Collyer & Adams, 2007). For each population, we plotted (a) the magnitude of phenotypic change across the HSPs between TR treated and nontreated populations; (b) and the direction of change in the phenotypic trajectory (Collyer & Adams, 2007; Dennis, Carter, Hentley, & Beckerman, 2011). Significant differences ($p < 0.05$) in magnitude and direction of phenotypic change between population pairs were derived from a residual randomization procedure following (Collyer & Adams, 2007). Scripts used for this analysis are in Ref. (Adams & Collyer, 2009). Individual HSPs differential expression was visualized through univariate reaction norms.

Population differential expression in Hb and supporting statistics are in Ref. (Cambroner et al., 2017). Here, we visualize these data as univariate reaction norms.

3 | RESULTS

3.1 | Physiological responses: body size and CT_{max}

3.1.1 | CGE1 (temperature)

The effect of temperature treatment on CT_{max} was significantly different among populations, whereas body size did not vary significantly among populations (Table 1—CGE1, Figure 1). Temperature treatment induced significant plastic response in CT_{max} and body size in the three populations. Neither CT_{max} nor body size showed significant evolution of plasticity over the five decades examined (Table 1—CGE1).

The univariate reaction norms, supporting the ANOVA analysis, showed that CT_{max} was constitutively higher in the most recent populations (CWP) than in the other two populations; the three populations showed a higher CT_{max} at 24°C than at 18°C (Figure 1—CGE1). Body size was smaller at 24°C than at 18°C in all populations, and

TABLE 1 Analysis of variance ANOVA for CT_{max} and body size on the three populations of *Daphnia magna* after exposure in common garden experiments to temperature (CGE1), temperature and food levels (CGE2) and temperature and insecticide loads (CGE3). The population (*P*) term indicates genetic differences among populations in CT_{max} and body size; the terms temperature (*T*), food (*F*) and insecticide (*I*) indicate plastic responses of CT_{max} and body size to these treatment(s); the interaction terms indicate evolution of plasticity. Significant *p*-values ($p < 0.05$) are shown in bold

| | df | CT_{max} (°C) | | Body size (mm) | |
|--|----|-----------------|------------------|----------------|------------------|
| | | F | p | F | p |
| CGE1 | | | | | |
| Population (<i>P</i>) | 2 | 3.25 | 0.05 | 0.02 | 0.98 |
| Temperature (<i>T</i>) | 1 | 12.86 | <0.001 | 11.14 | 0.001 |
| <i>P</i> \times <i>T</i> | 2 | 0.43 | 0.65 | 0.66 | 0.52 |
| CGE2 | | | | | |
| Population (<i>P</i>) | 2 | 2.96 | 0.06 | 4.82 | 0.01 |
| Temperature (<i>T</i>) | 1 | 16.80 | <0.001 | 7.63 | 0.01 |
| Food (<i>F</i>) | 1 | 36.72 | <0.001 | 793.14 | <0.001 |
| <i>T</i> \times <i>F</i> | 2 | 5.87 | 0.02 | 1.31 | 0.26 |
| <i>P</i> \times <i>T</i> | 2 | 6.88 | 0.001 | 0.02 | 0.98 |
| <i>P</i> \times <i>F</i> | 1 | 4.67 | 0.01 | 0.31 | 0.74 |
| <i>F</i> \times <i>T</i> \times <i>P</i> | 2 | 0.86 | 0.43 | 0.03 | 0.98 |
| CGE3 | | | | | |
| Population (<i>P</i>) | 2 | 0.81 | 0.45 | 0.35 | 0.71 |
| Temperature (<i>T</i>) | 1 | 18.40 | <0.001 | 38.34 | <0.001 |
| Insecticide (<i>I</i>) | 1 | 0.26 | 0.61 | 2.93 | 0.09 |
| <i>T</i> \times <i>I</i> | 2 | 0.12 | 0.74 | 4.92 | 0.03 |
| <i>P</i> \times <i>T</i> | 2 | 2.41 | 0.10 | 0.76 | 0.47 |
| <i>P</i> \times <i>I</i> | 1 | 9.28 | <0.001 | 0.28 | 0.76 |
| <i>I</i> \times <i>T</i> \times <i>P</i> | 1 | 2.09 | 0.15 | 1.88 | 0.18 |

the plastic response was comparable across the three populations (Figure 1—CGE1).

3.1.2 | CGE2 (temperature and food)

The effect of temperature combined with food levels varied significantly among populations for body size but not for CT_{max} (Table 1—CGE2). We observed a significant plastic response of both traits to temperature and to food levels as single stressors (Table 1—CGE2, *T* and *F*). However, a significant interaction between temperature and food (*T* \times *F*) was observed only for CT_{max} , indicative of additive effect of these two factors (Table 1—CGE2). Evolution of plasticity was observed in CT_{max} with significant interaction terms "temperature \times population" and "food \times population" (Table 2—CGE2; *P* \times *T* and *P* \times *F*).

The univariate reaction norms showed lower CT_{max} in low food levels than in high food levels and a different plastic response of the populations within food levels (Figure 1—CGE2). In high food levels,




FIGURE 1 Reaction norms of physiological responses Population reaction norms, based on population means and SD ($n = 10$), showing CT_{max} and body size changes after exposure in common garden experiments to temperature (CGE1), temperature combined with food levels (CGE2) and temperature combined with insecticide Carbaryl levels (CGE3). The populations are colour coded as follows: eutrophic population (EP, 1960–1970) in blue; pesticides population (PP, 1975–1985) in green; and clear water phase population (CWP, >1999) in red. The reaction norms for high Carbaryl are shown for two populations only as the PP population experienced 100% mortality at 18°C in high Carbaryl. High food and high Carbaryl are represented by circle symbols and dotted lines; low food and low Carbaryl are represented by triangle symbols and solid lines

CT_{max} was higher at 24°C than at 18°C for CWP and EP, whereas it remained unchanged in PP (Figure 1—CGE2; dotted lines). In low food levels, CT_{max} was higher at 24°C only in the CWP population, whereas it was lower in the other two populations (Figure 1—CGE2; solid lines). Body size remained largely unaffected by temperature within the same food regime, whereas it significantly differed between food levels (Figure 1—CGE2).

3.1.3 | CGE3 (temperature and pesticides)

The effect of temperature combined with the insecticide Carbaryl did not differ among populations in both physiological traits (Table 1—CGE3). We observed a significant plastic response of both traits to temperature but not to insecticide as single stressors; the interaction between temperature and insecticide significantly affected body size (Table 1—CGE3). Significant evolution of plasticity was observed in CT_{max} (Table 1—CGE3; $P \times I$).

The reaction norms showed comparable plastic responses of CT_{max} at both insecticide levels in the three populations (Figure 1—CGE3). The temperature of maximum tolerance was generally higher at 24°C than at 18°C, except for the PP population in low Carbaryl and the EP population in high Carbaryl (Figure 3—CGE3). Plastic responses in body size differed between high and low Carbaryl (Figure 1—CGE3). Specifically, body size was larger at 24°C than at 18°C in the presence of low Carbaryl in all three populations, whereas plastic response in body size differed among populations in high Carbaryl (Figure 1—CGE3, solid lines). In the presence of high Carbaryl, the PP population went extinct at 18°C. Therefore, we were unable to visualize the reaction norm for this population.

3.2 | Molecular response: HSPs and Hb

In the following, we present results on the HSPs multivariate analysis. In this analysis, the differential expression of the four HSPs is shown at 18°C and 24°C, and measured between genotypes exposed to the temperature ramping treatment (increase of 1°C every 20 ± 5 until swimming stops) and the same genotypes not exposed to this treatment (control).

3.2.1 | Warming

The effect of temperature ramping treatment (TR) on the HSPs did not vary significantly among populations both at 18°C and 24°C (Table 2—Warming). We observed a significant plastic response of the HSPs in the three populations of *D. magna* to TR treatment at both temperatures (Table 2—Warming). We did not observe significant evolution of plasticity under a warming only scenario (Table 2—Warming).

The magnitude and direction of change in the phenotypic trajectories measured across the HSPs in response to TR treatment were similar among populations at 18°C (Figure 2—Warming; Table S3). Conversely, at 24°C, the direction of change significantly differed between the clear water (CWP) and the eutrophic (EP) population, whereas the magnitude of phenotypic change was comparable among populations (Figure 2—Warming, 24°C; Table S3).

The univariate statistics (ANOVA) confirmed that HSPs differential expression did not vary significantly among populations at 18°C, except for HSP20, and that the expression of HSP60, HSP70 and HSP90 differed significantly among populations at 24°C

TABLE 2 Multivariate analysis of variance MANOVA showing multivariate response of four heat shock proteins to the temperature ramping treatment (TR). The TR treatment was performed after exposure of the genotypes to warming, temperature combined with food limitation (Warming and food), and temperature combined with one concentration of the insecticide Carbaryl (Warming and insecticide) in common garden experiments. The population (P) term indicates genetic differences in the overall HSP expression among populations; the TR term indicates a plastic response of the HSPs to temperature ramping treatment; the interaction term between P and TR indicates evolution of plasticity. Significant *p*-values (*p* < 0.05) are shown in bold

| | Warming | | | Warming and food limitation | | | | | | Warming and insecticide | | | | | | | | | | |
|----------------|---------|-------|----------|-----------------------------|-------|----------|------|-------|----------|-------------------------|--------|----------|-------|--------|----------|------|-------|----------|------|--------|
| | 18°C | | | 24°C | | | 18°C | | | 24°C | | | 18°C | | | 24°C | | | | |
| | df | F | <i>p</i> | df | F | <i>p</i> | df | F | <i>p</i> | df | F | <i>p</i> | df | F | <i>p</i> | df | F | <i>p</i> | | |
| Population (P) | 2 | 1.46 | 0.19 | 2 | 1.95 | 0.07 | 2 | 3.67 | 0.001 | 1.99 | 0.07 | 2 | 1.77 | 0.10 | 1.43 | 2 | 1.77 | 0.10 | 1.43 | 0.20 |
| TR | 1 | 17.94 | <0.001 | 1 | 11.90 | <0.001 | 1 | 38.28 | <0.001 | 6.14 | <0.001 | 1 | 26.21 | <0.001 | 9.24 | 1 | 26.21 | <0.001 | 9.24 | <0.001 |
| P × TR | 2 | 1.56 | 0.16 | 2 | 1.88 | 0.08 | 2 | 3.33 | 0.001 | 1.39 | 0.22 | 2 | 2.26 | 0.04 | 2.28 | 2 | 2.26 | 0.04 | 2.28 | 0.04 |

(Figure 3—Warming, Table S4). Significant plastic response of the HSPs to TR treatment observed in the MANOVA was confirmed by the ANOVAs on individual proteins. We observed significant upregulation of HSP20, HSP60 and HSP70 at 18°C, and of HSP60, HSP70 and HSP90 at 24°C (Figure 3—Warming, Table S4). The univariate analysis also identified significant interaction between “TR” and “population” in HSP70 at 24°C (Figure 3—Warming, Table S4).

3.2.2 | Warming and food limitation

The effect of TR treatment on the HSPs varied significantly among populations at 18°C, but not at 24°C (Table 2—Warming and food limitation). We observed a significant plastic response of the HSPs to TR treatment at both temperatures (Table 2—Warming and food limitation). We observed significant evolution of plasticity in the HSPs at 18°C (Table 2—Warming and food limitation).

At 18°C, the magnitude of change in the phenotypic trajectory of the HSPs in response to TR treatment did not significantly differ among populations. Conversely, the direction of change significantly differed in the pairwise comparisons involving the historical population and the two most recent populations (Figure 2—Warming and food limitation; Table S3). At 24°C, the magnitude of change of the phenotypic trajectory significantly differed between the eutrophic population (EP) and the pesticide population (PP) (Figure 2 Warming and food limitation; Table S3). In this treatment, the direction of the phenotypic trajectory did not significantly differ among populations (Figure 2 Warming and food limitation; Table S3).

The univariate statistics confirmed a significant constitutive difference among populations in the expression of three HSPs at 18°C (HSP60, HSP70 and HSP90) and at 24°C (HSP20, HSP60 and HSP70) (Figure 3—Warming and food limitation; Table S4). Significant plastic responses identified in the MANOVA were overall confirmed by the ANOVAs on individual proteins; however, whereas all proteins showed a significant downregulation at 18°C, only two of the four proteins (HSP60 and HSP90) were significantly differentially expressed at 24°C (Figure 3—Warming and food limitation; Table S4). A significant interaction term between “TR” and “population” was observed for all HSPs at 18°C and for HSP70 at 24°C (Figure 3—Warming and food limitation; Table S4).

3.2.3 | Warming and insecticide

The effect of TR treatment on the HSPs did not vary significantly among populations at both 18°C and 24°C (Table 2—Warming and insecticide). We observed a significant plastic response of the HSPs to TR treatment and evolution of plasticity in both temperatures (Table 2—Warming and insecticide).

At 18°C, the magnitude of change in the phenotypic trajectory significantly differed between the eutrophic (EP) and the clear water population (CWP), whereas the direction of change significantly differed between the pesticide (PP) and the eutrophic population (EP) (Figure 2—Warming and insecticide; Table S3). At 24°C, there was no significant difference in the magnitude of phenotypic trajectories




FIGURE 2 Phenotypic trajectory analysis PTA on the three populations of *Daphnia magna* resurrected from Lake Ring, resulting from multivariate response of four heat shock proteins to temperature ramping treatment at 18°C and 24°C. Patterns for PC1 and PC2 are shown for three combinations of stressors to which the populations were exposed prior to the TR treatment: (a) warming; (b) warming and food limitation; (c) warming and insecticide. Full circles represent the control (animals not exposed to temperature ramping) and open circles represent the temperature ramping (TR) treatment. HSP expression is shown for each genotype and its replicates. Population centroids are connected by reaction norms (solid lines). Differences among populations, in terms of magnitude (M) and direction (θ) of plastic response, are shown for significant pairwise population comparisons. ns: nonsignificant. Populations are colour coded as in Fig. 1: EP—blue; PP—green; CWP—red. The statistics supporting the PTA are in Table S3

among populations, whereas the direction of change was significantly different between the eutrophic (EP) and the pesticide (PP) population (Figure 2—Warming and insecticide; Table S3).

The univariate statistics confirmed that the constitutive expression of individual HSPs did not vary significantly among populations at both 18°C and 24°C (Figure 3—Warming and insecticide; Table S4). Significant plastic responses identified by the MANOVA were confirmed by the univariate statistics (Table S4). We observed significant upregulation of HSP20, HSP60 and HSP70 at 18°C, and significant differential expression of HSP70 and HSP90 at 24°C (Figure 3—Warming and insecticide; Table S4). A significant interaction term, “TR” × “population,” was observed in HSP20 and HSP60 at 18°C, and in HSP60 and HSP70 at 24°C (Figure 3, Table S4).

3.2.4 | Haemoglobin

Previously, Cambroner et al. (2017) showed that the constitutive Hb protein content did not significantly vary among populations at 20°C and 30°C, and that the synthesis of haemoglobin was higher at 30°C (Cambroner et al., 2017). These results are visualized here as population reaction norms (Figure 4).

4 | DISCUSSION

Although tolerance to extreme temperatures is critical to survival and organisms' fitness, little progress has been made to elucidate whether it evolves in natural populations and by what mechanisms.

Many studies investigating heat tolerance tend to minimize confounding factors, searching for trends in relatively undisturbed systems (Anderson, Inouye, McKinney, Colautti, & Mitchell-Olds, 2012; Geerts et al., 2015; Hoffmann & Sgro, 2011) but see (Brans et al., 2017). Limiting confounding factors has obvious advantages but can lead to wrong estimates of evolvability in the wild.

We studied the evolution of thermal tolerance in populations of the waterflea *D. magna* resurrected from different time points of the same biological archive. These populations had been exposed to an average increase in ambient temperature and occurrence of heat waves, as well as to changes in water chemistry, over five decades. Capitalizing on common garden experiments previously conducted on the resurrected populations, we assessed the impact of temperature as single stressor and in combination with biotic (food) and abiotic (insecticide) stress on physiological and molecular mechanisms of thermal tolerance.

In experimental conditions mimicking warming as single stressor, the three populations separated in time showed a plastic response in CT_{max} , with a positive correlation between upper thermal limit and plasticity in response to high temperature. Moreover, the most recent population in Lake Ring (CWP) showed a constitutive higher CT_{max} than the two historical populations. Therefore, the most recent population showed both a constitutively higher thermal maximum and comparable plasticity to the other populations. This result suggests a positive correlation between constitutive and induced thermal tolerance, at least in the presence of warming as single stressor. This evolutionary response likely occurred in response to increase in average ambient temperature and occurrence of heath




FIGURE 3 Differential expression of individual HSPs Population average ($n = 3$ genotypes \times three technical replicates) and SD of the differential expression of four heat shock proteins after temperature ramping (TR) treatment at 18°C and 24°C. Expression of HSPs is measured on the same genotypes in absence of TR and after TR treatment. The TR treatment was imposed after exposure to (1) warming; (2) warming and food limitation; (3) warming and insecticide. Dotted lines indicate significant constitutive differences in gene expression among populations (population term in Table S4). Full circles indicate a significant plastic response in gene expression induced by TR treatment (TR term in Table S4). Asterisks (*) indicate significant interaction terms "population" \times "TR" ($P \times TR$ in Table S4). Populations are colour coded as in Figure 1: EP—blue; PP—green; CWP—red




FIGURE 4 Haemoglobin differential expression population reaction norms based on population means ($n = 10$) and SD are shown for overall haemoglobin content measured at control temperature ($20 \pm 1^\circ\text{C}$) and after exposure to hyperthermal stress ($30 \pm 1^\circ\text{C}$). Populations are colour coded as in Figure 1: EP—blue; PP—green; CWP—red. Statistics supporting this plot and the Hb data used to generate these plots are from (Cambronero et al., 2017)

waves in the five decades studied here, recorded by a weather station adjacent to Lake Ring and supported by climate records in Europe (Committee on Climate Change 2017). In a previous study,

the evolution of the critical thermal maximum over few decades was associated with the capacity of *D. magna* to tolerate higher temperatures (Geerts et al., 2015). Our results corroborate these previous findings and provide a further line of evidence supporting the evolution of CT_{max} in *D. magna*, at least in a warming only scenario.

The evolution of CT_{max} over time in a warming only scenario was not reflected in the evolution of higher Hb synthesis and HSP expression. Because of the direct link between temperature changes and oxygen solubility, changes in Hb content have been studied in association with thermal stress in ectotherms (Lamkemeyer et al., 2003; Paul et al., 2004; Pörtner, 2002; Verberk et al., 2016). Previous studies have shown that higher Hb content enables ectotherms to cope with hyperthermal stress after acclimation to high temperature (Cambronero et al., 2017; Lamkemeyer et al., 2003; Paul et al., 2004; Pörtner & Knust, 2007; Verberk et al., 2016), whereas the ability to cope with this stress is dampened in absence of temperature acclimation (Cambronero et al., 2017; Seidl, Pirow, & Paul, 2005; Williams, Dick, & Yampolsky, 2012). These findings, supported by the evidence presented here, suggest that haemoglobin expression and CT_{max} evolve on different time scales and/or under different constraints.

The evolution of the upper thermal limit (CT_{max}) in a warming only scenario was not reflected in the evolution of individual HSPs proteins. However, the phenotypic trajectory analysis, providing an overall response across the HSPs, showed further insights. The trajectories and magnitude of plastic change across the HSPs in response to temperature ramping treatment did not differ among populations at 18°C, whereas they were significantly different between the historical and the two most recent populations at 24°C (PTA analysis). This finding suggests that prior acclimation to high temperature leads to divergent constitutive regulation of HSPs proteins between the two modern populations and the historical population. This response may be interpreted as acclimation to higher temperatures enhancing responses to extreme temperature events. Alternatively, exposure prior to dormancy to higher average ambient temperature and occurrence of heat waves may explain the divergent trajectories (Committee on Climate Change 2017; Desai, 2009; Parmesan et al., 2000). Overall, the observed patterns in a warming only scenario hint to a link between the evolution of CT_{max} and the expression of HSPs. However, we interpret these results with caution as expression of HSP proteins may have been driven both by the experimental increase in temperature (24°C) and by the temperature ramping (TR) treatment. A genome-wide transcriptome analysis and/or a genome-wide association study (GWAS) conducted separately on heated and nonheated scenarios is required to validate this link and identify additional molecular markers underpinning CT_{max} evolution.

In natural ecosystems and especially in enclosed habitats (e.g., lakes and ponds), factors such as eutrophication or by-products of land use may play an important role in driving evolutionary responses by altering solubility of nutrient, conductivity and oxygen levels (Feuchtmayr et al., 2009). These factors may interact additively or synergistically with temperature. There is increasing evidence that multiple factors may alter trait–environment and genotype–environment interactions influencing responses to climate change (Chown et al., 2010; Karl & Trenberth, 2003); for example, CT_{max} has been shown to change with diet (Bujan & Kaspari, 2017). Here, we show that in the presence of multiple stressors, a complex interplay among plastic and evolutionary responses both at physiological and molecular level underpinned population responses to extreme temperatures. Furthermore, we show that the evolutionary advantage of the most recent population, apparent in the constitutive higher CT_{max} in the presence of warming as single stressor, was no longer evident when temperature co-occurred with food or insecticide. Moreover, the regulation of HSPs in the presence of multiple stressors showed unexpected patterns. Although exposure to extreme temperature stress, such as the temperature ramping treatment used here to mimic heat waves, may be expected to induce a shut-down of the regulatory machinery (Kristensen, Loeschcke, & Hoffmann, 2007), we observed both a downregulation and an upregulation of HSPs after the temperature ramping treatment in the presence of multiple stressors. These patterns have been previously observed in a temporal natural population and in an

experimental population of *D. magna* exposed to temperature ramping treatment (Jansen et al., 2017). Our results and the findings of Jansen et al. (2017) indicate that whereas the HSPs seem to play an important role in the response to extreme temperatures showing a high degree of plasticity, their direction of change may be both up- and downregulation. This may be the result of diverse stressors affecting HSP expression in different directions.

Overall, the direction and the magnitude of plastic changes at molecular and physiological levels in response to extreme temperatures in multi-stress environments were not predictable from populations' response to warming as single stressor. Our results contrasting physiological and molecular responses in single and multiple stress scenarios showed that the co-occurrence of other environmental stressors with temperature has the potential to affect the evolution of thermal tolerance in natural populations of *D. magna*, either physiologically or at molecular level or both, in ways not directly predictable from the response to warming alone.

Our study is pioneer in studying trade-offs between constitutive and induced thermal tolerance over evolutionary times and in investigating the effect of multiple stressors on thermal tolerance. However, it suffers from a common limitation to resurrection ecology studies. Whereas these studies provide important insights into the evolutionary processes underlying adaptation in the wild, they require large efforts, and, for this reason, suffer from low replication. These limitations require caution when extrapolating results to other species.

Our results suggest that underestimating the effect of multiple stressors on thermal tolerance can lead to wrong estimates of species evolvability and persistence to future global change. To develop more realistic predictions about the biological impacts of climate change on species persistence, interactions between the mean and variance of environmental temperature, as well as the impact of biotic and abiotic stressors on thermal tolerance, should be considered (Bozinovic, Medina, Alruiz, Cavieres, & Sabat, 2016; Nadeau, Urban, & Bridle, 2017).

ACKNOWLEDGEMENTS

This work was supported by a NERC highlights grant (NE/N016777/1). We thank Ian Sewell for technical support in the Daphnia facility at the University of Birmingham. The dating of the Lake Ring sediment was performed by Ensis Ltd, Environmental Scientific Services, Environmental Change Research Centre, University College London. The temperature data were provided by the Danish Meteorological Institute. The authors declare no conflict of interests.

DATA ACCESSIBILITY

Data associated with this study are deposited in the DRYAD database at the following entry: <https://doi.org/10.5061/dryad.3fb854n>.

AUTHOR CONTRIBUTION

M.C.C. carried out the experiments and performed data analysis with input from J.B. S.K. and J.B. generated the HSP data. L.O. conceived the study and coordinated data analysis. L.O. and M.C.C. wrote the first version of the paper; all authors contributed to the editing of later versions.

ORCID

Luisa Orsini  <http://orcid.org/0000-0002-1716-5624>

REFERENCES

- Adams, D. C., & Collyer, M. L. (2009). A general framework for the analysis of phenotypic trajectories in evolutionary studies. *Evolution*, *63*, 1143–1154. <https://doi.org/10.1111/j.1558-5646.2009.00649.x>
- Altshuler, I., Demiri, B., Xu, S., Constantin, A., Yan, N. D., & Cristescu, M. E. (2011). An integrated multi-disciplinary approach for studying multiple stressors in freshwater ecosystems: *Daphnia* as a model organism. *Integrative and Comparative Biology*, *51*, 623–633. <https://doi.org/10.1093/icb/1093>
- Anderson, J. T., Inouye, D. W., McKinney, A. M., Colautti, R. I., & Mitchell-Olds, T. (2012). Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. *Proceedings of the Royal Society B-Biological Sciences*, *279*, 3843–3852. <https://doi.org/10.1098/rspb.2012.1051>
- Angilletta, M. J. (2009). *Thermal adaptation: A theoretical and empirical synthesis*. New York: Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780198570875.001.1>
- Appleby, P. G. (2001). *Chronostratigraphic techniques in recent sediments* Kluwer Academic Publisher. Dordrecht: The Netherlands.
- Atkinson, D. (1994). Temperature and organism size: A biological law for ectotherms? *Advances in Ecological Research*, *25*, 1–58.
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., & Courchamp, F. (2012). Impacts of climate change on the future of biodiversity. *Ecology Letters*, *15*, 365–377. <https://doi.org/10.1111/j.1461-0248.2011.01736.x>
- Bettencourt, B. R., Feder, M. E., & Cavicchi, S. (1999). Experimental evolution of Hsp70 expression and thermotolerance in *Drosophila melanogaster*. *Evolution*, *53*, 484–492. <https://doi.org/10.1111/j.1558-5646.1999.tb03783.x>
- Bozinovic, F., Medina, N. R., Alruiz, J. M., Cavieres, G., & Sabat, P. (2016). Thermal tolerance and survival responses to scenarios of experimental climatic change: changing thermal variability reduces the heat and cold tolerance in a fly. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, *186*, 581–587. <https://doi.org/10.1007/s00360-016-0980-6>
- Brans, K. I., Jansen, M., Vanoverbeke, J., Tuzun, N., Stoks, R., & De Meester, L. (2017). The heat is on: Genetic adaptation to urbanization mediated by thermal tolerance and body size. *Global Change Biology*, *3*, 5218–5227. <https://doi.org/10.1111/gcb.13784>
- Bujan, J., & Kaspary, M. (2017). Nutrition modifies critical thermal maximum of a dominant canopy ant. *Journal of Insect Physiology*, *102*, 1–6. <https://doi.org/10.1016/j.jinsphys.2017.08.007>
- Cambroner, C. M., Marshall, H., & De Meester, L., Davidson, A., Beckerman, A. P., & Orsini, L. (in review) Multiple stressors in a changing world and ecological unpredictability. *Scientific Reports*.
- Cambroner, C. M., Zeis, B., & Orsini, L. (2017). Haemoglobin-mediated response to hyper-thermal stress in the keystone species *Daphnia magna*. *Evolutionary Applications*.
- Chown, S. L., Hoffmann, A. A., Kristensen, T. N., Angilletta, M. J., Stenseth, N. C., & Pertoldi, C. (2010). Adapting to climate change: a perspective from evolutionary physiology. *Climate Research*, *43*, 3–15. <https://doi.org/10.3354/cr00879>
- Colbourne, J. K., Pfrender, M. E., Gilbert, D., Thomas, W. K., Tucker, A., Oakley, T. H., ... Boore, J. L. (2011). The ecoresponsive genome of *Daphnia pulex*. *Science*, *331*, 555–561. <https://doi.org/10.1126/science.1197761>
- Collyer, M. L., & Adams, D. C. (2007). Analysis of two-state multivariate phenotypic change in ecological studies. *Ecology*, *88*, 683–692. <https://doi.org/10.1890/06-0727>
- Committee on Climate Change. (2017). *UK climate change risk assessment 2017* (ed. Johns D), London, UK: Committee on Climate Change.
- Cousyn, C., De Meester, L., Colbourne, J. K., Brendonck, L., Verchuren, D., & Volckaert, F. (2001). Rapid, local adaptation of zooplankton behavior to changes in predation pressure in the absence of neutral genetic changes. *PNAS*, *98*, 6256–6260. <https://doi.org/10.1073/pnas.111606798>
- Cuenca Cambroner, M., & Orsini, L. (2018). Resurrection of dormant *Daphnia magna*: protocol and applications. *JoVE*, (131), e56637. <https://doi.org/10.3791/56637>
- Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in aquatic ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, *106*, 12788–12793. <https://doi.org/10.1073/pnas.0902080106>
- Decaestecker, E., Gaba, S., Raeymaekers, J., Stoks, R., Van Kerckhoven, L., Ebert, D., & De Meester, L. (2007). Host-parasite Red Queen dynamics archived in pond sediment. *Nature*, *450*, 870–874. <https://doi.org/10.1038/nature06291>
- Dennis, S. R., Carter, M. J., Hentley, W. T., & Beckerman, A. P. (2011). Phenotypic convergence along a gradient of predation risk. *Proc Biol Sci*, *278*, 1687–1696. <https://doi.org/10.1098/rspb.2010.1989>
- Desai, M. M. (2009). Reverse evolution and evolutionary memory. *Nature Genetics*, *41*, 142–143. <https://doi.org/10.1038/ng0209-142>
- Easterling, D. R., Meehl, G. A., Parmesan, C., Changnon, S. A., Karl, T. R., & Mearns, L. O. (2000). Climate extremes: Observations, modeling, and impacts. *Science*, *289*, 2068–2074. <https://doi.org/10.1126/science.289.5487.2068>
- Ebert, D. (2005). *Ecology, epidemiology, and evolution of parasitism in Daphnia*. Bethesda, MD: National Library of Medicine (US), National Center for Biotechnology.
- Feuchtmayr, H., Moran, R., Hatton, K., Connor, L., Heyes, T., Moss, B., ... Atkinson, D. (2009). Global warming and eutrophication: effects on water chemistry and autotrophic communities in experimental hypertrophic shallow lake mesocosms. *Journal of Applied Ecology*, *46*, 713–723. <https://doi.org/10.1111/j.1365-2664.2009.01644.x>
- Frisch, D., Morton, P. K., Chowdhury, P. R., Culver, B. W., Colbourne, J. K., Weider, L. J., & Jeyasingh, P. D. (2014). A millennial-scale chronicle of evolutionary responses to cultural eutrophication in *Daphnia*. *Ecology Letters*, *17*, 360–368. <https://doi.org/10.1111/ele.12237>
- García-Robledo, C., Kuprewicz, E. K., Staines, C. L., Erwin, T. L., & Kress, W. J. (2016). Limited tolerance by insects to high temperatures across tropical elevational gradients and the implications of global warming for extinction. *Proceedings of the National Academy of Sciences of the United States of America*, *113*, 680–685. <https://doi.org/10.1073/pnas.1507681113>
- Geerts, A. N., Vanoverbeke, J., & Vanschoenwinkel, B., Van Doorslaer, W., Feuchtmayr, H., Atkinson, D., ... De Meester, L. (2015). Rapid evolution of thermal tolerance in the water flea *Daphnia*. *Nature Climate Change* *5*, 665–668. <https://doi.org/10.1038/nclimate2628>
- Gerke, P., Börding, C., Zeis, B., & Paul, R. J. (2011). Adaptive haemoglobin gene control in *Daphnia pulex* at different oxygen and temperature conditions. *Comparative Biochemistry and Physiology*, *159*, 56–65. <https://doi.org/10.1016/j.cbpa.2011.01.017>

- Ghalambor, C. K., McKay, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*, *21*, 394–407. <https://doi.org/10.1111/j.1365-2435.2007.01283.x>
- Gianuca, A. T., Pantel, J. H., & De Meester, L. (2016). Disentangling the effect of body size and phylogenetic distances on zooplankton top-down control of algae. *Proceedings of the Royal Society B*, *283*, 1–8.
- Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., ... de Kroon, H. (2017). More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE* *12*(10), e0185809.
- Heiri, O., Lotter, A. F., & Lemcke, G. (2001). Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *Journal of Paleolimnology*, *25*, 101–110. <https://doi.org/10.1023/A:1008119611481>
- Hendry, A. P. (2016). *Eco-evolutionary dynamics*. Princeton, NJ: University Press.
- Hoffmann, A. A., & Sgro, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*, 479–485. <https://doi.org/10.1038/nature09670>
- Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W. S., Reich, P. B., ... Loreau, M. (2011). High plant diversity is needed to maintain ecosystem services. *Nature*, *477*, 199–203. <https://doi.org/10.1038/nature10282>
- Jansen, M., Geerts, A. N., Rago, A., Spanier, K. I., Denis, C., De Meester, L., & Orsini, L. (2017). Thermal tolerance in the keystone species *Daphnia magna*—a candidate gene and an outlier analysis approach. *Molecular Ecology*, *26*, 2291–2305. <https://doi.org/10.1111/mec.14040>
- Karl, T. R., & Trenberth, K. E. (2003). Modern global climate change. *Science*, *302*, 1719–1723. <https://doi.org/10.1126/science.1090228>
- Kellermann, V., Overgaard, J., Hoffmann, A. A., Flojgaard, C., Svenning, J. C., & Loeschcke, V. (2012). Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences of the United States of America*, *109*, 16228–16233. <https://doi.org/10.1073/pnas.1207553109>
- Kerfoot, W. C., & Weider, L. J. (2004). Experimental paleoecology (resurrection ecology): Chasing Van Valen's Red Queen hypothesis. *Limnology and Oceanography*, *49*, 1300–1316. https://doi.org/10.4319/lo.2004.49.4_part_2.1300
- Ketola, T., Laakso, J., Kaitala, V., & Airaksinen, S. (2004). Evolution of Hsp90 expression in *Tetrahymena thermophila* (Protozoa, Ciliata) populations exposed to thermally variable environments. *Evolution*, *58*, 741–748. <https://doi.org/10.1111/j.0014-3820.2004.tb00407.x>
- Kobayashi, M., Fujiki, M., & Suzuki, T. (1988). Variation and oxygen-binding properties of *Daphnia magna* hemoglobin. *Physiological Zoology*, *61*, 415–419. <https://doi.org/10.1086/physzool.61.5.30161263>
- Kristensen, T. N., Loeschcke, V., & Hoffmann, A. A. (2007). Can artificially selected phenotypes influence a component of field fitness? Thermal selection and fly performance under thermal extremes. *Proceedings of the Royal Society B-Biological Sciences*, *274*, 771–778. <https://doi.org/10.1098/rspb.2006.0247>
- Lamkemeyer, T., Zeis, B., & Paul, R. J. (2003). Temperature acclimation influences temperature-related behaviour as well as oxygen transport physiology and biochemistry in the water flea *Daphnia magna*. *Canadian Journal of Zoology*, *81*, 237–249. <https://doi.org/10.1139/z03-001>
- Livingstone, D. M., & Lotter, A. F. (1998). The relationship between air and water temperatures in lakes of the Swiss Plateau: a case study with palaeolimnological implications. *Journal of Paleolimnology*, *19*, 181–198. <https://doi.org/10.1023/A:1007904817619>
- Mayer, M. P., & Bukau, B. (2005). Hsp70 chaperones: Cellular functions and molecular mechanism. *Cellular and Molecular Life Sciences*, *62*, 670–684. <https://doi.org/10.1007/s00018-004-4464-6>
- Meehl, G. A., & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*, *305*, 994–997. <https://doi.org/10.1126/science.1098704>
- Michels, H. (2007). *Micro-evolutionary response of *Daphnia magna* to changes in biotic stress associated with habitat degradation and restoration of a shallow lake*. Doctoral University of Leuven.
- Mikulski, A., Bernatowicz, P., Grzesiuk, M., Kloc, M., & Pijanowska, J. (2011). Differential levels of stress proteins (HSPs) in male and female *Daphnia magna* in response to thermal stress: a consequence of sex-related behavioral differences? *Journal of Chemical Ecology*, *37*, 670–676. <https://doi.org/10.1007/s10886-011-9969-5>
- Mikulski, A., Grzesiuk, M., Kloc, M., & Pijanowska, J. (2009). Heat shock proteins in *Daphnia* detected using commercial antibodies: description and responsiveness to thermal stress. *Chemoecology*, *19*, 69–72. <https://doi.org/10.1007/s00049-009-0010-1>
- Millien, V., Lyons, S. K., Olson, L., Olson, L., Smith, F. A., Wilson, A. B., & Yom-Tov, Y. (2006). Ecotypic variation in the context of global climate change: Revisiting the rules. *Ecology Letters*, *9*, 853–869. <https://doi.org/10.1111/j.1461-0248.2006.00928.x>
- Miner, B. E., De Meester, L., Pfrender, M. E., Lampert, W., & Hairston, N. G. (2012). Linking genes to communities and ecosystems: *Daphnia* as an ecogenomic model. *Proceedings of the Royal Society B-Biological Sciences*, *279*, 1873–1882. <https://doi.org/10.1098/rspb.2011.2404>
- Mitchell, K. A., Sgro, C. M., & Hoffmann, A. A. (2011). Phenotypic plasticity in upper thermal limits is weakly related to *Drosophila* species distributions. *Functional Ecology*, *25*, 661–670. <https://doi.org/10.1111/j.1365-2435.2010.01821.x>
- Nadeau, C. P., Urban, M. C., & Bridle, J. R. (2017). Climates past, present, and yet-to-come shape climate change vulnerabilities. *Trends in Ecology & Evolution*, *32*, 786–800. <https://doi.org/10.1016/j.tree.2017.07.012>
- Orsini, L., Brown, J. B., Shams Solari, O., Li, D., He, S., Podicheti, R., ... De Meester, L. (2018). Early transcriptional response pathways in *Daphnia magna* are coordinated in networks of crustacean-specific genes. *Molecular Ecology* *27*, 886–897.
- Orsini, L., Gilbert, D., Podicheti, R., Jansen, M., Brown, J. B., Solari, O. S., ... Frilander, M. J. (2016). *Daphnia magna* transcriptome by RNA-Seq across 12 environmental stressors. *Scientific Data*, *3*, 160030. <https://doi.org/10.1038/sdata.2016.30>
- Orsini, L., Marshall, H., Cambroner, M. C., Chaturvedi, A., Thomas, K. W., Pfrender, M. E., ... De Meester, L. (2016). Temporal genetic stability in natural populations of the waterflea *Daphnia magna* in response to strong selection pressure. *Molecular Ecology*, *25*, 6024–6038. <https://doi.org/10.1111/mec.13907>
- Orsini, L., Spanier, K. I., & De Meester, L. (2012). Genomic signature of natural and anthropogenic stress in wild populations of the waterflea *Daphnia magna*: validation in space, time and experimental evolution. *Molecular Ecology*, *21*, 2160–2175. <https://doi.org/10.1111/j.1365-294X.2011.05429.x>
- Oyen, K. J., Giri, S., & Dillon, M. E. (2016). Altitudinal variation in bumble bee (*Bombus*) critical thermal limits. *Journal of Thermal Biology*, *59*, 52–57. <https://doi.org/10.1016/j.jtherbio.2016.04.015>
- Parmesan, C., Root, T. L., & Willig, M. R. (2000). Impacts of extreme weather and climate on terrestrial biota. *Bulletin of the American Meteorological Society*, *81*, 443–450. [https://doi.org/10.1175/1520-0477\(2000\)081<0443:IOEWAC>2.3.CO;2](https://doi.org/10.1175/1520-0477(2000)081<0443:IOEWAC>2.3.CO;2)
- Paul, R. J., Lamkemeyer, T., Maurer, J., Pinkhaus, O., Pirow, R., Seidl, M., & Zeis, B. (2004). Thermal acclimation in the microcrustacean *Daphnia*: A survey of behavioural, physiological and biochemical mechanisms. *Journal of Thermal Biology*, *29*, 655–662. <https://doi.org/10.1016/j.jtherbio.2004.08.035>

- Pockley, A. G. (2003). Heat shock proteins as regulators of the immune response. *Lancet*, 362, 469–476. [https://doi.org/10.1016/S0140-6736\(03\)14075-5](https://doi.org/10.1016/S0140-6736(03)14075-5)
- Pörtner, H. O. (2002). Climate variations and the physiological basis of temperature dependent biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comparative Biochemistry and Physiology Part A Molecular Integrative Physiology*, 132, 739–761. [https://doi.org/10.1016/S1095-6433\(02\)00045-4](https://doi.org/10.1016/S1095-6433(02)00045-4)
- Pörtner, H. O., & Knust, R. (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science*, 315, 95–97. <https://doi.org/10.1126/science.1135471>
- Preudhomme, E. B., & Stefan, H. G. (1992). Relationship between water temperatures and air temperatures for central U.S. streams. University of Minnesota, St. Anthony Falls hydraulic Laboratory, Duluth, Minnesota.
- Riehle, M. M., Bennett, A. F., Lenski, R. E., & Long, A. D. (2003). Evolutionary changes in heat-inducible gene expression in lines of *Escherichia coli* adapted to high temperature. *Physiological Genomics*, 14, 47–58. <https://doi.org/10.1152/physiolgenomics.00034.2002>
- Roff, D. A. (1997). *Phenotypic plasticity and reaction norms*. Boston, MA: Springer. <https://doi.org/10.1007/978-1-4615-4080-9>
- Schnaider, T., Somogyi, J., Csermely, P., & Szamel, M. (2000). The Hsp90-specific inhibitor geldanamycin selectively disrupts kinase-mediated signaling events of T-lymphocyte activation. *Cell Stress and Chaperones*, 5, 52–61. [https://doi.org/10.1379/1466-1268\(2000\)005<0052:T HSIGS>2.0.CO;2](https://doi.org/10.1379/1466-1268(2000)005<0052:T HSIGS>2.0.CO;2)
- Schneider, P., & Hook, S. J. (2010). Space observations of inland water bodies show rapid surface warming since 1985. *Geophysical Research Letters*, 37, L22405.
- Schumpert, C., Handy, I., Dudyca, J. L., & Patel, R. C. (2014). Relationship between heat shock protein 70 expression and life span in *Daphnia*. *Mechanisms of Ageing and Development*, 139, 1–10. <https://doi.org/10.1016/j.mad.2014.04.001>
- Seidl, M. D., Pirow, R., & Paul, R. J. (2005). Acclimation of the microcrustacean *Daphnia magna* to warm temperatures is dependent on haemoglobin expression. *Journal of Thermal Biology*, 30, 532–544. <https://doi.org/10.1016/j.jtherbio.2005.06.004>
- Seo, J. S., Lee, Y. M., Park, H. G., & Lee, J. S. (2006). The intertidal copepod *Tigriopus japonicus* small heat shock protein 20 gene (Hsp20) enhances thermotolerance of transformed *Escherichia coli*. *Biochemical and Biophysical Research Communications*, 340, 901–908. <https://doi.org/10.1016/j.bbrc.2005.12.086>
- Søndergaard, M., Jeppesen, E., Mortensen, E., Dall, E., Kristensen, T. N., & Sortkjær, O. (1990). Phytoplankton biomass reduction after planktivorous fish reduction in a shallow, eutrophic lake: A combined effect of reduced internal P-loading and increased zooplankton grazing. *Hydrobiologia*, 200, 229–240. <https://doi.org/10.1007/BF02530342>
- Sorensen, J. G., Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and ecological role of heat shock proteins. *Ecology Letters*, 6, 1025–1037. <https://doi.org/10.1046/j.1461-0248.2003.00528.x>
- Team RC (2017) *R: A language and environment for statistical computing* (ed. Computing RFFS). Vienna, Austria: Team RC.
- Verberk, W. C. E. P., Overgaard, J., Ern, R., Bayley, M., Wang, T., Boardman, L., & Terblanche, J. S. (2016). Does oxygen limit thermal tolerance in arthropods? A critical review of current evidence. *Comparative Biochemistry and Physiology Part A Molecular Integrative Physiology*, 192, 64–78. <https://doi.org/10.1016/j.cbpa.2015.10.020>
- Williams, P., Dick, K. B., & Yampolsky, L. Y. (2012). Heat tolerance, temperature acclimation and canalization of haemoglobin expression in *Daphnia*. *Evolutionary Ecology*, 26, 591–609. <https://doi.org/10.1007/s10682-011-9506-6>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Cuenca Cambronero M, Beasley J, Kissane S, Orsini L. Evolution of thermal tolerance in multifarious environments. *Mol Ecol*. 2018;27:4529–4541. <https://doi.org/10.1111/mec.14890>