

## Effectiveness and cost effectiveness of pharmacist input at the ward level:

Dawoud, Dalia M.; Smyth, Maria; Ashe, Joanna; Strong, Thomas; Wonderling, David; Hill, Jennifer; Varia, Mihir; Dyer, Philip; Bion, Julian

DOI:

[10.1016/j.sapharm.2018.10.006](https://doi.org/10.1016/j.sapharm.2018.10.006)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Dawoud, DM, Smyth, M, Ashe, J, Strong, T, Wonderling, D, Hill, J, Varia, M, Dyer, P & Bion, J 2018, 'Effectiveness and cost effectiveness of pharmacist input at the ward level: a systematic review and meta-analysis', *Research in Social and Administrative Pharmacy*. <https://doi.org/10.1016/j.sapharm.2018.10.006>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Accepted Manuscript

Effectiveness and cost effectiveness of pharmacist input at the ward level: A systematic review and meta-analysis

Dalia M. Dawoud, Maria Smyth, Joanna Ashe, Thomas Strong, David Wonderling, Jennifer Hill, Mihir Varia, Philip Dyer, Julian Bion



PII: S1551-7411(18)30652-1

DOI: <https://doi.org/10.1016/j.sapharm.2018.10.006>

Reference: RSAP 1157

To appear in: *Research in Social & Administrative Pharmacy*

Received Date: 13 July 2018

Revised Date: 9 October 2018

Accepted Date: 12 October 2018

Please cite this article as: Dawoud DM, Smyth M, Ashe J, Strong T, Wonderling D, Hill J, Varia M, Dyer P, Bion J, Effectiveness and cost effectiveness of pharmacist input at the ward level: A systematic review and meta-analysis, *Research in Social & Administrative Pharmacy* (2018), doi: <https://doi.org/10.1016/j.sapharm.2018.10.006>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title:**

**Effectiveness and cost effectiveness of pharmacist input at the ward level: a systematic review and meta-analysis**

**Authors:**

**Dalia M Dawoud<sup>1,2\*</sup>**

<sup>1</sup> Lecturer, School of Life and Medical Sciences, University of Hertfordshire, Hatfield, United Kingdom.

<sup>2</sup> Associate Professor, Clinical Pharmacy Department, Faculty of Pharmacy, Cairo University, Cairo, Egypt.

**Maria Smyth<sup>3</sup>**

<sup>3</sup>Senior Research Fellow, National Guideline Centre, Royal College of Physicians, London, United Kingdom

**Joanna Ashe<sup>3</sup>**

<sup>3</sup>Senior Information Scientist, National Guideline Centre, Royal College of Physicians, London, United Kingdom

**Thomas Strong<sup>3</sup>**

<sup>3</sup>Technical Adviser, National Institute for Health and Care Excellence, London, United Kingdom

**David Wonderling<sup>3</sup>**

<sup>3</sup>Head of Health Economics, National Guideline Centre, Royal College of Physicians, London, United Kingdom

**Jennifer Hill<sup>3</sup>**

<sup>3</sup>Guideline Lead, National Guideline Centre, Royal College of Physicians, London, United Kingdom

**Mihir Varia<sup>4</sup>**

<sup>4</sup>Community pharmacist, NHS East and North Hertfordshire Clinical Commissioning Group, Hertfordshire, United Kingdom

**Philip Dyer<sup>5</sup>**

<sup>5</sup>Consultant, Diabetes, Endocrinology and Acute Internal Medicine, Diabetes Centre, Heart of England NHS Foundation Trust, Birmingham, United Kingdom

**Julian Bion<sup>6</sup>**

<sup>6</sup>Professor, Intensive Care Medicine, University of Birmingham; Honorary Consultant in Intensive Care medicine, Queen Elizabeth Hospital, Birmingham, United Kingdom

**\*Corresponding author:**

Dr Dalia Dawoud

**Address:**

School of Life and Medical Sciences  
University of Hertfordshire  
Health Research Building, Room 2F412  
College Lane, Hatfield, AL10 9AB  
United Kingdom

**Tel:** +44 (0) 1707 285155**Mobile:** +44 (0) 7747610292**Email:** [ddawoud@hotmail.com](mailto:ddawoud@hotmail.com)**ORCID ID:** [orcid.org/0000-0002-2105-1937](https://orcid.org/0000-0002-2105-1937)**Keywords**

Clinical pharmacy, Systematic review, Meta-analysis, Cost effectiveness, acute medicine

**Word Count****3925****Funding:**

This work was undertaken by the National Guideline Centre (NGC), Royal College of Physicians London which received funding from the National Institute for Health and Care Excellence (NICE). The views expressed in this publication are those of the authors and not necessarily those of the Institute. The funding body (NICE) did not play any direct role in the study design; the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article for publication. All researchers involved in this work were independent from the funding body at the time of completing this work. All authors, external and internal to the NGC, had full access to all of the data (including statistical reports and tables) in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Conflict of Interest:**

DD, MC, JA, TS, DW, JH, MV and JB have no interests to declare. PD reports support from Welsh Acute Physician Society Meeting 21st June 2018 <https://acutemedwales.org.uk/> during manuscript writing-up.

**ABSTRACT****Background**

Pharmacists play important role in ensuring timely care delivery at the ward level. The optimal level of pharmacist input, however, is not clearly defined.

**Objective**

To systematically review the evidence that assessed the outcomes of ward pharmacist input for people admitted with acute or emergent illness.

**Methods**

The protocol and search strategies were developed with input from clinicians. Medline, EMBASE, Centre for Reviews and Dissemination, The Cochrane Library, NHS Economic Evaluations, Health Technology Assessment and Health Economic Evaluations databases were searched.

Inclusion criteria specified the population as adults and young people (age >16 years) who are admitted to hospital with suspected or confirmed acute or emergent illness. Only randomised controlled trials (RCTs) published in English were eligible for inclusion in the effectiveness review. Economic studies were limited to full economic evaluations and comparative cost analysis. Included studies were quality-assessed. Data were extracted, summarised, and meta-analysed, where appropriate.

**Results**

Eighteen RCTs and 7 economic studies were included. The RCTs were from USA (n=3), Sweden (n=2), Belgium (n=2), China (n=2), Australia (n=2), Denmark (n=2), Northern Ireland, Norway, Canada, UK and Netherlands. The economic studies were from UK (n=2), Sweden (n=2), Belgium and Netherlands. The results showed that regular pharmacist input was most cost effective. It reduced length-of-stay (mean= -1.74 days [95% CI: -2.76, -0.72], and increased patient and/or carer satisfaction (Relative Risk (RR) =1.49 [1.09, 2.03] at discharge). At £20,000 per quality-adjusted life-year (QALY)-gained cost-effectiveness threshold, it was either cost-saving or cost-effective (Incremental Cost Effectiveness Ratio (ICER) =£632/ QALY-gained). No evidence was found for 7-day pharmacist presence.

**Conclusions**

Pharmacist inclusion in the ward multidisciplinary team improves patient safety and satisfaction and is cost-effective when regularly provided throughout the ward stay. Research is needed to determine whether the provision of 7-day service is cost-effective.

**KEY WORDS**

Clinical pharmacy, Systematic review, Meta-analysis, Cost effectiveness, acute medicine

## 1 INTRODUCTION

2 Adverse drug events (ADEs) are common in clinical settings, with a reported incidence from 2.3% in paediatric  
3 inpatients to 27.4% in adult outpatients.<sup>1 2</sup> In adult inpatients, the reported incidence is 6.5%.<sup>3</sup> These ADEs are  
4 direct causes of patient harm, dissatisfaction, prolonged hospital stay and increased costs. Pharmacists are  
5 considered the medication experts in the health care team. Their extensive training in and knowledge of  
6 pharmacology and therapeutics have placed them in the best position to undertake this role and to advise other  
7 health care professionals on matters relating to appropriate prescribing and safe use of medicines.<sup>4</sup>

8 The pharmacist role in the hospital setting has evolved over the years, moving from a wholly dispensary-based  
9 role to a more clinically-focused one based on the ward.<sup>5</sup> In fact, the presence of a ward-based pharmacist has  
10 become common practice in the UK.<sup>6</sup> More recently, pharmacists have been granted the authority to prescribe  
11 medications in a number of countries including the UK and Canada.<sup>7</sup> This has allowed clinical pharmacists who  
12 practise in hospitals to be more directly involved in patient care.

13 In the UK, medical wards have access to some level of pharmacist input; however, the pharmacist may be  
14 responsible for covering several areas concurrently, limiting the level of detail they can bring to medicines  
15 management and patient and staff communication.<sup>6</sup> This is particularly important for an ageing population with  
16 multiple co-morbidities for whom polypharmacy adds complexity and may indeed be the cause of the acute  
17 admission.<sup>8</sup> Additionally, it has been argued that the input of a ward-based pharmacist, particularly at discharge,  
18 can improve patient flow by expediting the discharge process and alleviating the pressure that the “exit block”,  
19 created by delayed discharge, can have on emergency department performance and the emergency access target  
20 achievement.<sup>9</sup>

21 In 2014, the National Institute for Health and Care Excellence (NICE) was commissioned to develop a guideline  
22 to advise the National Health Service (NHS) in England on various aspects of the delivery of emergency and  
23 acute medical care services.<sup>10</sup> One of the aspects identified as a priority to be examined in the guideline was the  
24 role of ward-based pharmacists with the aim of assessing the impact of their interventions on improving patient  
25 and process outcomes in the acute and emergency medical care pathway within NHS hospitals.

26 Hence, this systematic review was undertaken as part of the guideline development process to assess the  
27 outcomes of ward-based pharmacists’ interventions for patients admitted to hospital with a suspected or  
28 confirmed acute medical emergency.

## 29 METHODS

30 A systematic literature review was undertaken to synthesise the evidence that assessed the effectiveness and  
31 cost-effectiveness of the presence of ward-based pharmacists for patients with a suspected or confirmed acute  
32 medical emergency. It was undertaken in accordance with the standard methods for reviewing the clinical and  
33 economic evidence specified in the NICE guidelines development manual.<sup>11</sup> No ethics approval was required for  
34 this work.

### 35 Protocol development

36 The protocol for reviewing the effectiveness evidence was developed and approved by the guideline  
37 development group (GDG), a team of experts consisting of 19 health care professionals including acute care  
38 clinicians and a pharmacist in addition to two lay members and a technical team. The protocol specified the  
39 inclusion and exclusion criteria (including the population, interventions and comparators, outcomes and study  
40 design). These are briefly outlined below (Box 1).

41 The protocol for reviewing the economic evidence was aligned with this in terms of the population,  
42 interventions and comparators. Full economic evaluations (studies comparing costs and health consequences of  
43 alternative courses of action which include cost-utility, cost-effectiveness, cost-benefit and cost-consequences  
44 analyses) and comparative costing studies that addressed the review question in the relevant population were  
45 considered potentially includable as health economic evidence.

46 Exclusion criteria for the economic review included the following:

- 47 1- Economic studies that only reported cost per hospital (not per patient), or only reported average cost-  
48 effectiveness without disaggregated costs and effects.
- 49 2- Studies published before 2005, because health services change rapidly and therefore the costs and  
50 benefits of treatments soon become out of date.
- 51 3- Studies from non-OECD countries or the USA were also excluded, on the basis that the applicability of  
52 such studies to the present UK NHS context is likely to be too low for them to be helpful for decision-  
53 making.

54 Remaining health economic studies were prioritised for inclusion based on their relative applicability to the  
55 guideline context and the study limitations (see Quality Assessment below).



56 The clinical and economic review protocols are presented in Appendix 1 in the Supplementary Material.

57

ACCEPTED MANUSCRIPT

## **Box 1: Population, Intervention, Comparator, Outcomes (PICO) and inclusion/exclusion criteria of the clinical review**

### *Population*

The population of interest was defined as adults and young people (16 years and over) admitted to hospital with a suspected or confirmed acute medical emergency (AME).

### *Interventions and comparators*

The intervention was defined as “presence of medical ward-based pharmacists” and the comparator as “No ward-based pharmacists”. The intervention was further stratified as either for less than 7 days a week or for 7 days a week.

### *Outcomes*

- Mortality during the study period,
- Avoidable adverse events during the study period,
- Quality of life during the study period,
- Patient and/or carer satisfaction during the study period,
- Length of stay in hospital during the study period,
- Readmissions within 30 days, future admissions to hospital (over 30 days),
- Discharges during the study period,
- Prescribing errors during the study period,
- Missed medications during the study period,
- Medicines reconciliation during the study period,
- Staff satisfaction during the study period.

### *Inclusion and exclusion criteria*

The key population inclusion criterion was:

- Adults (18 years and over) and young people (16-17 years) who seek, or are referred for, emergency NHS care for a suspected or confirmed acute medical emergency.

The key population exclusion criteria were:

- Children
- People with acute obstetric emergencies
- People with acute mental health emergencies, once a diagnosis has been made
- People with acute surgical emergencies, once a diagnosis has been made
- People who have experienced major trauma, complex or non-complex fractures or spinal injury
- People in hospital who are not there for an acute medical emergency (i.e. elective admissions) and do not develop an acute medical emergency during their stay
- People already in hospital with acute deterioration
- People with chronic conditions who are being managed as outpatients but who require an elective admission for treatment from specialists who may be involved in the acute pathway.

Literature reviews, posters, letters, editorials, comment articles, unpublished studies and studies not in English were excluded.

**59 Information sources and search strategies**

60 Databases were searched using relevant medical subject headings, free-text terms and study-type filters where  
61 appropriate. Searches were restricted to papers published in English and were conducted in Medline, EMBASE,  
62 Centre for Reviews and Dissemination (CRD) and The Cochrane Library.

63 The economic evidence was identified by conducting a search in Medline and EMBASE, using economic filters.  
64 Searches were also conducted in the economics-specific databases NHS Economic Evaluation Database (NHS  
65 EED) and Health Technology Assessment database (HTA); which were searched via CRD.

66 Search strategies were quality assured by cross-checking reference lists of highly relevant papers, analysing  
67 search strategies in other systematic reviews, and asking the GDG members to highlight any additional studies.  
68 Searches were quality assured by a second information scientist before being run and were updated in December  
69 2016. All search strategies are listed in Appendix 2 of the Supplementary Material.

**70 Study selection**

71 The titles and abstracts of records retrieved were sifted for relevance, with potentially significant publications  
72 obtained in full text. These were assessed against the inclusion criteria (see the review protocols in Appendix 1  
73 of the supplementary materials). For the effectiveness evidence, parallel randomised controlled trials (RCTs)  
74 were included. A sample of 10% of the abstract lists was double-sifted by a second reviewer and any  
75 discrepancies were rectified.

**76 Data extraction and synthesis**

77 Data were extracted from the included studies into standard evidence tables. Meta-analyses of the efficacy data  
78 were conducted using Cochrane Review Manager (RevMan5)<sup>2</sup> software to combine the data given in all studies  
79 for each of the outcomes of interest. Fixed-effects (Mantel-Haenszel) techniques (using an inverse variance  
80 method for pooling) were used to calculate risk ratios (relative risk (RR)) for the binary outcomes, which  
81 included: mortality, admission, readmission and adverse events. The absolute risk difference was calculated  
82 using GRADEpro software,<sup>12</sup> using the median event rate in the control arm of the pooled results. For binary  
83 variables where there were zero events in either arm or a less than 1% event rate, Peto odds ratios, rather than  
84 risk ratios, were calculated.

85 Continuous outcomes were analysed using an inverse variance method for pooling weighted mean differences.  
86 These outcomes included: quality of life, length of stay in hospital (LOS), patient and/or carer satisfaction.

87 Where the studies within a single meta-analysis had different scales of measurement, standardised mean  
88 differences were used (providing all studies reported either change from baseline or final values rather than a  
89 mixture of both); each different measure in each study was 'normalised' to the standard deviation value pooled  
90 between the intervention and comparator groups in that same study.

91 Statistical heterogeneity was assessed by considering the chi-squared test for significance at  $p < 0.1$  or an I-  
92 squared ( $I^2$ ) inconsistency statistic (with an I-squared value of more than 50% indicating significant  
93 heterogeneity) as well as the distribution of effects. Where significant heterogeneity was present, predefined  
94 subgrouping of studies was carried out as per the protocols.

95

96 NICE economic evidence profile tables were used to summarise cost and cost-effectiveness estimates from the  
97 included studies. These show the incremental costs, incremental effects (for example, quality-adjusted life-years  
98 [QALYs]) and incremental cost-effectiveness ratio (ICER) for the base-case analysis in the study, as well as  
99 information about the assessment of uncertainty in the analysis. When a non-UK study was included, the results  
100 were converted into pounds sterling using the appropriate purchasing power parity.<sup>13</sup> Cost effectiveness was  
101 assessed based on a cost-effectiveness threshold of £20,000 per QALY gained; in line with the NICE reference  
102 case; where ICERs less than the specified threshold indicate cost effectiveness.<sup>11</sup>

### 103 **Quality assessment**

104 The evidence for outcomes from the included RCTs were evaluated using an adaptation of the 'Grading of  
105 Recommendations Assessment, Development and Evaluation (GRADE) toolbox' developed by the international  
106 GRADE working group (<http://www.gradeworkinggroup.org/>). The software (GRADEpro) was used to assess  
107 the quality of the evidence for each outcome, taking into account individual study quality and the meta-analysis  
108 results.<sup>12</sup> Each outcome was first examined for each of the quality elements (see Supplementary material,  
109 Appendix 3, Table 3.1 for details). Publication bias was only taken into consideration in the quality assessment  
110 if it was apparent.

111 The methodological quality of the economic evidence and its applicability to the UK context were assessed  
112 using the economic evaluation checklist from the NICE guidelines manual, and included in the economic  
113 evidence profile (see Appendix 3 in the Supplementary Material for the possible ratings for each dimension and  
114 their criteria).<sup>11</sup>

#### 115 **Patient involvement**

116 Two lay members were part of the guideline development group and contributed to the development of the  
117 review protocol. The choice of the outcome measures was informed by their views of which outcomes were  
118 critical from a patient perspective. The analysis methods and results were regularly presented to and validated  
119 by all the group members including the two lay members.

#### 120 **RESULTS**

121 The search for RCTs retrieved 3196 records. Of these, 20 papers reporting on 18 RCTs were included in the  
122 review.<sup>14-33</sup> A list of the excluded studies with reasons for exclusion are presented in Appendix 4 in the  
123 Supplementary Material. The economic search retrieved 918 records, of which 7 papers reporting on 7 studies  
124 were included.<sup>17 21 34-38</sup> The PRISMA flow diagrams of both searches are presented in Appendix 5, Figure 5.1  
125 and Figure 5.2.

126 The studies were split into 3 strata: regular ward-based pharmacist input (where the ward-based pharmacist  
127 provided interventions throughout the patient stay on the ward, which included both admission and discharge  
128 services), pharmacist input at admission, and pharmacist input at discharge. The interventions and comparators  
129 were often not well defined and there was variation across the studies in their composition.

130 The characteristics of the included RCTs and economic studies are summarised in Tables 1 and 2, respectively.

**Table 1: Characteristics of the included studies- clinical evidence**

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
<b>1.Regular ward-based pharmacist input</b>						
Claus 2014 <sup>17</sup>	Belgium	Surgical ICU admissions (n=69) within a university hospital.  <b>Inclusion</b> - over 16 years of age, length of stay greater than 48 hours.  <b>Exclusion</b> - none stated.	RCT	Pharmacist present on the ward. Duties included making active recommendations and performing patient follow-up.	Pharmacist is present on the ward but recommendations were not passed on to the primary care giver.	In-hospital mortality.
Iowa Continuity of Care Study trial: Farris 2014 (Farley 2014) <sup>19 20</sup>	USA	General medicine, family medicine, cardiology or orthopaedic admissions (n=631) within an academic tertiary care hospital.  <b>Inclusion</b> - patients with certain disease classifications: hypertension, hyperlipidaemia, heart failure, coronary artery disease, myocardial infarction, stroke, transient ischemic attack, asthma, chronic obstructive pulmonary disease or receiving oral anticoagulation.	RCT	Pharmacy case manager. Duties included medication reconciliation, ward visits and discharge service.	Nurse based medication reconciliation and discharge service.	Preventable adverse drug events in-hospital; post-discharge (90 days) hospital Readmission at 30 days; Admission at 90 days Medication appropriateness index (MAI) at discharge; 30 days; 90 days.
Gillespie 2009 <sup>21</sup>	Sweden	Patients (n=400) admitted to the 2 acute internal study wards at a University teaching hospital.	RCT	Pharmacist present on the ward. Duties included taking part in the rounding team,	No pharmacist involvement in the healthcare team at	Overall survival at 12 months, reported as hazard ratio.

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		<p><b>Inclusion</b> - 80 years of age.</p> <p><b>Exclusion</b> - previously been admitted to the study wards during the study period or had scheduled admissions.</p>		documenting medication history, and discharge counselling.	the ward level.	Admission at 12 months
Kucukarslan 2003 <sup>23</sup>	USA	<p>All patients (n=165) admitted to 1 of the 2 internal medicine study wards within a tertiary care hospital.</p> <p><b>Inclusion</b> - admitted to the internal medicine service and remained in the same patient care unit until discharge.</p> <p><b>Exclusion</b> – none given.</p>	Quasi-RCT	Pharmacist present on the ward. Duties included taking part in the rounding team, documenting medication history, and discharge counselling.	Standard care from 1 pharmacist (implication in paper that this is not ward-based).	<p>Avoidable adverse drug events until discharge.</p> <p>Length of stay in-hospital (reported as mean difference).</p> <p>Re-admission (unclear follow-up time, reported as percentage reduction).</p>
Shen 2011 <sup>29</sup>	China	<p>n=354 inpatients in 2 respiratory wards diagnosed with respiratory tract infections.</p> <p><b>Exclusion:</b> transferred from other medical departments; transferred to other medical departments for further treatment; already received antibiotics before admission; did</p>	RCT	Clinical pharmacist part of the treating team – communicated any potentially inappropriate antibiotic use (indication, choice, dosage, dosing schedule, duration, conversion) with the	Standard treatment strategies performed by the physicians and nurses without pharmacist involvement.	Length of stay.

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		not receive antibiotics during hospitalisation.		physician to discuss and make recommendations.		
Scullin 2007 <sup>28</sup>	Northern Ireland	Admitted patients (n=762) to the 4 medical study wards within 3 general hospitals.  <b>Inclusion:</b> taking at least 4 regular medication, were taking a high risk drug(s), were taking antidepressants and were 65 years old or older, had a hospital admission within the last 6 months, prescribed antibiotics on day 1 of admission.  <b>Exclusion</b> - scheduled admissions and patients admitted from private nursing homes.	RCT	Pharmacist present on the ward. Duties included admission services, in-patient monitoring, and discharge services	Traditional clinical pharmacy services (no further details given).	Admission at 12 months.  Mortality at 12 months.  Length of stay.
Spinewine 2007 <sup>30</sup>	Belgium	All eligible patients (n=186) admitted to the Geriatric Evaluation and Management (GEM) unit within a university teaching hospital.  GEM unit accepted patients over 70 years of age.	RCT	Pharmacist present on the ward. Duties included taking part in the rounding team, documenting medication history, and discharge counselling.	Usual care (no details of any clinical pharmacist involvement).	Rate of death at 1 year follow-up.  Satisfaction with information received.  Admission at 12 months. Medical appropriateness index.
Zhao 2015 &	China	n=90 patients admitted to the	RCT	Interventions by	Conventional	Avoidable adverse



Study	Country	Population	Study design	Intervention	Comparator	Outcomes
Zhao 2015B <sup>32</sup> 33		cardiology ward in a hospital.  <b>Inclusion:</b> diagnosis of CHD by physician, accepted $\geq 4$ kinds of drugs, $\geq 18$ years, primary high school education, able to complete the study, available for telephone follow up.  <b>Exclusion:</b> pregnant/lactating women, patients enrolled in other studies, severe co-morbidities, family history of psychosis, and barriers to communication.		clinical pharmacists including individual drug regimens, attending daily medical rounds, advice to physicians, education of medical staff, patient education on lifestyle changes, psychological interventions such as stress reduction, medication counselling at discharge, monthly follow up telephone calls post-discharge.	medical treatment without pharmacist participation.	events (adverse drug reactions).  Patient and/or carer satisfaction.
<b>2. Ward-based pharmacist input at admission</b>						
Aag 2014 <sup>14</sup>	Norway	Consecutively admitted patients (n=201) to the Cardiology study ward at a University hospital.  <b>Inclusion</b> - aged 18 and over.  <b>Exclusion</b> - terminal illness, isolated due to an infectious disease, unable to communicate in either Norwegian or English.	RCT	Pharmacist medication reconciliation.	Nurse medication reconciliation.	Medication discrepancies identified at admission.  Prescribing physician agreement to act upon medication discrepancies identified

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
Khalil 2016 <sup>22</sup>	Australia	n=110 adult medical patients admitted to the acute assessment and admission (AAA) unit via the ED during pharmacy operating hours (8.30am – 5pm).  <b>Exclusion:</b> not admitted to the AAA ward within 24 hours; no medications prior to admission; not a general medical patient.	RCT	Pharmacist-initiated medication reconciliation – pharmacist obtained a ‘best possible medication history’ from the patient and/or other sources, undertook admission medication reconciliation, reviewed current medications and the need for new medications in relation to the admission diagnosis, developed a medication management plan with the referring senior medical officer and charted on the electronic medication administration record	Usual care – medication orders charted by medical staff.	Prescribing errors.
Lind 2016 <sup>24</sup>	Denmark	n=448 patients arriving at the acute admission unit on weekdays 9am-4.15pm.	RCT	Clinical pharmacist intervention - obtaining medication history (using a	Standard care – on arrival, patients triaged by a nurse, then seen by a	Length of stay on the acute admission unit (defined as interval in minutes between

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		<p><b>Inclusion:</b> <math>\geq 18</math> years, taking <math>\geq 4</math> drugs daily (including over-the-counter, herbals and supplements).</p> <p><b>Exclusion:</b> terminal or intoxicated; assigned to triage level 1; referred to acute outpatient clinic; unable to give informed consent; interviewed by physician prior to giving informed consent; unexpected overnight stay.</p>		minimum of 2 sources, 1 of which was an interview with the patient and/or relatives where possible), entering prescriptions into the electronic medication module (EMM), medication reconciliation, reviewing overall medication treatment and writing a note in the electronic medical record.	physician who was responsible for obtaining medication history, reconciling and assessing medication treatment and entering prescriptions in the EMM.	arrival and discharge or transfer to a hospital ward).
Lisby 2010 <sup>25</sup>	Denmark	<p>Consecutively admitted patients (n=100) to acute internal medicine study ward within 1 regional hospital.</p> <p><b>Inclusion</b> - patients were 70 years or older.</p>	RCT	Pharmacist admission review.	Senior physician admission review.	<p>Self-experienced quality of health at 3 months.</p> <p>Length of stay in hospital.</p> <p>Admission rate at 3 months.</p> <p>Mortality.</p>
Nester 2002 <sup>26</sup>	USA	Consecutively admitted patients (n=100) to a tertiary care	Quasi-RCT	Pharmacist medication reconciliation.	Nurse medication reconciliation.	Medication discrepancies

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		referral centre.  <b>Inclusion</b> - over 18, responsive and able to speak English.  <b>Exclusion</b> - intensive care, ambulatory surgical and labour-and-delivery units.				identified at admission.
Tong 2016 <sup>31</sup>	Australia	n=881 patients admitted to the general medical unit (GMU) and emergency short stay unit (ESSU) during pharmacist working hours (7am-9pm).  <b>Exclusion:</b> medication chart written by a doctor before pharmacist review; admitted to ESSU and not reviewed by a pharmacist.	RCT	Early medication review and charting on admission involving a partnership between a pharmacist and a medical officer – pharmacist took medical history, VTE risk assessment and discussed medical and medication problems with admitting medical officer to agree a medication management plan.	Standard medication charting by medical officers of relevant teams, with subsequent medication reconciliation performed by pharmacist within 24 hours of admission.	Prescribing errors.
<b>3. Ward-based pharmacist input at discharge</b>						
Al-Rashed 2002 <sup>15</sup>	UK	n=83 patients admitted to 2 care of the elderly wards.	RCT	Pre-discharge counselling (24 hours before discharge) by	Normal hospital discharge policy – all patients, their GPs,	Readmission.

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		<b>Inclusion:</b> >65 years, prescribed 4 or more regular items, were to be discharged to their own home and had an abbreviated mental score >7/10, English as a first language, and routine clinical pharmacist assessment that they could have problems with their medicines after discharge.		the clinical pharmacist attached to that ward.	district nurses and carers received a copy of the patient's medication and information discharge summary sheet (MIDS) and patients received a medicine reminder card. Nurse went through (MIDS) with patients.	
Bladh 2011 <sup>16</sup>	Sweden	<p>Patients (n=345) admitted on weekdays to the 2 internal medicine study wards at a university hospital.</p> <p><b>Inclusion</b> - capable of assessing their HRQL and giving written informed consent.</p> <p><b>Exclusion</b> - poor Swedish language, planned discharge before intervention can be performed, transferred during their stay to other hospitals or wards not belonging to the Department of Medicine.</p>	RCT	Pharmacist discharge review	Usual care, which was received from the same group of physicians and nurses. No other details given.	EQ-5D summarised index at 6 months follow-up.
Eggink 2010 <sup>18</sup>	Netherlands	Patients (n=89) to be discharged (no criteria given) in the	RCT	Pharmacist discharge	Nurse discharge	Prescription errors identified during first

Study	Country	Population	Study design	Intervention	Comparator	Outcomes
		<p>Cardiology study ward within a teaching hospital.</p> <p><b>Inclusion</b> - patients have prescribed 5 or more medicines (from any class) at discharge.</p> <p><b>Exclusion</b> - none stated.</p>		review.	review.	outpatient follow-up.
Nickerson 2005 <sup>27</sup>	Canada	<p>n=253 patients admitted to 2 family practice units.</p> <p><b>Inclusion:</b> not discharged to another hospital, prescribed at least 1 medication at discharge, provided consent, agreement from community pharmacy, no previous study enrolment.</p> <p><b>Exclusion:</b> unable to answer study questions, unavailable for follow-up.</p>	RCT	Seamless care pharmacist at discharge including medication reconciliation, review of drug regime as part of comprehensive pharmaceutical care work-up, identification of problems and communication to community pharmacy, hospital staff and family physician, medication discharge counselling and a medication compliance chart	Standard care at discharge - discharge counselling and manual transcription of discharge notes from medical chart by nurse.	Prescriber errors-unresolved drug therapy inconsistencies and omissions.

Abbreviations: CHD: chronic heart disease; EQ-5D: EuroQol 5 Dimensions questionnaire; GP: general practitioner; RCT: randomised controlled trial,

1 **Table 2: Characteristics of the included studies- economic evidence**

Study	Country	Population	Study design	Follow-up/time horizon	Intervention 1	Intervention 2
<b>1. Regular ward-based pharmacist input</b>						
Claus 2014. <sup>17</sup>	Belgium	Critically ill patients (>16 years of age and with minimum length of ICU stay of 2 days) and in a 22-bed, surgical ICU at Ghent University Hospital, Belgium.	Within RCT analysis of individual patient level data  Cost-effectiveness analysis	Length of ICU stay	No clinical pharmacist direct involvement in patient care	A clinical pharmacist is directly involved in patient care
Ghatnekar 2013 <sup>35</sup>	Sweden	Elderly hospital inpatients at Skane University Hospital in Lund, Sweden	Decision tree model  Cost-utility analysis	3 months	Standard care (not defined)	Multidisciplinary team including clinical pharmacist undertakes systematic medication review and reconciliation from admission to discharge (the Lund Integrated Medicines Management [LIMM])
Gillespie 2009 <sup>21</sup>	Sweden	Elderly inpatients (80 years or older) admitted to 2 acute internal medicine wards at a University Hospital of Uppsala,	Within-RCT analysis  Cost-effectiveness analysis	12 months	No pharmacist involvement in the healthcare team at the ward level.	Pharmacist present on the ward.

Study	Country	Population	Study design	Follow-up/time horizon	Intervention 1	Intervention 2
		Sweden.				
Karnon 2008 <sup>36</sup>	UK	inpatients at 400 beds acute hospital (average hospital size) with around 14 wards	Decision tree model  Cost-utility analysis	5 years	No ward-based pharmacist (a pharmacist covers 2 wards of about 30 patients over a morning to provide basic level of pharmaceutical care and in the afternoons they have departmental commitments)	Ward-based senior pharmacist (grade 7/8a) attends rounds with residents, nurses, attending staff each morning; is present in the ward for consultation and assistance to nursing staff during the rest of the morning and is available on call as necessary during the rest of the day.
Klopotowska 2010 <sup>37</sup>	Netherlands	Patients in an adult surgical and medical 28-bed ICU of an academic medical centre	Before and after comparative interventional study  Cost-consequences analysis	Length of ICU stay.	Standard pharmacy services provided by the hospital pharmacy department.	Two experienced hospital pharmacists present on the ICU daily and attending multidisciplinary patient review meeting.
<b>2. Ward-based pharmacist input at admission</b>						
Fertleman 2005 <sup>34</sup>	UK	Medical patients admitted within the preceding 24 hours to a general medical ward	Before-and-after observational study  Comparative cost	3 days	Ward-based pharmacist provide pharmaceutical care for 1-2 hours at some	Senior pharmacist present on post-admission (post-take) ward rounds (PTWR)



Study	Country	Population	Study design	Follow-up/time horizon	Intervention 1	Intervention 2
		at a district general hospital (Northwick Park hospital in north-west London)	analysis		time during the day (usual care)	in addition to the usual care
<b>3. Ward-based pharmacist input at discharge</b>						
Wallerstedt 2012 <sup>38</sup>	Sweden	Elderly inpatients on 2 internal medicine wards at Sahlgrenska University Hospital, Sweden.	Within-RCT analysis (linked trial: Bladh 2011)  Cost-utility analysis	6 months	Usual care, which was received from the same group of physicians and nurses.	Clinical pharmacists delivering a composite intervention consisting of medication review including feedback to physicians on prescribing, drug treatment discussion with the patient at discharge, medication report including summary of drug treatment changes to be sent to the GP

2 Abbreviations: GP: general practitioner; ICU: intensive care unit; RCT: randomised controlled trial

3

#### 4 **Regular ward-based pharmacist input**

5 Eight RCTs (n= 2,303) evaluated the outcomes of the presence of a ward-based pharmacist providing regular  
6 input.<sup>17 19-21 23 28-30 32 33</sup> In these studies, the pharmacist in the intervention arm was involved in all stages of the  
7 patient journey from admissions to monitoring, follow-up and discharge. The evidence suggested reduced  
8 mortality (RR= 0.92 (95% CI: 0.72 to 1.16), 3 studies, very low quality), reduced preventable ADEs in hospital  
9 (RR= 0.74 (95% CI: 0.06 to 8.57), 2 studies, very low quality) and at 90 days follow up (RR= 0.77 (95% CI:  
10 0.29 to 2.05), 1 study, very low quality), reduced LOS (-1.74 days (95% CI: -2.76 to -0.72), 2 studies, moderate  
11 quality), reduced prescribing errors at discharge (- 0.02 (95% CI: -0.12 to 1.08), 2 studies, low quality) and  
12 increased patient and/or carer satisfaction at discharge (RR= 1.49 (95% CI: 1.09 to 2.03) and at one month  
13 follow-up (RR= 1.79 (95% CI: 1.38 to 2.32), 1 study, low quality). It also reduced hospital admission (RR= 0.93  
14 (95% CI: 0.83 to 1.04), 4 studies, moderate quality) and readmission (RR= 0.92 (95% CI: 0.62 to 1.37), 1 study,  
15 very low quality). However, there were increased prescribing errors (measured by medication appropriateness  
16 index) at 30 days (2.1 higher (95% CI: 0.45 to 3.75 higher), 1 study, moderate quality) and adverse drug events  
17 at 3 to 6 months post discharge (RR= 1.47 (0.26 to 8.33), 1 study, very low quality). The results are summarised  
18 in the clinical evidence profile in Table 5.3 and the Forest plots presented in Appendices 5 and 6 of the  
19 Supplementary Materials, respectively.

20 Five economic evaluations were included in this stratum.<sup>17 21 35-37</sup> These were conducted in Belgium (n=3),  
21 Netherland (n=1) and the UK (n=1). Three studies reported that the ward-based pharmacist input was dominant  
22 (more effective and less costly) compared to usual care. One cost-utility analysis (CUA) showed that the ward-  
23 based pharmacist intervention was cost-effective with an ICER of £632 per QALY-gained. One study showed  
24 that regular ward-based pharmacist input was less effective and less costly, with no clear conclusion regarding  
25 cost effectiveness given the absence of a cost-effectiveness threshold for the reported outcomes. All five studies  
26 were assessed as partially applicable with potentially serious limitations. The results are summarised in Table  
27 5.4, Appendix 5 and the quality assessment rationale in Appendix 7 in the Supplementary Material.

#### 28 **Ward-based pharmacist input at admission**

29 Six RCTs (n=401) evaluated the role of pharmacists at admission for improving outcomes.<sup>14 22 24-26 31</sup> The  
30 pharmacists in the intervention arms in these studies were mainly involved at the admission stage of the patient  
31 journey, for example participating in post-take ward rounds, medicines reconciliation and taking medication  
32 history. The evidence suggested that pharmacist input at admission may provide benefit in improving

33 identification of medication discrepancies during medicines' reconciliation at admission (+0.36 (95% CI: 0.07  
34 to 0.65), 2 studies, low quality), reducing medication errors within 24 hours of admission (RR= 0.05 (95% CI:  
35 0.03 to 0.08), 1 study, moderate quality) and increasing physician agreement to act upon medication  
36 discrepancies identified (RR= 1.35 (95% CI: 1.13 to 1.63), 1 study, very low quality). However, there was no  
37 difference for quality of life (EQ-5D visual analogue scale (VAS): + 6.2 (95% CI: -5.7 to 18.1 higher), 1 study,  
38 low quality), LOS (+1.3 hours (-108.96 to 111.56), 1 study, moderate quality), or number of future hospital  
39 admissions (- 0.1 admissions per patient (95% CI: -0.38 to 0.18), 1 study, low quality) and a possible increase in  
40 mortality at 3 months (RR= 1.57 (95% CI: 0.55 to 4.46), 1 study, very low quality). The results are summarised  
41 in the clinical evidence profile in Table 5.3 in Appendix 5 and the Forest plots presented in Appendix 6 of the  
42 Supplementary Materials.

43 One comparative cost analysis (CCA) conducted in the UK showed that pharmacist input at admission was cost  
44 saving compared to usual care (mean saving of £142 per patient).<sup>34</sup> The analysis was assessed as partially  
45 applicable with potentially serious limitations. The results are summarised in Table 5.4, Appendix 5 and the  
46 quality assessment rationale in Appendix 7 in the Supplementary Material.

#### 47 **Ward-based pharmacist input at discharge**

48 Four RCTs (n=770) evaluated provision of ward-based pharmacists' input at discharge.<sup>15 16 18 27</sup> The pharmacists  
49 in the intervention arm in these studies were involved only at the discharge stage, for example preparing  
50 patients' medications and providing counselling before discharge. The evidence suggested a benefit in terms of  
51 reduced prescription errors (RR 0.57 (95% CI: 0.37 to 0.88), 1 study, low quality), reduced readmissions up to  
52 22 days post discharge (RR 0.36 (95% CI: 0.14 to 0.91), 1 study, very low quality) and drug therapy  
53 inconsistencies and omissions at discharge (RR 0.06 (95% CI: 0.01 to 0.44), 1 study, moderate quality). There  
54 was no evidence of effect on quality of life (EQ-5D VAS: 2.8 (95% CI: -1.83 to 7.43), EQ-5D index: 0.05  
55 higher (95% CI: -0.05 to 0.15), 1 study, very low to low quality). The results are summarised in the clinical  
56 evidence profile in Appendix 5, Table 5.3, and the Forest plots presented in Appendix 6 of the Supplementary  
57 Materials.

58 One CUA, conducted in Sweden, showed that the ward-based pharmacist input at discharge was not cost  
59 effective, with an ICER of £327,378 per adjusted QALY gained.<sup>38</sup> The analysis was assessed as partially  
60 applicable with minor limitations. The results are summarised in Appendix 5, Table 5.4 and the quality  
61 assessment rationale in Appendix 7 of the Supplementary Materials.

62 **DISCUSSION**

63 Medication prescribing is the most common healthcare intervention for a patient, and is normally the main  
64 course of treatment for the vast majority. The hospital pharmacist is central to ensuring the quality and safety of  
65 this process. Pharmacist input can be crucial at all stages of the patient journey with different interventions at  
66 each stage. Hence, we stratified the evidence by whether the pharmacist input occurred throughout the patient  
67 stay or was only at admission or discharge. The reviewed evidence for all three strata demonstrated some  
68 benefits for ward-based hospital pharmacist input, although there was variation in the intensity of the  
69 interventions and composition of the comparators. The evidence was of very low to moderate quality due to risk  
70 of bias, imprecision and inconsistency for regular ward-based pharmacist input and ward-based pharmacist input  
71 at discharge. The evidence reviewed for ward-based pharmacist input at admission was of very low to moderate  
72 quality due to risk of bias, imprecision and outcome indirectness as the outcome 'agreement with prescriber'  
73 was used as a surrogate outcome for staff satisfaction and was considered an indirect outcome.

74 The health economic evidence was assessed to be partially applicable (with only 2 studies from the UK and 3  
75 reporting QALYs, which is the outcome measure preferred by NICE). However, it is acknowledged that quality  
76 of life is an outcome that may not be sensitive to pharmacist interventions. Hence, studies reporting other  
77 outcomes were also considered by the committee when making the recommendations.

78 The evidence was also considered to have potentially serious limitations with none of the studies being based on  
79 a review of the evidence base and the cost components included being variable. No clinical or economic  
80 evidence was found relating to 7-day provision of ward-based pharmacist input.

81 Studies assessing the clinical and economic outcomes of the ward-based, clinical pharmacist role have been  
82 accumulating over the years. These studies have generally focused on the effect of pharmacist interventions on  
83 medication errors, medicines reconciliation and savings achieved from reduced medication waste and more  
84 appropriate prescribing. A number of reviews have assessed this evidence in an attempt to draw conclusions  
85 regarding impact on patient outcomes.<sup>39 40</sup> In line with our findings, these reviews have generally shown positive  
86 outcomes including reduced prescribing errors, reduced LOS, reduced admission, and improved patient  
87 satisfaction and physician agreement to act upon medication discrepancies identified. However, overall, the  
88 evidence was relatively weak. The evidence was based mainly on studies with small sample sizes, which  
89 contributed to the high risk of bias in the study outcomes and imprecision around the effect estimates.

90 The mechanism by which pharmacists might improve patient outcomes would most likely be through  
91 minimising prescribing errors, by ensuring appropriate prescribing and also by deprescribing/discontinuation of  
92 drugs. Pharmacist education and input is also likely to improve patient and/or carer satisfaction.<sup>41</sup> Evidence was  
93 found for these outcomes, though not in all strata and with some inconsistencies. For example, some evidence  
94 showed increased prescribing errors at 30 days post discharge, measured by research (rather than intervention)  
95 pharmacists according to the medication appropriateness index and adverse drug events at 3 to 6 months post  
96 discharge. These findings, though unexpected, suggest that the experience of the pharmacist and their  
97 integration in the ward team are likely important factors in achieving positive outcomes, because the  
98 pharmacists in the study reporting this findings were junior pharmacists and new to the ward team. The impact  
99 on quality of life was also modest, which is likely to be due to the acute nature of illness and the short follow-up  
100 periods.

101 Prescription and administration errors are amongst the most commonly identified adverse events during a  
102 patient's stay in hospital.<sup>42</sup> Pharmacists, as part of the multidisciplinary team, can reduce these errors and ensure  
103 that the patient gets the correct treatment, as well as discontinuing drugs which are no longer required in both  
104 the short and long term. The pharmacist has an important educational role which has the potential to improve  
105 patients' adherence after discharge. These activities allow doctors to focus on other key patient care priorities.

106 It is also acknowledged with the aging population that there is an increasing number of patients with multi-  
107 morbidities who are exposed to poly-pharmacy.<sup>8</sup> In this situation the pharmacist can play a vital role in advising  
108 the medical team regarding drugs and how to prescribe treatment optimally. Involving the pharmacist at hospital  
109 discharge may have reduced the need for junior doctors to explain prescribing regimens, and the need for the  
110 patient to visit their general practitioner following discharge for drug review. This would improve patient and/or  
111 carer satisfaction and have a potential cost benefit.

112 Pharmacists are also gradually acquiring independent prescribing rights.<sup>43 44</sup> This allows them to correct  
113 prescribing errors or make changes directly without the need for doctor involvement. Streamlining the  
114 prescribing of medications to take home at the end of hospital stay could also facilitate earlier discharge and  
115 allow junior doctors to focus on other tasks produced from the ward rounds.<sup>45</sup> Assessment of the cost  
116 effectiveness of prescribing pharmacists in hospital should include these considerations.

117 The cost effectiveness of the ward-based pharmacist role has been assessed in a number of published economic  
118 evaluations. However, unlike the evidence for clinical effectiveness, the generalisability of the findings of these

119 evaluations from one health care system to another might be limited due to the different funding arrangements  
120 and the perspectives used in the analysis.<sup>46-48</sup>

121 The economic evidence in our review was in favour of the provision of ward-based pharmacist input but the  
122 interventions, and therefore results, varied from one country to another. Clinical pharmacists in the reviewed  
123 UK studies were generally experienced (band 7/8) and have specialist knowledge in the medications they  
124 managed. They also were completely integrated in their clinical teams.<sup>36</sup> This may not be the same profile in the  
125 non-UK studies. Additionally, the standard care/control arm in the included economic studies was not always  
126 clearly defined and was variable in terms of the level of pharmacist input. Some studies included a specified  
127 level of clinical pharmacist input in the control group which was enhanced in the intervention group (for  
128 example, by attendance at ward rounds) while others described the introduction of a completely new service.  
129 These differences might explain the differences in the findings of these studies, which has also been highlighted  
130 by other reviews of this evidence.<sup>47</sup>

131 With the exception of the UK economic modelling study,<sup>36</sup> all economic studies had a follow-up of 12 months  
132 or less and hence would not have assessed the long term impact of the ward-based pharmacist intervention.  
133 Additionally, the majority of the studies assessed a limited number of cost categories; focusing on medication  
134 costs, pharmacist time and less on other staff time (e.g. freeing up or release of junior doctor time) and patient-  
135 related downstream costs.

136 There was evidence that pharmacist input throughout the hospital stay would achieve saving in terms of  
137 medications costs, which was the most frequently assessed cost category in the included studies. One study  
138 found the pharmacist cost was completely offset by medication cost savings.<sup>21</sup> The evidence was less clear in  
139 terms of impact on other staff time and on long-term patient outcomes, which were not always assessed in the  
140 included studies. Where this impact was quantified, the results showed potential for cost saving. Avoiding  
141 medication errors that have severe consequences is also an important positive outcome in terms of avoiding  
142 litigation costs.<sup>36</sup> Overall, the economic evidence suggested that the regular input by ward-based pharmacists is  
143 cost-effective. Pharmacist input only at discharge was not cost effective, but the evidence for this was limited to  
144 one Swedish study.<sup>38</sup>

145 This systematic review demonstrates the potential benefits for patient safety of including ward-based  
146 pharmacists in the multidisciplinary team in hospital. Our focus on higher-quality studies permits robust  
147 conclusions. However, sample sizes tended to be small, there was some heterogeneity between the interventions

148 studied, and we did not formally assess publication bias. Nevertheless, our findings are consistent with earlier  
149 reviews and have strong face-validity, allowing the guideline committee to recommend the routine inclusion of  
150 ward-based pharmacists in the multidisciplinary team managing acutely ill hospitalised patients.<sup>49</sup>

## 151 **CONCLUSION**

152 Evaluations of the ward-based pharmacist input have largely found it to be both effective and cost effective,  
153 particularly when provided throughout the different stages of the patient journey by experienced pharmacists  
154 who are integrated in the ward team. The effectiveness evidence, however, was generally of low quality. The  
155 economic evidence had potentially serious limitations. The interventions and comparators were often not well  
156 defined and there was variation across the studies in their composition.

157 Nevertheless, the collective body of the available evidence suggests that recommending regular ward-based  
158 pharmacist input and inclusion in the multidisciplinary team would offer additional value to the provision of  
159 care for those admitted for a suspected or confirmed medical emergency. However, further research is needed to  
160 determine the optimal level of involvement of ward-based pharmacists and to assess whether the provision of a  
161 7-day service is cost-effective.

## 162 **ACKNOWLEDGMENT**

163 The guideline development group members who contributed to the protocol development and commented to the  
164 results, informing the discussion of our findings. Mr Joseph Runicles, Information Scientist, for his help in  
165 compiling the bibliography. National Institute for Health and Care Excellence (2018). Emergency and acute  
166 medical care in over 16s: service delivery and organisation Available from  
167 <https://www.nice.org.uk/guidance/ng94>.

## 168 **FUNDING**

169 This work was undertaken by the National Guideline Centre (NGC), Royal College of Physicians London which  
170 received funding from NICE. The views expressed in this publication are those of the authors and not  
171 necessarily those of the Institute. The funding body (NICE) did not play any direct role in the study design; the  
172 collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article  
173 for publication. All researchers involved in this work were independent from the funding body at the time of  
174 completing this work. All authors, external and internal to the NGC, had full access to all of the data (including

175 statistical reports and tables) in the study and take responsibility for the integrity of the data and the accuracy of  
176 the data analysis.

177

ACCEPTED MANUSCRIPT



178 **References**

- 179 1. Gandhi TK, Weingart SN, Borus J, et al. Adverse drug events in ambulatory care. *N Engl J Med*  
180 2003;348:1556-1564.
- 181 2. Kaushal R, Bates DW, Landrigan C, et al. Medication errors and adverse drug events in pediatric inpatients.  
182 *JAMA* 2001;285:2114-20.
- 183 3. Bates DW, Cullen DJ, Laird N, et al. Incidence of adverse drug events and potential adverse drug events.  
184 Implications for prevention. ADE Prevention Study Group. *JAMA* 1995;274:29-34.
- 185 4. Hepler CD, Strand LM. Opportunities and responsibilities in pharmaceutical care. *Am J Hosp Pharm*  
186 1990;47:533-43.
- 187 5. Perez A, Doloresco F, Hoffman JM, et al. ACCP: economic evaluations of clinical pharmacy services: 2001-  
188 2005. *Pharmacotherapy* 2009;29:128.
- 189 6. Olalekan K, Young D, Baker J, et al. Hospital pharmacists' contribution: a perspective. Hospital Pharmacy  
190 Europe. [http://www.hospitalpharmacyeurope.com/featured-articles/hospital-pharmacists%E2%80%99-](http://www.hospitalpharmacyeurope.com/featured-articles/hospital-pharmacists%E2%80%99-contribution-perspective)  
191 [contribution-perspective](http://www.hospitalpharmacyeurope.com/featured-articles/hospital-pharmacists%E2%80%99-contribution-perspective) ; 2015 Accessed 10.07.18.
- 192 7. Baqir W, Miller D, Richardson G. A brief history of pharmacist prescribing in the UK. *Eur J Hosp Pharm Sci*  
193 *Pract* 2012;19:487-488.
- 194 8. National Institute for Health and Care Excellence. Multimorbidity: clinical assessment and management.  
195 NICE guideline 56. London, 2016.
- 196 9. Landeiro F, Roberts K, Gray AM, et al. Delayed Hospital Discharges of Older Patients: A Systematic Review  
197 on Prevalence and Costs. *Gerontologist* 2017; doi: 10.1093/geront/gnx028. [Epub ahead of print].
- 198 10. National Institute for Health and Care Excellence. Acute medical emergencies, service guidance scope.  
199 London, 2015.
- 200 11. National Institute for Health and Care Excellence. Developing NICE guidelines: the manual. London, 2014.
- 201 12. Group GW. The Grading of Recommendations Assessment, Development and Evaluation (GRADE)  
202 Working Group website. <http://www.gradeworkinggroup.org/> ; 2011 Accessed 10.07.18.
- 203 13. OECD. Purchasing Power Parities (PPP), 2016.
- 204 14. Aag T, Garcia BH, Viktil KK. Should nurses or clinical pharmacists perform medication reconciliation? A  
205 randomized controlled trial. *Eur J Clin Pharmacol* 2014;70:1325-32.
- 206 15. Al-Rashed SA, Wright DJ, Roebuck N, et al. The value of inpatient pharmaceutical counselling to elderly  
207 patients prior to discharge. *Br J Clin Pharmacol* 2002;54:657-64.

- 208 16. Bladh L, Ottosson E, Karlsson J, et al. Effects of a clinical pharmacist service on health-related quality of  
209 life and prescribing of drugs: a randomised controlled trial. *BMJ Qual Saf* 2011;20:738-46.
- 210 17. Claus BOM, Robays H, Decruyenaere J, et al. Expected net benefit of clinical pharmacy in intensive care  
211 medicine: a randomized interventional comparative trial with matched before-and-after groups. *J Eval Clin*  
212 *Pract* 2014;20:1172-79.
- 213 18. Eggink RN, Lenderink AW, Widdershoven JG, et al. The effect of a clinical pharmacist discharge service on  
214 medication discrepancies in patients with heart failure. *Pharm World Sci* 2010;32:759-766.
- 215 19. Farley TM, Shelsky C, Powell S, et al. Effect of clinical pharmacist intervention on medication  
216 discrepancies following hospital discharge. *Int J Clin Pharm* 2014;36:430-437.
- 217 20. Farris KB, Carter BL, Xu Y, et al. Effect of a care transition intervention by pharmacists: an RCT. *BMC*  
218 *Health Serv Res* 2014;14:406.
- 219 21. Gillespie U, Alassaad A, Henrohn D, et al. A comprehensive pharmacist intervention to reduce morbidity in  
220 patients 80 years or older: a randomized controlled trial. *Arch Intern Med* 2009;169:894-900.
- 221 22. Khalil V, deClifford JM, Lam S, et al. Implementation and evaluation of a collaborative clinical pharmacist's  
222 medications reconciliation and charting service for admitted medical inpatients in a metropolitan hospital. *J Clin*  
223 *Pharm Ther* 2016;4:662-666.
- 224 23. Kucukarslan SN, Peters M, Mlynarek M, et al. Pharmacists on rounding teams reduce preventable adverse  
225 drug events in hospital general medicine units. *Arch Intern Med* 2003;163:2014-2018.
- 226 24. Lind KB, Soerensen CA, Salamon SA, et al. Impact of clinical pharmacist intervention on length of stay in  
227 an acute admission unit: a cluster randomised study. *Eur J Hosp Pharm Sci Pract* 2016;23:171-176.
- 228 25. Lisby M, Thomsen A, Nielsen LP, et al. The effect of systematic medication review in elderly patients  
229 admitted to an acute ward of internal medicine. *Basic Clin Pharmacol Toxicol* 2010;106:422-427.
- 230 26. Nester TM, Hale LS. Effectiveness of a pharmacist-acquired medication history in promoting patient safety.  
231 *Am J Health Syst Pharm* 2002;59:2221-225.
- 232 27. Nickerson A, MacKinnon NJ, Roberts N, et al. Drug-therapy problems, inconsistencies and omissions  
233 identified during a medication reconciliation and seamless care service. *Healthcare Quart* 2005;8:65-72.
- 234 28. Scullin C, Scott MG, Hogg A, et al. An innovative approach to integrated medicines management. *J Eval*  
235 *Clin Pract* 2007;13:781-788.
- 236 29. Shen J, Sun Q, Zhou X, et al. Pharmacist interventions on antibiotic use in inpatients with respiratory tract  
237 infections in a Chinese hospital. *Int J Clin Pharm* 2011;33:929-933.
- 238 30. Spinewine A, Swine C, Dhillon S, et al. Effect of a collaborative approach on the quality of prescribing for  
239 geriatric inpatients: a randomized, controlled trial. *J Am Geriatr Soc* 2007;55:658-665.

- 240 31. Tong EY, Roman C, Mitra B, et al. Partnered pharmacist charting on admission in the General Medical and  
241 Emergency Short-stay Unit - a cluster-randomised controlled trial in patients with complex medication  
242 regimens. *J Clin Pharm Therap* 2016;41:414-418.
- 243 32. Zhao S, Zhao H, Du S, et al. The impact of clinical pharmacist support on patients receiving multi-drug  
244 therapy for coronary heart disease in china: a long-term follow-up study. *Eur J Hosp Pharm* 2015;22:323-327.
- 245 33. Zhao S, Zhao H, Du S, et al. Impact of pharmaceutical care on the prognosis of patients with coronary heart  
246 disease receiving multidrug therapy. *Pharm Care Res* 2015;15:179-181.
- 247 34. Fertleman M, Barnett N, Patel T. Improving medication management for patients: the effect of a pharmacist  
248 on post-admission ward rounds. *Qual Saf Health Care* 2005;14:207-121.
- 249 35. Ghatnekar O, Bondesson A, Persson U, et al. Health economic evaluation of the Lund Integrated Medicines  
250 Management Model (LIMM) in elderly patients admitted to hospital. *BMJ Open* 2013;3:e001563. doi:  
251 10.1136/bmjopen-2012-001563
- 252 36. Karnon J, McIntosh A, Dean J, et al. Modelling the expected net benefits of interventions to reduce the  
253 burden of medication errors. *J Health Servs Res Policy* 2008;13:85-91.
- 254 37. Klopotoska JE, Kuiper R, vanKan HJ, et al. On-ward participation of a hospital pharmacist in a Dutch  
255 intensive care unit reduces prescribing errors and related patient harm: an intervention study. *Crit Care*  
256 2010;14:R174.
- 257 38. Wallerstedt SM, Bladh L, Ramsberg J. A cost-effectiveness analysis of an in-hospital clinical pharmacist  
258 service. *BMJ Open* 2012;2:e00032. doi: 10.1136/bmjopen-2011-000329
- 259 39. Graabaek T, Kjeldsen LJ. Medication reviews by clinical pharmacists at hospitals lead to improved patient  
260 outcomes: a systematic review. *Basic Clin Pharmacol Toxicol* 2013;112:359-373.
- 261 40. Zegers M, Hesselink G, Geense W, et al. Evidence-based interventions to reduce adverse events in hospitals:  
262 a systematic review of systematic reviews. *BMJ Open* 2016;6: e012555. doi: 10.1136/bmjopen-2016-012555
- 263 41. Dunn SP, Birtcher KK, Beavers CJ, et al. The Role of the Clinical Pharmacist in the Care of Patients With  
264 Cardiovascular Disease. *J Am Col Cardio* 2015;66:2129-2139.
- 265 42. Slawomirski L, Auraaen A, Klazinga N. The Economics of Patient Safety: Strengthening a value-based  
266 approach to reducing patient harm at national level: Organisation for Economic Co-operation and Development:  
267 Paris, France; 2017.
- 268 43. Dawoud D, Griffiths P, Maben J, et al. Pharmacist supplementary prescribing: a step toward more  
269 independence? *RSAP* 2011;7:246-256.
- 270 44. Law MR, Ma T, Fisher J, et al. Independent pharmacist prescribing in Canada. *Can Pharm J* 2012;145:17-  
271 23.

- 272 45. Bourne RS, Baqir W, Onatade R. Pharmacist independent prescribing in secondary care: opportunities and  
273 challenges. *Int J Clin Pharm* 2016;38:1-6.
- 274 46. Touchette DR, Doloresco F, Suda KJ, et al. Economic evaluations of clinical pharmacy services: 2006-2010.  
275 *Pharmacotherapy* 2014;34:771-793.
- 276 47. Gallagher J, McCarthy S, Byrne S. Economic evaluations of clinical pharmacist interventions on hospital  
277 inpatients: a systematic review of recent literature. *Int J Clin Pharm* 2014;36:1101-1114.
- 278 48. Gammie T, Vogler S, Babar ZU. Economic Evaluation of Hospital and Community Pharmacy Services. *Ann*  
279 *Pharmacother* 2017;51:54-65.
- 280 49. National Clinical Guideline Centre. Emergency and acute medical care in over 16s: service delivery and  
281 organisation [Chapter 30: Pharmacist support]. London, 2018.
- 282