

A broad-spectrum synthesis of Tetravinylethylenes

Horvath, Kelsey; Newton, Christopher; Roper, Kimberley; Ward, Jas S; Sherburn, Michael

DOI:

[10.1002/chem.201900550](https://doi.org/10.1002/chem.201900550)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Horvath, K, Newton, C, Roper, K, Ward, JS & Sherburn, M 2019, 'A broad-spectrum synthesis of Tetravinylethylenes', *Chemistry: A European Journal*, vol. 25, no. 16, pp. 4072-4076.
<https://doi.org/10.1002/chem.201900550>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility 08/02/2019

Published in *Chemistry: A European Journal*
<https://doi.org/10.1002/chem.201900550>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

CHEMISTRY

A European Journal

A Journal of



Accepted Article

Title: A Broad-Spectrum Synthesis of Tetravinylenes

Authors: Michael Sherburn, Kelsey Horvath, Kimberley Roper,
Christopher Newton, and Jas S Ward

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Eur. J.* 10.1002/chem.201900550

Link to VoR: <http://dx.doi.org/10.1002/chem.201900550>

Supported by
ACES

WILEY-VCH

A Broad-Spectrum Synthesis of Tetravinylethylenes

Kelsey L. Horvath,^[a] Christopher G. Newton,^[a] Kimberley A. Roper,^[a] Jas S. Ward^[a] and Michael S. Sherburn^{*[a]}


Abstract: The first general synthesis of compounds of the tetravinylethylene (TVE) family is reported. Ramirez-type dibromo-olefination of readily accessible penta-1,4-dien-3-ones generates 3,3-dibromo[3]dendralenes, which undergo twofold Negishi, Suzuki-Miyaura or Mizoroki-Heck reactions with a wide variety of olefinic coupling partners. This route delivers a broad range of unsymmetrically-substituted tetravinylethylenes with up to three different alkenyl substituents attached to the central C=C bond. The extensive scope of the approach is demonstrated by the preparation of the first higher order oligo-alkenic through-conjugated/cross-conjugated hybrid compounds. An unsymmetrically-substituted TVE is shown to undergo a domino electrocyclic-cycloaddition with high site-selectivity and diastereoselectivity, thereby demonstrating the substantial synthetic potential of substituted TVEs for controlled, rapid structural complexity generation.

Tetravinylethylene (TVE) is one of the smallest structures exhibiting both through-conjugation and cross-conjugation (Scheme 1). Specifically, the structure comprises intersecting pairs of (*E*)-1,3,5-hexatrienes, (*Z*)-1,3,5-hexatrienes and 2-methylene-1,4-pentadienes ([3]dendralenes). Nothing is known about the impact of substitution upon the behavior of TVEs and how this might be exploited. Furthermore, related structures containing more than five C=C units have not been reported in the literature. The first synthesis of TVE, a landmark contribution from the classical era of unsaturated hydrocarbon chemistry, was reported by Skattebøl and co-workers in the 1960s.¹⁻³ We recently disclosed the first preparatively useful approach to TVE, a one-step multi-gram scale synthesis, and demonstrated that the hydrocarbon is bench-stable when stored as a neat liquid.^{4,5} The one-step synthesis of TVE (Scheme 1, **1a**) utilized a Pd(0)-catalyzed fourfold Stille cross-coupling sequence between tetrachloro-ethylene **2** and vinyltributylstannane **3a**. In addition to the parent TVE, five symmetrically-substituted TVEs **1b** were similarly prepared using substituted alkenyltributylstannanes **3b** in the exhaustive Stille cross-coupling process.^{4,5}


As a starting point for future synthetic applications, the parent TVE has been shown to rapidly generate multi-cyclic systems.^{4,5} Additionally, the hybrid through/cross-conjugated structure of the TVE unit represents the smallest organic four-directional branch point in a possible conducting molecular wire.⁶ To realize the potential of the TVE family in target synthesis and materials applications, synthetic access to unsymmetrically-substituted systems must be granted. Until now, no unsymmetrically-substituted TVEs have been reported. Our previous synthetic approach (Scheme 1), in common with all other substituted TVE

syntheses reported in the literature,⁷ is only suited to the attachment of four identical substituted alkenyl-groups to the central C=C core.

introduction to TVE




previous work^{4,5}



• fourfold Stille approach to TVE and symmetrically-substituted TVEs

this work



• first synthesis of unsymmetrically-substituted TVEs
 • di-, tetra-, hexasubstituted TVEs with up to 3 different substituents
 • no organotin: Negishi, Suzuki-Miyaura, Mizoroki-Heck reactions

Scheme 1. The existing synthetic route to TVEs with its limitations, and the new, broad scope synthesis.

Herein we report a broad-reaching solution, delivering the first 29 unsymmetrically substituted TVEs carrying up to three different substituents (Scheme 1, bottom). We also report the first compounds exhibiting extended π -systems based upon TVE subunits. Moreover, we establish the synthetic potential of these unsymmetrically-substituted TVEs, by way of a one-pot, site-selective and diastereoselective domino pericyclic sequence that generates three C–C bonds and six contiguous stereocenters. The new synthetic approach (Scheme 1, bottom) involves the conversion of penta-1,4-dien-3-ones **4** (readily accessed through twofold aldol-type condensations) into 3,3-dibromo-[3]dendralenes **5**, which in turn undergo a range of twofold cross-couplings to furnish diversely-substituted tetravinylethylenes **6**.

Twelve substituted penta-1,4-dien-3-ones **4** were prepared, by way of sequential Claisen-Schmidt condensations⁸ (see SI for details). There are only three examples of Ramirez dibromo-olefinations of penta-1,4-dien-3-ones **4** in the literature⁹⁻¹¹ and, in our hands, low yields were obtained under standard Ramirez reaction conditions.¹² Gratifyingly, in the majority of cases, the modified protocol introduced by Lautens and co-workers¹³

[a] Kelsey L. Horvath, Dr Christopher G. Newton, Dr Kimberley A. Roper, Dr Jas S. Ward, Prof Michael S. Sherburn*
 Research School of Chemistry
 Australian National University
 Canberra, ACT 2601 (Australia)
 E-mail: michael.sherburn@anu.edu.au

Supporting information for this article is given via a link at the end of the document.

furnished workable yields, and twelve substituted 3,3-dibromo[3]dendralenes **5** were prepared in this manner (Table 1).





Table 1. Lautens-modified Ramirez dibromo-olefination of penta-1,4-dien-3-ones **4** permits the first general synthesis of 3,3-dibromo[3]dendralenes **5**.

Unlike some other dendralenic structures, which exhibit a propensity to dimerize and polymerize,¹⁵ 3,3-dibromo[3]dendralenes **5** are generally white or yellow crystalline solids that can be stored neat without appreciable decomposition, and are easily handled using standard laboratory techniques.

While there are many examples of cross-coupling reactions of 1,1-dibromoalkenes in the literature,¹⁶ there is only one example of a cross-coupling of a 3,3-dibromo[3]dendralene, involving the Sonogashira sp^2 - sp coupling of a cyclic system with trimethylsilylacetylene.¹⁰ In light of the lack of precedent, both for sp^2 - sp^2 cross-couplings and also for reactions of acyclic systems, we were delighted to find that twofold Negishi reactions of 3,3-dibromo[3]dendralenes **5** with vinylzinc bromide proceed well using [Pd(dppf)Cl₂] as pre-catalyst to form the first unsymmetrical TVEs **6** (Table 2).





Table 2. Twofold Negishi cross-coupling of 3,3-dibromo[3]dendralenes **5** with vinylzinc bromide furnishes the first unsymmetrical tetra- and hexa-substituted TVEs **6**.

3,3-Dibromo[3]dendralenes comprising the same (**5a-g**) or different (**5h-l**) alkenyl substituents performed equally well, to generate the first examples of TVEs possessing two or three different alkenyl groups about the central C=C bond. TVEs bearing carbocyclic aromatic substituents with different electronic characteristics (**6j-l**) and the first TVEs carrying heterocyclic aromatic substituents (**6f-i**) are included in this selection.

Twofold cross-couplings of 3,3-dibromo[3]dendralenes **5** are by no means limited to the introduction of a pair of unsubstituted vinyl groups: in fact, alkenyl groups bearing substituents at any position are tolerated. The syntheses of eighteen of these more highly substituted TVEs are summarized in Table 3. Double Negishi couplings of 3,3-dibromo[3]dendralene **5a** with 2-propenyl-, (*E*)-styrenyl- and (2-methylprop-1-en-1-yl)zinc bromides furnished six novel tetra- and hexa-substituted TVEs **6n-s**. Cross-couplings other than Negishi reactions are also successful, as demonstrated by the products of double Suzuki-Miyaura¹⁷ and Mizoroki-Heck¹⁸ processes, **6t-v** (3 examples)

and **6w-ae** (9 examples), respectively. These results establish the wide scope of the 3,3-dibromo[3]dendralene cross-coupling approach to TVE synthesis and its broad tolerance of substitution upon the incoming alkenyl-groups.






Table 3. Twofold Negishi, Suzuki-Miyaura or Mizoroki-Heck reactions grant access to unsymmetrical, multi-substituted tetravinylethylenes. ^aX-ray crystal structures of these four TVEs²² are in the SI.

Following the trend previously seen,^{4,5} the 25 new TVEs depicted in Tables 2 and 3 are bench-stable substances. No special equipment or methods are required when manipulating these compounds: they survive storage neat at ambient temperature for several weeks. The lack of susceptibility of this new family of π -bond rich systems towards uncatalyzed Diels-Alder dimerization and autoxidation (typical of related cross-conjugated¹⁵ and through-conjugated¹⁹ polyenic systems) bodes well for future applications.

Access to unsymmetrically-substituted TVEs promotes new opportunities in rapid structure complexity generation, and an interesting question of diene site selection in [4+2]cycloadditions of polyenes (Scheme 2). Heating diphenyl-di-(1-naphthyl)-TVE **6aa** in benzene brings about a disrotatory electrocycloaddition to generate cyclic [4]dendralene **7**, with a *cis*-disposition between the two cyclohexadiene substituents. When performed in the presence of dienophile *N*-*tert*-butylmaleimide **8**, a subsequent *endo*-mode cycloaddition occurs, with approach to the less sterically encumbered π -diastereoface of [4]dendralene **7**. The two non-equivalent semi-cyclic dienes of [4]dendralene **7** (highlighted in different colors in Scheme 2) are expected to be more reactive than the cyclohexadiene site,⁵ and the phenyl-substituted diene is preferred over the 1-naphthyl-substituted site. Overall, this operationally trivial domino sequence generates densely-functionalized decalin **9**, with three new C–C bonds, six contiguous stereocenters, and a cyclic [3]dendralene subunit that invites additional synthetic manipulation.²⁰



Scheme 2. Stereoselective domino [6 π]electrocyclization/[4+2]cycloaddition of TVE **6aa** proceeds with diene site selectivity to generate multicyclic product **9**. H atoms not attached to stereocenters are omitted from the X-ray crystal structure^{22,23} for clarity.



Scheme 3. First preparation of higher order acyclic C=C-based π -frameworks related to TVE. The new π -systems are highlighted in green. Ethyl groups are truncated in the X-ray crystal structure^{22,23} for clarity.

The robust nature of the new pentaene-based TVE compounds, combined with their ease of synthesis, prompted the question of whether larger structures with unprecedented acyclic π -frameworks were within reach. The successful extension of these methods to the syntheses of hexaenes **14** and **15**, and heptaenes **13**, **16** and **17** are depicted in Scheme 3. Dibromotetraene **10**, dibromo-pentaene **11** and tetrabromo-triene **12**, prepared through Lautens-modified Ramirez dibromo-olefination reactions of commercially available ketonic precursors (see SI), underwent twofold or fourfold Negishi or Mizoroki-Heck couplings to furnish the bench-stable polyenic products. The UV absorption spectrum of heptaene **17** exhibits features attributable to the longest through-conjugated subunit (see SI). The solid state molecular structure of tetra-ester **17**, obtained from single crystal X-ray analysis, is consistent with this observation, revealing an essentially in-plane 2,4,6,8,10-dodecapentaenoate unit (Scheme 3). Acyclic, hybrid cross-conjugated/through-conjugated C=C structures are very poorly represented in the literature.²¹ The structures depicted in Scheme 3 represent the first examples of substances of this type.

In summary, we have described the first general synthetic approach to TVEs and related through/cross-conjugated acyclic C=C-based systems. This work reports 34 new TVEs and related structures, including the first examples exhibiting unsymmetrical substitution patterns, extended through-conjugation and/or cross-conjugation. The synthetic approach is robust, involving twofold Negishi, Suzuki-Miyaura or Mizoroki-Heck couplings of acyclic 3,3-dibromo[3]dendralenes. The new structures are bench stable compounds that undergo operationally simple, selective, complexity-generating domino sequences. The acyclic C=C-bond-rich structures described herein are representatives of a sizeable region of structural space that was previously inaccessible but is now available for exploration and investigation. Applications in target synthesis and in conducting materials will follow.

Acknowledgements

This work was supported by the Australian Research Council.

Keywords: hydrocarbons • polyenes • cross-coupling • electrocyclic reactions • cycloaddition


- [1] L. Skattebøl, *Acta. Chem. Scand.* **1963**, *17*, 1683–1693.
- [2] L. Skattebøl, S. Solomon, *J. Am. Chem. Soc.* **1965**, *87*, 4506–4513.
- [3] L. Skattebøl, J. L. Charleton, P. deMayo, *Tetrahedron Lett.* **1966**, *7*, 2257–2260.
- [4] E. J. Lindeboom, A. C. Willis, M. N. Paddon-Row, M. S. Sherburn, *Angew. Chem. Int. Ed.* **2014**, *79*, 5440–5443.
- [5] E. J. Lindeboom, A. C. Willis, M. N. Paddon-Row, M. S. Sherburn, *J. Org. Chem.* **2014**, *53*, 11496–11507.
- [6] To our knowledge, only aromatic ring-based (and not C=C bond-based) branch points have been reported. Despite their significant potential, it was recently noted that branched molecular wires are rarely fabricated in a precise manner: R. Sakamoto, S. Katagiri, H. Maeda, Y. Nishimori, S. Miyashita, H. Nishihara, *J. Am. Chem. Soc.* **2015**, *137*, 734–741.
- [7] (a) R. W. Bost, J. A. Krynitsky, *J. Am. Chem. Soc.* **1948**, *70*, 1027–1029; (b) F. Bohlmann, K. Kieslich, *Abh. Braunschweig. Wiss. Gess.* **1957**, *147*; (c) P. Cadiot, W. Chodkiewicz, J. Pauss-Godineau, *Bull. Soc. Chim. Fr.* **1961**, 2176–2193; (d) R. Kuhn, B. Schulz, *Chem. Ber.* **1963**, *96*, 3200–3208; (e) R. Kuhn, B. Schulz, *Angew. Chem.* **1963**, *75*, 452–453; (f) L. Lombardo, F. Sondheimer, *Synthesis* **1980**, 950–952; (g) J.-M. Pons, M. Santelli, *J. Org. Chem.* **1989**, *54*, 877–884.
- [8] (a) L. Claisen, A. Claparède, *Ber.* **1881**, *14*, 349–353; (b) J. G. Schmidt, *Ber.* **1881**, *14*, 1459–1461.
- [9] (a) R. Kuhn, H. Fischer, D. Rewicki, H. Fischer, *Liebigs Ann. Chem.* **1965**, *689*, 1–20; (b) C. Raulet, E. Levas, *Bull. Soc. Chim. Fr.* **1971**, 2598–2606.
- [10] H. Hopf, J. Kampen, P. Bubenitschek, P. G. Jones, *Eur. J. Org. Chem.* **2002**, 1708–1721.
- [11] Dichloro-olefination of acyclic penta-1,4-dien-3-ones with dichloroketene proceeds in low yields: J. Ojima, K. Itagawa, S. Hamai,

- T. Nakada, S. Kuroda, *J. Chem. Soc., Perkin Trans. 1* **1983**, 2997–3004.
- [12] N. B. Desai, N. McKelvie, F. Ramirez, *J. Am. Chem. Soc.* **1962**, *84*, 1745–1747.
- [13] (a) Y.-Q. Fang, O. Lifchits, M. Lautens, *Synlett* **2008**, *3*, 413–417; (b) C. Bryan, V. Aurregi, M. Lautens, *Org. Synth.* **2009**, *86*, 36–46.
- [14] For a review on dendralenes, see: H. Hopf, M. S. Sherburn, *Angew. Chem., Int. Ed.* **2012**, *51*, 2298–2338.
- [15] (a) A. T. Blomquist, J. A. Verdol, *J. Am. Chem. Soc.* **1955**, *77*, 81–83; (b) J. Mulzer, G. Brüntrup, U. Kühl and G. Hartz, *Chem. Ber.* **1982**, *115*, 3453–3469; (c) H. Toombs-Ruane, E. L. Pearson, M. N. Paddon-Row, M. S. Sherburn, *Chem. Commun.* **2012**, *48*, 6639–6641.
- [16] Reviews: (a) F. Legrand, K. Jouvin, G. Evano, *Isr. J. Chem.* **2010**, *50*, 588–604; (b) G. Chelucci, *Chem. Rev.* **2012**, *112*, 1344–1462.
- [17] These transformations adapt conditions reported in S. Lou, G. C. Fu, *Adv. Synth. Catal.* **2010**, *352*, 2081–2084.
- [18] These reactions adapt the protocol described in A. F. Littke, G. C. Fu, *J. Am. Chem. Soc.* **2001**, *123*, 6989–7000.
- [19] F. Sondheimer, D. A. Ben-Efraim, R. Wolovsky, *J. Am. Chem. Soc.* **1961**, *83*, 1675–1681.
- [20] For important synthetic contributions involving cyclic [3]dendralenes and their oxa-[3]dendralene analogs, see the following representative papers (a) K. M. Brummond, H. Chen, P. Sill, L. You, *J. Am. Chem. Soc.* **2002**, *124*, 15186–15187; (b) S. Werner, D. M. Turner, P. G. Chambers, K. M. Brummond, *Tetrahedron* **2008**, *64*, 6997–7007; (c) C. Spino, G. Liu, *J. Org. Chem.* **1993**, *58*, 817–819; (d) S. Perreault, C. Spino, *Org. Lett.* **2006**, *8*, 4385–4388.
- [21] *Cross Conjugation: Modern Dendralene, Radialene and Fulvene Chemistry*, H. Hopf, M. S. Sherburn, eds., Wiley-VCH 2016. 464 pages. ISBN 978-3-527-33437-7.
- [22] CCDC 1890206, 1890207, 1890208, 1890213, 1890214 and 1890215 contain the supplementary crystallographic data for this paper. These data are provided free of charge by The Cambridge Crystallographic Data Centre.
- [23] Molecular structures from single crystal X-ray analyses were visualized using CYLview 1.0b; C. Y. Legault, Université de Sherbrooke, 2009 (<http://www.cylview.org>)

Entry for the Table of Contents

COMMUNICATION

Master key for TVE. The first general synthesis of tetravinylethylenes (TVEs) involves the preparation and cross-coupling of dibromo-[3]dendralenes. Their value in controlled, rapid structural complexity generation is demonstrated. The first higher order acyclic oligo-alkenic through-conjugated/cross-conjugated hybrid compounds are also reported.



*Kelsey L. Horvath, Christopher G. Newton, Kimberley A. Roper, Jas S. Ward, Michael S. Sherburn**

Page No. – Page No.

A Broad-Spectrum Synthesis of Tetravinylethylenes