

GALEX observations of "Passive Spirals" in the Cluster Cl0024+17

Moran, Sean M.; Ellis, Richard S.; Treu, Tommaso; Salim, Samir; Rich, R. Michael; Smith, Graham; Kneib, Jean-Paul

DOI:
[10.1086/504078](https://doi.org/10.1086/504078)

License:
Other (please specify with Rights Statement)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Moran, SM, Ellis, RS, Treu, T, Salim, S, Rich, RM, Smith, G & Kneib, J-P 2006, 'GALEX observations of "Passive Spirals" in the Cluster Cl0024+17: clues to the formation of S0 galaxies', *The Astrophysical Journal*, vol. 641, no. 2, pp. L97-100. <https://doi.org/10.1086/504078>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:
© 2006. The American Astronomical Society. All rights reserved. Printed in U.S.A.

Published in *Astrophysical Journal Letters* on 03/04/2006

<https://doi.org/10.1086/504078>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

GALEX OBSERVATIONS OF “PASSIVE SPIRALS” IN THE CLUSTER CI 0024+17: CLUES TO THE FORMATION OF S0 GALAXIES

SEAN M. MORAN,¹ RICHARD S. ELLIS,¹ TOMMASO TREU,² SAMIR SALIM,³
R. MICHAEL RICH,³ GRAHAM P. SMITH,^{1,4} AND JEAN-PAUL KNEIB⁵

Received 2006 January 10; accepted 2006 March 9; published 2006 April 3

ABSTRACT

We present new results from deep *GALEX* UV imaging of the cluster CI 0024+17 at $z \sim 0.4$. Rest-frame far-UV emission is detected from a large fraction of so-called passive spiral galaxies—a significant population that exhibits spiral morphology with little or no spectroscopic evidence of ongoing star formation. This population is thought to represent infalling galaxies whose star formation has been somehow truncated by environmental processes, possibly in morphological transition to S0 galaxies. Compared to normal cluster spirals, we find that passive spirals are redder in FUV-optical color, while exhibiting much stronger UV emission than cluster E/S0 galaxies—as expected for recently truncated star formation. By modeling the different temporal sensitivities of UV and spectroscopic data to recent activity, we show that star formation in passive spirals decayed on timescales of less than 1 Gyr, consistent with “gas starvation”—a process where the cluster environment prevents cold gas from accreting onto the spiral disk. Intriguingly, the fraction of spirals currently observed in the passive phase is consistent with the longer period expected for the morphological transformation and the subsequent buildup of cluster S0 galaxies observed since $z \approx 0.4$.

Subject headings: galaxies: clusters: individual (CI 0024+1654) — galaxies: evolution — galaxies: spiral — galaxies: stellar content — ultraviolet: galaxies

1. INTRODUCTION

In normal spiral galaxies, the ultraviolet (UV) luminosity depends sensitively on the presence of two components: young, hot stars and dust. UV continuum observations are sensitive to star formation on timescales of less than a few times 100 Myr and can thus place strong constraints on both current and recent activity, particularly if independent estimates of dust extinction are available. Due to its wide field of view, the *Galaxy Evolution Explorer* (*GALEX*) satellite allows for an efficient census of the UV content of galaxies (Martin et al. 2005). Here we use *GALEX* imaging to address a key issue in galaxy evolution: the search for objects in morphological transition in intermediate-redshift clusters.

It has been known for many years that galaxies in rich clusters are generally bluer and more active at intermediate redshift (Butcher & Oemler 1978). With the aid of *Hubble Space Telescope* (*HST*) imaging, recent quantitative studies of the evolving morphology-density relation (Dressler et al. 1997; Smith et al. 2005; Postman et al. 2005) lend support to the suggestion that blue star-forming cluster spirals have been somehow transformed into red S0 galaxies (Couch & Sharples 1987). In order to understand how such transformations might occur, several spectroscopic and photometric surveys of intermediate-redshift clusters have sought to pinpoint samples of galaxies that are in the (presumed short-lived) *transitional phase* (e.g., Couch et al. 1998; Poggianti et al. 1999; Balogh et al. 1999).

Several candidate object classes have been identified from spectroscopic data utilizing the combination of [O II] emission

and Balmer absorption, which are respectively sensitive to ongoing star formation and recent star formation on timescales of ≈ 1 Gyr. These include the “E+A” or “k+a” galaxies that exhibit a deep Balmer H δ line (Dressler & Gunn 1983, 1992) and the “e(a)” galaxies that combine [O II] emission with strong Balmer absorption (Poggianti et al. 1999). Recent studies indicate that e(a) spectra are largely associated with dusty starburst galaxies (Poggianti & Wu 2000) and that (k+a) spectra are associated with the poststarburst remnants of a merger or close encounter (Goto 2005; Tran et al. 2003; but see also Poggianti et al. 2004). However, there are few intermediate-redshift S0 galaxies with e(a) or poststarburst spectral signatures (Poggianti et al. 1999). Indeed, Moran et al. (2005) found no difference between the stellar populations of E and S0 galaxies at $z = 0.4$ —an unexpected result if S0 galaxies were recently created from starbursting systems. Clearly, our understanding of the evolutionary link between e(a)/k+a galaxies and S0 galaxies remains incomplete.

Recently, a further interesting class has been identified—the so-called passive spirals (Couch et al. 1998; Dressler et al. 1999; Poggianti et al. 1999; Goto et al. 2003). These objects exhibit spiral morphology in *HST* images, but they reveal weak or no [O II] emission. Some authors have suggested that these are spiral-to-S0 transition objects for which the cessation of star formation occurs on a faster timescale than the transformation of spiral morphology. Such a delay in the morphological transformation would naturally explain the older stellar populations inferred from optical spectra of S0 galaxies (Poggianti et al. 1999; Moran et al. 2005). Theoretical models by Bekki et al. (2002) have shown that this scenario is consistent with gas “starvation,” an interaction with the intracluster medium that serves to inhibit star formation by halting the accretion of cold halo gas onto the galaxy disk (see Larson et al. 1980, Quilis et al. 2000, and Treu et al. 2003). Without further gas accretion, Bekki et al. (2002) found that the spiral arms fade within ≈ 3 Gyr.

Goto et al. (2003) find that passive spirals in the Sloan Digital

¹ California Institute of Technology, Department of Astronomy, Mail Code 105-24, Pasadena, CA 91125; smm@astro.caltech.edu, rse@astro.caltech.edu.

² Department of Physics, University of California, Santa Barbara, CA 93106; tt@physics.ucsb.edu.

³ University of California at Los Angeles, Department of Physics and Astronomy, Los Angeles, CA 90095.

⁴ School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

⁵ Laboratoire d’Astrophysique de Marseille, Traverse du Siphon, BP 8, 13376 Marseille Cedex 12, France.




FIG. 1.—Normalized co-added spectra of E+S0 galaxies (*lower spectrum*), passive (*middle spectrum*), and active (*upper spectrum*) spiral galaxies, offset vertically, in arbitrary units, created as in Moran et al. (2005). Key spectral lines are marked.

Sky Survey reside preferentially in intermediate-density environments, thus confirming this is a cluster-related phenomenon. Balmer H δ absorption is weaker in passive spirals than in the overall spiral population (Goto et al. 2003; Poggianti et al. 1999), suggesting that their stellar populations already resemble the older populations found in E+S0 (“early-type”) galaxies. To date, however, little else is known about the detailed star formation histories of passive spirals.

In this Letter, we combine *GALEX* UV imaging of Cl 0024+17, a rich cluster at $z = 0.4$, with *HST* imaging and Keck spectroscopy to constrain the recent and ongoing star formation rates (SFRs) of passive spirals. By comparing our measurements of UV-optical colors and key spectral lines with model star formation histories, we can for the first time distinguish between several explanations for the nature of the passive spirals.

2. OBSERVATIONS AND SAMPLE SELECTION

2.1. Data

We make use of *HST* imaging and Keck spectroscopy of Cl 0024+17 from the comprehensive wide-field survey fully described in Treu et al. (2003) and Moran et al. (2005). Briefly, the survey builds on a sparsely sampled mosaic of 39 WFPC2 images taken in the F814W filter ($\sim I$ band), providing coverage to a projected radius >5 Mpc. Treu et al. (2003) reported reliable morphological classifications to $I = 21.1$, and we combine these classifications with Keck spectroscopy in order to define three samples of cluster members: passive spiral, “active” or star-forming spiral, and E+S0.

Cl 0024+17 was observed for 13.3 ks with *GALEX* in 2004 October (GO program 22; Cycle 1; PI: T. Treu) in both near-(NUV) and far-ultraviolet (FUV) filters (Martin et al. 2005; Morrissey et al. 2005). As *GALEX*’s field of view is $\sim 1^\circ 2$, the images readily cover the full *HST* mosaic. The NUV band closely matches the rest-frame FUV at $z \approx 0.4$ ($\lambda_c \sim 1620$ Å), and so we focus on that band. Galaxy fluxes were measured within $6''$ circular apertures, centered on the optical position, and comparable to the measured NUV FWHM ($5''.5$). Such a fixed-size aperture makes the detection limits easier to interpret. We apply a fixed 20% aperture correction for agreement with SExtractor-derived total magnitudes (MAG_AUTO) and for comparison to MAG_AUTO magnitudes in F814W (Bertin & Arnouts 1996). We use the *kcorrect* software version 4.1.2 (Blanton et al. 2003) to convert observed NUV and F814W fluxes to rest-frame FUV and *V* luminosities. We assume a Galactic extinction of $E(B - V) = 0.056$ (Schlegel et al. 1998).

Observations with the DEIMOS (DEep Imaging Multi-Object Spectrograph) on Keck II from 2001 October to 2005 October secured spectra for over 500 members of Cl 0024+17 (300 with *HST* imaging). Details are, or will be, provided in Treu et al. (2003), Moran et al. (2005), and S. M. Moran et al. (2006, in preparation). We augment our sample with cluster members derived from the Canada-France-Hawaii Telescope (CFHT) survey of Czoske et al. (2001), who reported the strengths of (or the absence of) several spectral lines including [O II]. For both the Keck and CFHT spectra, we use Lick-style indices that are measured as described in Moran et al. (2005) and references therein.

2.2. Sample Selection

As faint emission line objects are easier to robustly identify than those without emission, there is a potential bias against identifying “passive” objects. To minimize this effect, we adopt a bright magnitude limit $F814W \leq 21.1$, where the sample is nearly unbiased (Moran et al. 2005). Reliable visual morphologies are available to the same limit (Treu et al. 2003).

We define as “passive” any Sa–Sd spirals that have $EW([O II]) > -5$ Å (Dressler et al. 1999; Poggianti et al. 1999), with “active” spirals having $EW([O II]) \leq -5$ Å; a more stringent limit is precluded by signal-to-noise ratio considerations. While this definition does not exclude (k+a) or e(a) spirals as separate classes, our Cl 0024 sample contains only two spirals with $EW(H\delta) \equiv H\delta_\lambda > 5$ Å, both with [O II] emission. To be consistent in comparing passive spirals with E+S0 galaxies, we likewise exclude all early-type galaxies with $EW([O II]) < -5$ Å ($\sim 25\%$). We manually examined all optical and UV galaxy images to ensure correct associations. We removed from the sample all galaxies where the UV flux is centered on a neighbor or is otherwise contaminated; this reduced the total sample by 12%. The final sample consists of 68 cluster spirals, of which 24 are passive and 44 active. The comparison sample of passive E+S0 galaxies contains 75 objects. In Figure 1, we show co-added, normalized spectra for each type.

3. UV EMISSION IN PASSIVE CLUSTER SPIRALS

While the optical spectra of passive spirals are noteworthy for their resemblance to those of red early-type galaxies (Fig. 1; Dressler et al. 1999), their FUV properties reveal a very different picture. A large fraction of passive spirals in Cl 0024 emit vigorously in the FUV. The fraction of UV-detected passive spirals ($>3\sigma$) is $67\% \pm 17\%$, not significantly different from that of active spirals ($80\% \pm 13\%$). However, only $25\% \pm 7\%$ of the early-type galaxies are similarly detected in the UV, clearly indicating that the stellar populations of passive spirals differ from those of the early-type galaxies.

Moreover, Figure 2 shows that the UV-detected passive spirals have $FUV - V$ colors intermediate between the bluer active spirals and early-type galaxies, which span a wide range extending the furthest to the red. A Kolmogorov-Smirnov test confirms that the $FUV - V$ colors of passive spirals differ from those of E+S0 galaxies with $>99\%$ confidence. These intermediate colors strongly suggest that passive spirals bridge the gap between the blue and red galaxy sequences. Below, we evaluate this suggestion by considering model star formation histories.

4. MODEL STAR FORMATION HISTORIES

The detection of strong FUV emission from passive spirals is perhaps surprising in a simple picture where the UV light and optical emission lines are generated simultaneously from




FIG. 2.—Distribution of rest-frame $(FUV - V)_{AB}$ colors for active spiral (blue histogram), passive spiral (hatched green histogram), and early-type (filled red histogram) galaxies.

the same star-forming regions. Here we explore possible explanations. The key to understanding the observation lies in the hierarchy of lifetimes of the stars responsible for the strong $H\delta$ absorption (10^9 yr), FUV flux (10^7 – 10^8 yr), and hard ionizing photons (10^7 yr).

It is useful at this point to introduce the $FUV - V$ versus $H\delta_A$ diagram, which we will use as our main diagnostic tool. First, by comparing signatures of star formation over different timescales, this diagram allows us to explore the recent star formation history of active and passive spirals and to assess whether they can be connected in an evolutionary sequence. Second, since $FUV - V$ is sensitive to dust extinction while $H\delta$ is not (e.g., MacArthur 2005; Sato & Martin 2006), this diagram allows us to explore the possibility that the redder $FUV - V$ colors of passive spirals with respect to active spirals could be due to dust instead of differences in the recent star formation history.

Figure 3 shows the location of passive and active spirals in the $FUV - V$ versus $H\delta_A$ diagram. Active spirals (open blue triangles) are clustered at a blue $FUV - V$ color with moderately strong Balmer absorption ($H\delta \sim 4 \text{ \AA}$), whereas the passive spirals (filled green triangles) extend to redder $FUV - V$ colors with weaker $H\delta$. The mean location of each population is marked as an error-barred point; errors indicate uncertainty in the mean. The error-barred point with the weakest $H\delta$ is a composite of 13 passive spirals that cannot be plotted individually, due to either a lack of reliable $H\delta$ or contamination of FUV flux. We co-added available spectra for these galaxies to measure $H\delta$, and we adopt the mean of available $FUV - V$. This composite lies near the locus of individually plotted passive spirals at low $H\delta$ and red $FUV - V$.

Evolutionary tracks from the population synthesis code of Bruzual & Charlot (2003) are overlaid in Figure 3. We consider only solar metallicity and only plot model points reached in less than the time from formation at $z \sim 5$ to observation at $z = 0.4$ (~ 8 Gyr). Tracks are in red for regimes where $EW([O II]) > -5 \text{ \AA}$; we use SFRs and L_b from the Bruzual & Charlot (2003) models to estimate $EW([O II])$ according to the relation in Barbaro & Poggianti (1997). We calculate $[O II]$ attenuation by dust following Calzetti et al. (1994), with nebular emission extinguished more strongly than the stellar continuum ($E_*/E_g \sim 0.5$), but we caution that this is only a representative model. We subtract 0.9 \AA from $H\delta$ for all model tracks to account for infalling by nebular emission (Barbaro & Poggianti 1997).

Solid lines represent exponentially declining SFRs. Those marked with characteristic timescales $\tau = 3, 5,$ and 10 Gyr have been found to reproduce the optical spectra of Sa-, Sb-, and Sc-type galaxies (“active” spirals), respectively (Poggianti




FIG. 3.—Rest-frame $(FUV - V)_{AB}$ color vs. $H\delta_A$, for passive (filled green triangles) and active (open blue triangles) spirals, with typical error at upper right. Arrows indicate lower limits on $FUV - V$, color-coded in the same manner. Black crosses indicate, from lower right to upper left, the mean location of the active spirals (with error on the mean), the same for passive spirals, and a composite of 13 passive spirals that cannot be plotted individually. Solid, dotted, dashed, and dash-dotted lines, respectively, mark several classes of model star formation histories, as labeled on the plot and discussed in the text. Tracks are colored red for regimes where $EW([O II]) > -5 \text{ \AA}$. Model ages increase generally from the lower right to the upper left.

& Barbaro 1996). Dashed lines include dust extinction of $A_V = 0.6$, corresponding to the mean difference in $FUV - V$ between the passive and active spirals, for a Cardelli et al. (1989) extinction law with $R_V = 3.1$. The lower dashed line is a representative $\tau = 5$ Gyr model with dust. The uppermost dashed and solid lines represent a $\tau = 1$ Gyr SFR (with and without dust) and will be discussed below. Active spirals largely occupy the region in between tracks with zero and moderate extinction.

While adding extinction to the active spiral models can serve to reproduce the lack of emission lines and the redder $FUV - V$ colors of the passive spirals, as expected it has very little effect on the $H\delta$ values. Thus, while a model with extinguished star formation is sufficient to explain those passive spirals with higher $H\delta$, it is unable to account for the bulk of the population observed with weaker $H\delta$. Conceivably, more complex dust screens could account for this deficiency, but we will explore a more straightforward explanation in which the SFR is suppressed as a result of environmental processes.

We first consider “truncated” star formation: in Figure 3, we plot (as dotted lines) several representative $\tau = 3$ – 10 Gyr models where SFR instantaneously falls to zero at an age between $t = 5$ and 7 Gyr, representing a range of initial formation ages. Such models eventually generate both lower $H\delta$ values and redder $FUV - V$ colors after star formation is turned off, but the UV emission decays too rapidly, falling below our GALEX detection limit well before low- $H\delta$ values are reached.

Only models with a relatively fast exponential decline in SFR ($\tau \leq 1$ Gyr, the upper solid and dashed lines in Fig. 3) can match the $FUV - V$ and $H\delta$ of the bulk of passive spirals. Alternatively, any of the active spiral models with $\tau = 3$ – 10 Gyr can migrate toward the $\tau = 1$ Gyr track if their star formation begins to

decline at a much steeper exponential rate (thick dash-dotted line in Fig. 3), entering into a spectroscopically passive phase. The two scenarios are essentially equivalent, differing only in the age of the galaxy when it enters the passive phase and in the amount of dust extinction applied.

5. DISCUSSION

Remarkably, the star formation histories with a sharp decline in SFR resemble those expected for spirals affected by “starvation” (Bekki et al. 2002; Larson et al. 1980; Quilis et al. 2000). As the accretion of gas is rapidly cut off, star formation decays on a short timescale as the remaining disk gas is used up ($0.8 \text{ Gyr} < \tau < 2 \text{ Gyr}$, according to Larson et al. 1980). More generally, we might expect that any “gentle” physical mechanism that inhibits star formation without immediately affecting the spiral morphology would behave similarly.

A gentle mechanism is also easier to reconcile with the observed homogeneity of the stellar population and dynamical properties of S0 galaxies (Treu et al. 2003; Moran et al. 2005). Models with no explicit break in the SFR, but with low values of τ ($\sim 1 \text{ Gyr}$), reproduce the passive spiral data as well as the starvation model, and they seem not to require the action of any physical mechanism. Yet passive spirals seem to be most prevalent in clusters (Goto et al. 2003), suggesting that a cluster-related mechanism contributes to their formation.

Since few passive spirals are observed with an FUV upper limit ($8/24$), we might expect that any morphological transition occurs soon after a passive spiral fades below our FUV limiting magnitude. For the model starvation track plotted as a thick dash-dotted line in Figure 3, the total lifetime from the halt of gas accretion to the point where the FUV luminosity fades below our detection limit is $\sim 3 \text{ Gyr}$, in good agreement with the morphological transformation timescale predicted by Bekki et al. (2002).

This particular evolutionary track spends $\sim 1.5 \text{ Gyr}$ in a phase with $\text{EW}([\text{O II}]) \geq -5.0 \text{ \AA}$ and detectable FUV (for the mean $V = 20.6$). The fraction of all bright spirals in this UV-detected, spectroscopically passive phase is $27\% \pm 7\%$ in Cl 0024, although some contamination is likely from galaxies under-

going dust-obscured star formation. These statistics enable us to consider whether the fraction of passive spirals observed is consistent with the buildup of the S0 fraction from $z = 0.4$ to $z = 0.0$.

Following Smith et al. (2005), we define the ratio of S0 to elliptical galaxies today, $N_{\text{S0}, z=0}/N_{\text{E}, z=0}$, in terms of the passive spiral fraction and S0 fraction at $z = 0.4$:

$$\frac{N_{\text{S0}, z=0}}{N_{\text{E}, z=0}} = \left(\frac{N_{\text{S0}, z=0.4}}{N_{\text{E}, z=0.4}} + \frac{\Delta t}{\tau} \frac{N_{\text{psp}, z=0.4}}{N_{\text{E}, z=0.4}} \right) \frac{N_{\text{E}, z=0.4}}{N_{\text{E}, z=0.4} + \Delta N_{\text{E}}}, \quad (1)$$

where Δt is the look-back time to $z = 0.4$, 4 Gyr , and τ is the passive spiral lifetime of 1.5 Gyr . The last term accounts for the buildup of elliptical galaxies through other means, which we set equal to 1 for this simple analysis. Taking $N_{\text{S0}, z=0.4}/N_{\text{E}, z=0.4} = 0.65$ (Dressler et al. 1997) and $N_{\text{psp}, z=0.4}/N_{\text{E}, z=0.4} = 0.25$,⁶ we calculate $N_{\text{S0}, z=0}/N_{\text{E}, z=0} = 1.3$, in rough agreement with $N_{\text{S0}, z=0}/N_{\text{E}, z=0} \sim 1.7 \pm 0.6$ from Dressler et al. (1997). This shows that passive spirals undergoing starvation can evolve into present-day S0 galaxies, consistent with the observed evolution of the morphology-density relation.

While starved spirals cannot build up the *entire* population of local S0 galaxies due to differences in the stellar mass functions of local S0 galaxies and spiral galaxies (Kodama & Smail 2001), our result provides strong evidence that the abundant passive spiral population at intermediate redshift is an important transition population that contributes to the present-day cluster S0 population. By combining the different postburst timescales probed by *GALEX* UV imaging and optical spectroscopy, we have delineated a duty cycle of gradual decline in activity consistent with the recent growth in the S0 population.

We thank L. MacArthur and I. Smail for valuable comments. We acknowledge financial support from NSF grant AST-0307859, STScI grants HST-GO-08559.01-A and HST-GO-09836.01-A, and NASA for *GALEX* Cycle 1 GO program 22.

⁶ According to Dressler et al. (1997), the ratio of elliptical galaxies to spiral galaxies is nearly unity in Cl 0024, and so $N_{\text{psp}, z=0.4}/N_{\text{E}, z=0.4} \approx N_{\text{psp}}/N_{\text{sp}}$.

REFERENCES

- Balogh, M. L., Morris, S. L., Yee, H. K. C., Carlberg, R. G., & Ellingson, E. 1999, *ApJ*, 527, 54
- Barbaro, G., & Poggianti, B. M. 1997, *A&A*, 324, 490
- Bekki, K., Couch, W. J., & Shioya, Y. 2002, *ApJ*, 577, 651
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Blanton, M. R., et al. 2003, *AJ*, 125, 2348
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Butcher, H., & Oemler, J., Jr. 1978, *ApJ*, 226, 559
- Calzetti, D., Kinney, A. L., & Storchi-Bergmann, T. 1994, *ApJ*, 429, 582
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Couch, W. J., Barger, A. J., Smail, I., Ellis, R. S., & Sharples, R. M. 1998, *ApJ*, 497, 188
- Couch, W. J., & Sharples, R. M. 1987, *MNRAS*, 229, 423
- Czoske, O., Kneib, J.-P., Soucail, G., Bridges, T. J., Mellier, Y., & Cuillandre, J.-C. 2001, *A&A*, 372, 391
- Dressler, A., & Gunn, J. E. 1983, *ApJ*, 270, 7
- . 1992, *ApJS*, 78, 1
- Dressler, A., Smail, I., Poggianti, B., Butcher, H., Couch, W. J., Ellis, R. S., & Oemler, A., Jr. 1999, *ApJS*, 122, 51
- Dressler, A., et al. 1997, *ApJ*, 490, 577
- Goto, T. 2005, *MNRAS*, 357, 937
- Goto, T., et al. 2003, *PASJ*, 55, 771
- Kodama, T., & Smail, I. 2001, *MNRAS*, 326, 637
- Larson, R. B., Tinsley, B. M., & Caldwell, C. N. 1980, *ApJ*, 237, 692
- MacArthur, L. A. 2005, *ApJ*, 623, 795
- Martin, D. C., et al. 2005, *ApJ*, 619, L1
- Moran, S. M., Ellis, R. S., Treu, T., Smith, G. P., Smail, I., Dressler, A., & Coil, A. L. 2005, *ApJ*, 634, 977
- Morrissey, P., et al. 2005, *ApJ*, 619, L7
- Poggianti, B., & Barbaro, G. 1996, *A&A*, 314, 379
- Poggianti, B. M., Bridges, T. J., Komiyama, Y., Yagi, M., Carter, D., Mobasher, B., Okamura, S., & Kashikawa, N. 2004, *ApJ*, 601, 197
- Poggianti, B. M., Smail, I., Dressler, A., Couch, W. J., Barger, A. J., Burtcher, H., Ellis, R. S., & Oemler, A., Jr. 1999, *ApJ*, 518, 576
- Poggianti, B. M., & Wu, H. 2000, *ApJ*, 529, 157
- Postman, M., et al. 2005, *ApJ*, 623, 721
- Quilis, V., Moore, B., & Bower, R. 2000, *Science*, 288, 1617
- Sato, T., & Martin, C. L. 2006, *ApJ*, submitted
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Smith, G. P., Treu, T., Ellis, R. S., Moran, S. M., & Dressler, A. 2005, *ApJ*, 620, 78
- Tran, K. H., Franx, M., Illingworth, G., Kelson, D. D., & van Dokkum, P. 2003, *ApJ*, 599, 865
- Treu, T., Ellis, R. S., Kneib, J.-P., Dressler, A., Smail, I., Czoske, O., Oemler, A., & Natarajan, P. 2003, *ApJ*, 591, 53