

Draft whole-genome sequences of 10 atypical enteropathogenic Escherichia coli strains isolated in Brazil

Santos, Fernanda F; Richardson, Emily J; Bryant, Jack A; Yamamoto, Denise; Elias, Waldir P; Gomes, Tânia A T; Henderson, Ian R

DOI:

[10.1128/MRA.01432-18](https://doi.org/10.1128/MRA.01432-18)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Santos, FF, Richardson, EJ, Bryant, JA, Yamamoto, D, Elias, WP, Gomes, TAT & Henderson, IR 2018, 'Draft whole-genome sequences of 10 atypical enteropathogenic Escherichia coli strains isolated in Brazil', *Microbiology resource announcements*, vol. 7, no. 22, e01432-18. <https://doi.org/10.1128/MRA.01432-18>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 12/03/2019

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.


Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.



Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Draft Whole-Genome Sequences of 10 Atypical Enteropathogenic *Escherichia coli* Strains Isolated in Brazil

Fernanda F. Santos,^a Emily J. Richardson,^b Jack A. Bryant,^b Denise Yamamoto,^{a*}  Waldir P. Elias,^c  Tânia A. T. Gomes,^a Ian R. Henderson^b

^aDepartamento de Microbiologia, Imunologia e Parasitologia, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil

^bInstitute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom

^cInstituto Butantan, Laboratório de Bacteriologia, São Paulo, Brazil

ABSTRACT The number of diarrhea cases caused by atypical enteropathogenic *Escherichia coli* (aEPEC) has been increasing worldwide. Here, we report the draft whole-genome sequences of 10 aEPEC strains isolated in Brazil. These sequences will provide an important source for future studies concerning aEPEC pathogenicity and genetic markers of potentially virulent strains.

Globally, diarrheal diseases are the second leading cause of death and the leading cause of malnutrition in children under 5 years old (1, 2). One of the most significant etiological agents of moderate-to-severe diarrhea is enteropathogenic *Escherichia coli* (EPEC), which is one of the diarrheagenic *E. coli* pathotypes. EPEC is subdivided into typical EPEC (tEPEC) and atypical EPEC (aEPEC) based on the presence of the EPEC adherence factor (EAF) plasmid in the former group and its absence in the latter group (3, 4).

The main pathogenic mechanism of both tEPEC and aEPEC is the formation of attaching and effacing (AE) lesions in the intestinal epithelium, in which intimate adherence between bacteria and host cells occurs (5–8). The genes required for the establishment of AE lesions are located on the locus of enterocyte effacement (LEE) island, which is approximately 35 kb long (9).

The number of cases of diarrhea caused by aEPEC has increased even in industrialized countries. This information reveals the need for a thorough knowledge of the genetic traits of aEPEC organisms and their mechanisms of pathogenicity in order to develop appropriate control strategies (4, 10, 11).

The 10 aEPEC strains we sequenced were isolated from feces of children under 5 years old during epidemiological studies carried out in São Paulo, Brazil (1985 to 1986 and 1989 to 1990) by the Laboratory of Microbiology of the Universidade Federal de São Paulo (UNIFESP) and in Salvador (2003 to 2004) by the Laboratory of Bacteriology of the Butantan Institute (12, 13). All the strains were stored at –80°C in lysogeny broth (LB) and 15% glycerol and were routinely cultured in LB for 18 to 20 h at 37°C.

DNA extraction was performed using the QIAamp DNA minikit (Qiagen) according to the manufacturer's instructions. DNA libraries from each aEPEC genome were prepared using the Nextera XT library prep kit (Illumina, San Diego, CA) and sequenced at MicrobesNG (University of Birmingham, UK) on the Illumina HiSeq 2500 platform using 2- × 250-bp paired-end reads, achieving between 34 and 157× depth (Table 1). Trimmomatic version 0.30 (14) was used to adapt and quality trim the reads with a sliding window quality cutoff of Q15. The *de novo* assembly and annotation of the genomes were performed using the software SPAdes version 3.9.1 (with the –careful flag) (15) and Prokka version 1.12 (16), respectively. The genome assembly metric, calculated using QUAST, and the number of

Received 17 October 2018 **Accepted** 13 November 2018 **Published** 6 December 2018

Citation Santos FF, Richardson EJ, Bryant JA, Yamamoto D, Elias WP, Gomes TAT, Henderson IR. 2018. Draft whole-genome sequences of 10 atypical enteropathogenic *Escherichia coli* strains isolated in Brazil. Microbiol Resour Announc 7:e01432-18. <https://doi.org/10.1128/MRA.01432-18>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2018 Santos et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Fernanda F. Santos, ff.santos@unifesp.br, or Tânia A. T. Gomes, tatg.amaral@unifesp.br.

* Present address: Denise Yamamoto, Universidade de Santo Amaro, Campus I, São Paulo, Brazil.

TABLE 1 Genome assembly metrics and number of CDS of 10 aEPEC genomes

Strain	Mean coverage (×)	No. of contigs	Largest contig (bp)	Total length (bp)	GC content (%)	GC content		L_{50} (contigs)	L_{75} (contigs)	No. of CDS	GenBank accession No.
						N_{50} (bp)	N_{75} (bp)				
1331-2	141.587	148	428,643	5,187,915	50.61	112,592	63,355	14	29	4,998	QYVD000000000
2012-1	78.7493	230	247,360	5,607,608	50.43	97,093	41,042	21	43	5,482	QYVC000000000
2531-13	157.673	265	242,288	5,055,689	50.59	73,162	36,314	21	45	4,843	QYVB000000000
3522-6	85.7236	144	317,272	4,730,622	50.57	97,356	51,903	15	32	4,494	QYVA000000000
3881-3	109.173	73	359,010	4,701,486	50.72	241,425	130,938	9	15	4,442	QYUZ000000000
3991-1	34.3279	239	196,577	4,812,795	50.79	73,367	32,807	22	46	4,569	QYUY000000000
4581-2	87.0432	78	406,594	4,696,360	50.41	136,790	95,011	11	20	4,382	QYVF000000000
4632-3	93.7625	174	288,111	4,855,364	50.6	90,810	36,963	16	38	4,617	QYUX000000000
BA2103	85.4008	252	285,340	5,412,725	50.48	102,510	48,630	18	36	5,231	QYUW000000000
BA4095	43.121	71	510,385	4,797,996	50.61	216,769	124,397	8	14	4,568	QYUV000000000

coding sequences (CDS) annotated in each aEPEC genome are shown in Table 1. Unless specified, all software was used with default settings.

These sequences will provide an important source for futures studies concerning aEPEC pathogenicity and genetic markers of potentially virulent strains. Furthermore, these data will permit comparative studies to be carried out with aEPEC strains isolated in Brazil and others isolated in different countries.

Data availability. The reads used for assembly of the 10 aEPEC genomes were deposited in the Sequence Read Archive (SRA) at the National Center for Biotechnology Information (NCBI) under the accession number [PRJNA490882](https://www.ncbi.nlm.nih.gov/sra/PRJNA490882), and the whole-genome shotgun sequences were deposited in the GenBank database under the accession numbers shown in Table 1. The versions described in this paper are the first versions.

ACKNOWLEDGMENTS

This study was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, grant number 420561/2016-1), Brazil (to T.A.T.G.), and by the Institute of Microbiology and Infection (IMI) at the University of Birmingham, United Kingdom. Genome sequencing was provided by MicrobesNG, which was supported by the BBSRC (grant number BB/L024209/1). F.F.S. was supported by a scholarship from CNPq (140443/2014-2).

REFERENCES

- United Nations Children's Fund (UNICEF)/World Health Organization (WHO). 2009. Diarrhoea: why children are still dying and what can be done. whqlibdoc.who.int/publications/2009/9789241598415_eng.pdf.
- World Health Organization (WHO). 2017. Diarrhoeal disease/fact sheet. <http://www.who.int/mediacentre/factsheets/fs330/en/>.
- Kaper JB. Defining EPEC. 1996. *Rev Microbiol* 27:130–133.
- Donnenberg MS, Finlay BB. 2013. Combating enteropathogenic *Escherichia coli* (EPEC) infections: the way forward. *Trends Microbiol* 21: 317–319. <https://doi.org/10.1016/j.tim.2013.05.003>.
- Rothbaum R, McAdams AJ, Giannella R, Partin JC. 1982. A clinicopathologic study of enterocyte-adherent *Escherichia coli*: a cause of protracted diarrhea in infants. *Gastroenterology* 83:441–454.
- Moon HW, Whipp SC, Argenzio RA, Levine MM, Giannella RA. 1983. Attaching and effacing activities of rabbit and human enteropathogenic *Escherichia coli* in pig and rabbit intestines. *Infect Immun* 41:1340–1351.
- Hernandes RT, Elias WP, Vieira MAM, Gomes TAT. 2009. An overview of atypical enteropathogenic *Escherichia coli*. *FEMS Microbiol Lett* 297: 137–149. <https://doi.org/10.1111/j.1574-6968.2009.01664.x>.
- Wong AR, Pearson JS, Bright MD, Munera D, Robinson KS, Lee SF, Frankel G, Hartland EL. 2011. Enteropathogenic and enterohaemorrhagic *Escherichia coli*: even more subversive elements. *Mol Microbiol* 80: 1420–1438. <https://doi.org/10.1111/j.1365-2958.2011.07661.x>.
- McDaniel TK, Jarvis KG, Donnenberg MS, Kaper JB. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc Natl Acad Sci U S A* 92:1664–1668. <https://doi.org/10.1073/pnas.92.5.1664>.
- Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. 2013. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev* 26:822–880. <https://doi.org/10.1128/CMR.00022-13>.
- Ingle DJ, Tauschek M, Edwards DJ, Hocking DM, Pickard DJ, Azzopardi KI, Amarasekera T, Bennett-Wood V, Pearson JS, Tamboura B, Antonio M, Ochieng JB, Oundo J, Mandomando I, Qureshi S, Ramamurthy T, Hossain A, Kotloff KL, Nataro JP, Dougan G, Levine MM, Robins-Browne RM, Holt KE. 2016. Evolution of atypical enteropathogenic *E. coli* by repeated acquisition of LEE pathogenicity island variants. *Nat Microbiol* 1:15010. <https://doi.org/10.1038/nmicrobiol.2015.10>.
- Gomes TA, Irino K, Girão DM, Girão VB, Guth BE, Vaz TM, Moreira FC, Chinarelli SH, Vieira MA. 2004. Emerging enteropathogenic *Escherichia coli* strains? *Emerg Infect Dis* 10:1851–1855. <https://doi.org/10.3201/eid1010.031093>.
- Abe CM, Trabulsi LR, Blanco J, Blanco M, Dahbi G, Blanco JE, Mora A, Franzolin MR, Taddei CR, Martinez MB, Piazza RM, Elias WP. 2009. Virulence features of atypical enteropathogenic *Escherichia coli* identified by the *eae*(+) EAF-negative *stx*(-) genetic profile. *Diagn Microbiol Infect Dis* 64:357–365. <https://doi.org/10.1016/j.diagmicrobio.2009.03.025>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.