

Xuanweioxylon damogouense sp. nov., a gymnosperm stem from the Lopingian (late Permian) of southwestern China and its systematic and paleoecological implications

Yang, Yang; Xiao-Yuan, He; Hilton, Jason; Zhou, Fu-Guang; Chen, Xin-Shi; Wang, Shi-Jun

DOI:

[10.1016/j.revpalbo.2019.06.012](https://doi.org/10.1016/j.revpalbo.2019.06.012)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Yang, Y, Xiao-Yuan, H, Hilton, J, Zhou, F-G, Chen, X-S & Wang, S-J 2019, 'Xuanweioxylon damogouense sp. nov., a gymnosperm stem from the Lopingian (late Permian) of southwestern China and its systematic and paleoecological implications', *Review of Palaeobotany and Palynology*, vol. 269, pp. 94-103.
<https://doi.org/10.1016/j.revpalbo.2019.06.012>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 24/06/2019

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

***Xuanweioxylon damogouense* sp. nov., a gymnosperm stem from the Lopingian (late Permian) of southwestern China and its systematic and palaeoecological implications**

Yang Yang^a, Xiao-Yuan He^{a,*}, Jason Hilton^b, Fu-Guang Zhao^a, Xin-Shi Chen^a,
Shi-Jun Wang^c

^a *Institute of Deep Time Terrestrial Ecology, Yunnan University, Kunming 650091, China*

^b *School of Geography, Earth and Environmental Sciences and Birmingham Institute of Forest Research, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK*

^c *State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China.*

* Corresponding author. E-mail address hexiaoy68@163.com (X.Y. He)

Keywords: Conifer, wood anatomy, palaeoclimate, false rings, Xuanwei Formation, volcaniclastic tuff

ABSTRACT

A new species of permineralized gymnosperm stem is described from volcanoclastic tuffs in Lopingian (late Permian) Xuanwei Formation from eastern Yunnan province, China. The stem comprises well-preserved tissues of the pith, primary and secondary xylem. Pith is divided into a thick-walled parenchymatous peripheral zone and an inner parenchymatous part. Primary xylem strands are numerous and indistinct with mesarch maturation. Secondary xylem is pycnoxylic with scalariform bordered pits on radial tracheid walls, with rays normally uniseriate or partly biseriate. Cross-field pits are mixed and including circular, elliptical and scalariform pitting. Comparisons indicate a close affinity with the contemporaneous plant *Xuanweioxylon scalariforme*, but differences with it lead to the erection of the new *Xuanweioxylon damogouense* sp. nov. We evaluate the systematic affinity of both species of *Xuanweioxylon* and conclude they represent conifers based on features of their wood anatomy. We consider the Xuanwei Formation includes at least four species of conifer, and show that *X. damogouense* possessed weak growth rings and false rings that suggest it grew in a climate with low seasonality and occasional periods of drought.

1. Introduction

Fossil woods are important in the geological record in providing valuable information to help determine the organization, structure and habit of woody plants in the past and also providing a key line of evidence for palaeoclimate analysis. In addition, fossil woods can also help determine the composition of vegetation in

geological history (Zhang et al., 2006; Wang et al., 2011; Wei et al., 2016). In China, approximately 60 species belonging to 35 genera of **gymnospermous** wood have been recorded from the Permian (Li and Taylor, 1998, 1999; Wang et al., 2003, 2009, 2011; Zhang et al., 2006, 2007; Hilton et al., 2009a,b; Seyfullah et al., 2009; Feng et al., 2010, 2011, 2012; Feng, 2012; He et al., 2013; Shi et al., 2014; Wei et al., 2016; Wan et al., 2014, 2016, 2017a,b,c). **Most of these accounts** come from the northern China palaeo-landmass, and with no more than 10 species to date identified from the southern China palaeo-landmass.

Here we described a new species of **gymnospermous** wood that was collected from the upper Permian of Yunnan Province, southwestern China. Through detailed comparisons, we assigned it into *Xuanweioxylon* (He et al., 2013) and **establish** the new species *X. damogouense*. The discovery of this new species not only complements the anatomical characteristics of *Xuanweioxylon*, but also provides information on the palaeoecology and diversity of conifers from the Xuanwei Formation in the run up to Permian/Triassic mass **extinction** (Neregato et al., 2016).

2. Material and methods

Two permineralized specimens (**n**umbered YNUPB11002 and YNUPB11003) were collected from the Xuanwei Formation (Lopingian; late Permian) of the Damogou coalfield and Qinyun coalfield, Fuyuan County, Yunnan Province (fig. 1). The Xuanwei Formation was deposited during the late Wuchipingian to

Changhsingian stages of the Permian period, approximately 260–251 Ma (Wang et al., 2011).

Both specimens were cut to reveal cross, longitudinal and tangential sections using a rock saw with a diamond blade. Cut surfaces were then then prepared by the acetate peel method (Galtier and Phillips, 1999). Grinding was undertaken with #400 and #600 grade carborundum. Ground surfaces were etched in 5% HCL to leave the organic contents exposed on the etched surface. Peels were mounted on glass slides with coverslips using Canada Balsam. Photography was undertaken on an Axio Imager A2, and Nikon D3X digital camera with AF–S Micro 105mm 1:2.8 GED lens. Scanning Electron Microscopy (SEM) was undertaken at Yunnan University. Figures were constructed in CorelDraw. The specimens, peels and slides are deposited in the Institute of Deep Time Terrestrial Ecology of Yunnan University, China.

3. Systematic palaeobotany

Phylum Coniferophyta (Serbet et al., 2010)

Genus *Xuanweioxylon* He, Wang, Hilton et Shao, 2013

Type species: *Xuanweioxylon scalariform* He, Wang, Hilton et Shao, 2013

Species *Xuanweioxylon damogouense* sp. nov. (Plates I–V)

Holotype: YNUPB11002

Paratype: YNUPB11003

Depository: Institute of Deep Time Terrestrial Ecology, Yunnan University,

Kunming, China.

Collecting locality: Damogou coalfield and Qinyun coalfield in Fuyuan County of Eastern Yunnan Province, China.

Stratigraphic horizon and age: Xuanwei Formation, Late Wuchiapingian to Changhsingian stages, Lopingian series, Permian period (Wang et al., 2011).

Etymology: The specific *epiphet damogouense* is derived from the fossil location of Damogou coalfield, Yunnan Province.

Diagnosis: Gymnosperm wood with pith divided into two parts: thick-walled parenchymatous peripheral zone and parenchymatous central part. Sclerenchymatous cells absent. Primary xylem elements numerous and indistinct, maturation mesarch. Secondary xylem pycnoxylic, mainly with uniseriate scalariform bordered pits on radial tracheid walls. “Leaf gaps” absent. Ray cells homogeneous, uniseriate and partially biseriate, 1–13 cells high. Cross-fields with 3–19 circular, oval, polygonal and scalariform bordered pits with the pores nearly horizontal.

4. Description

4.1. General features

The species is represented by two decorticated stems in which the pith, primary xylem and secondary xylem are preserved. The two specimens vary in size from which the larger specimen (YNUPB11002; Plates I–IV) has a preserved length of c.

65 mm and an incomplete diameter in cross section of c. 68×126 mm (Plate I, 1). The smaller specimen (YNUPB11003; Plates V–VI) is 40 mm long and c. 12×14 mm in diameter in cross section (Plate V, 1). According to the preserved extent of the large stem, it is estimated that its entire diameter may be more than 20 cm. In the following account, we describe the specimens separately in order to characterise each first before considering their variation.

4.2. Specimen YNUPB11002

This is the larger of the two specimens described.

4.2.1. Pith

The pith is oval in the cross section, c. 12×18 mm in diameter (Plate I, 1–2) and is divided into a peripheral zone and central parenchyma region (Plate I, 3). The width of the peripheral zone varies slightly at different levels and is absent in some areas due to poor preservation. Cells of the peripheral zone are normally thick-walled parenchymatous and circular, oval and polygonal in cross section (Plate I, 3–4). The diameter of the thick-walled parenchyma cells decreases gradually in cross section from the inner part where they are 115×142–127×137 µm to 41×46–45×53 µm in the outer part (Plate I, 3–4). Secretory cells are scattered across the peripheral zone (Plate I, 4) and are circular, oval and polygonal. The secretory cells usually contain dark-coloured contents and form a discontinuous band located close to the internal

parenchyma region of the pith. In longitudinal section, the thick-walled parenchyma cells are usually longitudinally elongated (Plate I, 5), and many of them have circular pits on the walls (Plate I, 8).

The central parenchymatous region of the pith is composed of parenchyma cells that are isodiametric, polygonal and oval in cross section (Plate I, 6). Parenchyma cells are from 43×44 – 44×50 μm to 152×183 – 177×185 μm in diameter in cross section. In the longitudinal section, parenchyma cells usually are polygonal and rectangular and somewhat longitudinally elongated, but some of them are square, circular or even horizontally elongated (Plate I, 7).

4.2.2. Primary xylem

In cross section, primary xylem strands are indistinct as they are small, low and are almost at the same level with the innermost secondary xylem (Plate II, 1–3). Primary xylem strands are numerous but the exact number is unknown due to incomplete preservation of the stem. The primary xylem is mesarch (Plate II, 3) and in some places double primary xylem strands can be seen (Plate ?, ?). In longitudinal section, there are helical and scalariform thickenings on the primary xylem tracheid walls (Plate II, 4).

4.2.3. Secondary xylem

The secondary xylem is pycnoxylic, consisting of rays and tracheids. Growth rings are very weakly developed (Plate III, 1); a few false rings occur where they are formed by bands of broken tracheid walls (Plate II, 5–6).

In cross section, tracheids are sub-circular, square, rectangular with tangential dimensions larger than radial dimensions, or flat with tangential width smaller than radial width (Plate III, 2). Tracheid diameters varies from 29×25 to 79×62 µm. Rays are normally uniseriate, occasionally partly biseriate. Ray cells are rectangular, 17–82 µm in radial direction and 10–34 µm in tangential direction. There are no pits in the horizontal wall of ray cells. Rays density is 3–10 rays per mm².

In radial section, tracheid end walls are blunt and occasionally straight. Pitting on the radial tracheid walls is normally uniseriate scalariform (Plate III, 3–6). The width of the bordered pits is c. 9–37 µm and their height is c. 4–8 µm. Pits are usually located in the middle part of the tracheid, covering about 1/3 to 5/6 of the tracheid width. Occasionally, there are uniseriate oval or circular (Plate III, 7), or bi- to multiseriate opposite (Plate III, 8–10) bordered pits on the radial tracheid walls. Pits are usually contiguous. Ray cells are rectangular with thin walls. The horizontal walls of ray cells are often slightly wavy, and tangential walls are vertical or oblique (Plate III, 11). The cross-field show circular, oval, polygonal and scalariform bordered pits (Plate IV, 1–6); oval bordered pits are dominant, followed by scalariform bordered pits, while circular and polygonal bordered pits are rare. When the cross-field is occupied only by oval bordered pits, they are small and crowded, numbering about 5–19 with nearly horizontal apertures, and 3×4 to 12×15 µm in diameter. When the

cross-field pits are occupied only by scalariform bordered pits, they number about 3–6, each with diameters varying from 3×13 to 8×33 μm . Circular and polygonal bordered pits are rare and they are randomly distributed in the cross-field.

In tangential section, tracheid end walls are apiculate or partly straight. Individual tracheids are bent to varying degrees, and ornamentation on the tangential tracheid walls is absent. Rays are normally uniseriate and some are partially biseriate (Plate III, 12, 13) and 1–13 (typically 1–7) cells high. The width or tangential diameter of ray cells is 23–56 μm (mean = 36 μm , $n = 50$), while their height ranges from 18–61 μm (mean = 42 μm). Pits are absent on the walls of ray cells.

4.3. The smaller stem (YNUPB11003)

Features of this specimen largely agree with those of the larger stem. This stem is c. 40 mm long and c. 12×14 mm in diameter in cross section (Plate V, 1). In cross section, the pith consists of oval or isodiametric parenchyma cells with some secretory cells. In the middle of the pith, there is a band composed of broken cell walls (Plate V, 2). In longitudinal section, parenchyma cells are usually longitudinally elongated (Plate V, 3, 4). The primary xylem is indistinct and mesarch (Plate V, 5, 6), and secondary xylem is pycnoxylic (Plate V, 7) with uniseriate or occasionally bi- to multiseriate scalariform bordered pits on the radial tracheid walls (Plate V, 8, 9). Rays are normally uniseriate and partly biseriate. The cross-field pits are well-preserved, with oval, circular and scalariform bordered pits (Plate V, 10–12).

5. Discussion

The two stems investigated have common features allowing them to be considered as a single species including indistinct primary xylem strands, absence of a “leaf gap” in the secondary xylem, scalariform bordered pits on secondary tracheid walls, and well developed pits in cross-field. The larger specimen has an estimated diameter of >20 cm if it was completely preserved, suggesting it was a trunk or a large lateral branch of a larger tree. However, the small specimen has a stem diameter only 12×14 mm and while the external tissues are absent, was either a small trunk or a lateral branch of a larger tree.

5.1. Comparisons

The genus *Xuanweioxylon* was established by He et al. (2013) based on permineralized gymnosperm stems collected from the Lopingian of Panxian district, Guizhou Province, China, a short distance from the present locality (Fig. 1). The main features of *Xuanweioxylon* are that it possesses a heterocellular pith, which consists of parenchyma cells, transfusion tissue and sclerotic cells, and is divided into central part of thin-walled parenchymatous cells and peripheral part of thick-walled parenchyma cells. In *X. salariforme* primary xylem strands are numerous, secondary xylem is pycnoxylic and possesses scalariform bordered pits on radial tracheid walls

throughout the secondary xylem. The features of the specimens described here conform to these characters and allow their assignment to *Xuanweioxylon*.

Xuanweioxylon is monotypic and includes the sole species *X. scalariforme*. Although our specimens are quite similar to *X. scalariforme*, there are obvious differences between them. Firstly, in the pith of our specimens lacks sclerenchymatous cells but they are present in *X. scalariforme*. Secondly, the primary xylem strand in our specimens is indistinct and the margin between pith periphery and xylem is smooth without any projections of the primary xylem strands (Plate II, 1–3; Plate V, 5, 6), while the primary xylem strand of *X. scalariforme* is distinct and forms a triangle that extends into the pith (Plate VI, 1–3). Thirdly, cells of the peripheral zone of the pith are isodiametric in our specimens while they are mostly radially elongated in *X. scalariforme*, especially in places where the primary xylem strand is present. Fourthly, our specimens lack a “leaf gap” in the secondary xylem (Plate VI, 4–6), but in *X. scalariforme* “leaf gaps” are well-developed and obvious. These differences indicate that the present specimens differ from *X. scalariforme*, leading to the establishment of *Xuanweioxylon damogouense* sp. nov.

As outlined above, *Xuanweioxylon damogouense* had a diameter of > 20 cm if it was completely preserved, suggesting it was a trunk or a large lateral branch of a larger tree. However, *Xuanweioxylon scalariforme* has a stem with the diameter of only 2.8 x 3.2 cm, being much smaller than *X. damogouense* sp. nov. and was probably a small tree or a lateral branch of a larger tree.

He et al. (2013) made a detailed comparison between *Xuanweioxylon* and other

gymnosperm stems with scalariform pits on secondary xylem tracheid walls; this information will not be repeated here as with the addition of a second species the same features still distinguish *Xuanweioxylon* from these taxa. Recently Wan et al. (2017c) described a gymnosperm wood, *Yangquanoxylon miscellum*, from the Upper Pennsylvanian–lower Permian Taiyuan Formation of Yangquan City, Shanxi Province, North China. *Y. miscellum* shows scalariform pits on secondary xylem tracheid walls and mixed pits in the cross-fields, however differing from *Xuanweioxylon*. While scalariform pits in *Y. miscellum* are only occasionally present, in *X. damogouense* the scalariform pits are present on nearly all secondary xylem tracheid walls, and the mixed pits in cross-fields are mainly bordered. The mixed pits in cross-field in *Y. miscellum* are mainly simple. Thus *Y. miscellum* is obviously different from *Xuanweioxylon*.

5.2. Affinity of *Xuanweioxylon damogouense*

He et al. (2013) discussed the systematic position of *X. scalariforme* and considered it to be a coniferophyte of uncertain systematic affinity. In the secondary xylem of our specimens, there are some bi- to multiseriate opposite, circular to elliptical pits on radial tracheid walls, and the pits in cross field possess nearly horizontal pores. This suggests that *Xuanweioxylon* is a conifer, rather than a cordaitalean plant in which pits on radial tracheid walls of secondary xylem are typically alternatively arranged and the pits in cross-field possess oblique pores

(Wang et al., 2003; Hilton et al., 2009a, b).

Previous palaeobotanical studies from the Xuanwei Formation have documented four species of conifer from compression/impression fossils. These comprise leaves of *Podozamites lanceolatus* (L. et H.) Braun and *P. permica* Zhao, *Ullmannia bronni* Goeppert and *Ull. cf. bronni* Goeppert (Zhao et al., 1980; Zhao, 1990; Tian et al., 1993). In addition, interpreting *Xuanweioxylon scalariforme* as a conifer (see above), four species of conifer wood have also been reported from the Xuanwei Formation. These comprise *Walchiopremnon gaoi* Tian et al. (1993), *Xuanweioxylon scalariforme* He et al. (2013), *X. damogouense* sp. nov. and *Parapodocarpoxylon huopuense* He (2013). At present, there is no anatomical or co-occurrence data to help associate any of these conifer species with other conifer species from the Xuanwei Formation to develop whole-plant relationships (Bateman and Hilton, 2010). Furthermore, taphonomic studies of the fossil plant-bearing tuffaceous sediments from the Xuanwei Formation remain in their infancy (e.g. Neregato *et al.*, 2016), also not offering additional evidence of whole-plant reconstructions (Bateman and Hilton, 2010). The available evidence from adpression/impression foliage species and permineralized wood species demonstrate that at least four species of conifer were present in the Xuanwei Formation.

5.3. Environmental implications

Growth rings refer to a growth layer of wood and bark in cross section of a trunk

286 that represents a layer of wood and bark produced during a growing period. Growth
287 rings are an important tool to help evaluate palaeoclimate and tree habit (e.g., Creber
288 and Chaloner, 1984; Ash and Creber, 1992; Schweingruber, 1992; Zhou and Jiang,
289 1994; Falcon-Lang et al., 2000a,b; Brea et al., 2008, 2011; Wan et al., 2017c).
290 Through the study of the growth rings, growth conditions from the region in which
291 the tree grew can be deduced including also information on season (Creber, 1977;
292 Wan et al., 2017c). Growth rings are often formed on trees when the seasons vary
293 significantly. During the early stages of the growth season, as the light intensifies and
294 temperature generally raises, cambial activities increases forming early or spring
295 wood that has larger diameter of tracheid with thinner walls (Creber and Chaloner,
296 1984; Zhou and Jiang, 1994). Early or spring wood has a lighter colour. By contrast,
297 in the late stages of a growth season as the light weakness and temperature generally
298 decreases, cambial activity slows and forms late or summer wood that has smaller
299 diameter of tracheids. These thicker cell walls produce a dark colour (Creber and
300 Chaloner, 1984; Zhou and Jiang, 1994). Such features can be readily identified in
301 permineralized fossil woods. Change in growth ring colour, cell thickness and cell
302 wall thickness can reflect fluctuations in dryness, temperature and other
303 environmental factors during the period of plant growth (Zhou and Jiang, 1994; Shi et
304 al., 2015, 2017). By contrast, absence or weak growth rings could indicate that the
305 environmental factors affecting the growth of plants in the growth period is small,
306 having less noticeable seasonality. In our specimens, the growth rings are very weakly
307 developed (Plate III, 1), which indicates that variation in environmental factors that

308 affected plant growth when it was alive **was small, thus, suggests** low seasonality.

309 Both specimens of *Xuanweioxylon damogouense* have faint false rings which are
310 normally 1–5 cells wide and discontinuous. False rings are formed by the short-term
311 environmental disturbance during the growing period of a tree. Disturbance to growth
312 can be caused by phenomena such as flood, pathogen or insect attack, drought, light
313 availability, wind or fire damage and temperature fluctuations (Creber and Chaloner,
314 1984; Weaver et al., 1997; Falcon-Lang, 2003; Wan et al., 2017c). During the Permian
315 period, the Southern China tectonic plate was located in the region between the
316 tropics and subtropics, with a seasonal rainforest developing under a temperate and
317 humid **climate** (Tian and Zhang, 1980; Shen, 1995; Tian and Wang, 1995; He et al.,
318 2013). The structure of the false rings in *X. damogouense* were most likely caused by
319 temporary drought during the growing period (Wan et al., 2017c). However,
320 considering the abundance of volcanic ash horizons in the lower member of the
321 Xuanwei Formation (Wang et al., 2011; Neregato et al. 2016), this could also
322 represent episodes of **volcanic activity influencing environmental conditions that in**
323 **turn disrupted plant growth, for instance by producing gaseous emissions or ash fall**
324 **events**. Detailed palaeoenvironmental analysis of the tuffaceous units in the Xuanwei
325 Formation are now required to further assess the causal mechanisms of climate
326 change during the deposition of the Xuanwei Formation.

327 328 **Acknowledgements**

This work was supported by the National Natural Science Foundation of China (Awards No. 41162001, 41472021, 41530101, U1702242) and the Strategic Priority Research Program of Chinese Academy of Sciences (XDB26000000). We thank R. Rößler and an anonymous reviewer for constructive reviews of the manuscript.

References

- Ash, S.R., Creber, G.T., 1992. Palaeoclimatic interpretation of the wood structures of the trees in the Chinle Formation (Upper Triassic), petrified Forest National Park, Arizona, USA. *Palaeogeog., Palaeoclimatol., Palaeoecol.* 96, 299–317.
- Bateman, R. M. and Hilton, J., 2009. Palaeobotanical systematics for the phylogenetic age: applying organ–species, form–species and phylogenetic species concepts in a framework of reconstructed fossil and extant whole–plants. *Taxon* 58, 1254–1280.
- Brea, M., Artabe, A., Luis., Spalletti, A., 2008. Ecological reconstruction of a mixed Middle Triassic forest from Argentina. *Alcheringa* 32, 365–393.
- Brea, M., Matheos, S.D., Raigemborn, M.S., Iglesias, A., Zucol, A.F., Prámparo, M., 2011. Paleoecology and paleoenvironments of Podocarp trees in the Ameghino petrified forest (Golfo San Jore Basin, Patagonia, Argentina): constraints for early Paleogene paleoclimate. *Geologica Acta* 9, 13–28.
- Creber, G.T., 1977. Tree ring: a natural data–storage system. *Biol. Rev.* 52, 349–381.
- Creber, G.T., Chaloner, W.G., 1984. Influence of environmental factors on the wood structure of living and fossil trees. *Bot. Rev.* 50, 357–448

351 Falcon-Lang, H.J., 2000a. The relationship between leaf longevity and growth ring
 352 markedness in modern conifer woods and its implications for palaeoclimatic
 353 studies. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 160, 317–328.

354 Falcon-Lang, H.J., 2000b. A method to distinguish between woods produced by
 355 evergreen and deciduous coniferopsids on the basis of growth ring anatomy: a
 356 new palaeoecological tool. *Palaeontology* 43, 785–793.

357 Falcon-Lang, H.J., 2003. Growth interruptions in silicified conifer woods from the
 358 Upper Cretaceous Two Medicine Formation, Montana, USA: Implications for
 359 palaeoclimate and dinosaur palaeoecology. *Palaeogeogr., Palaeoclimatol.,*
 360 *Palaeoecol.* 199, 299–314.

361 Feng, Z., Wang, J., Rößler, R., 2010. *Palaeoginkgoxylon zhoui*, a new ginkgophyte
 362 wood from the Guadalupian (Permian) of China and its evolutionary implications.
 363 *Rev. Palaeobot. Palynol.* 162, 146–158.

364 Feng, Z., Wang, J., Rößler, R., 2011. A unique gymnosperm from the latest Permian of
 365 China, and its ecophysiological implications. *Rev. Palaeobot. Palynol.* 165,
 366 27–40.

367 Feng, Z., Wang, J., Liu, L.J., Rößler, R., 2012. A novel coniferous tree trunk with
 368 septate pith from the Guadalupian (Permian) of China: ecological and
 369 evolutionary significance. *Int. J. Plant Sci.* 173, 835–848.

370 Feng, Z., 2012. *Ningxiaites specialis*, a new woody gymnosperm from the uppermost
 371 Permian of China. *Rev. Palaeobot. Palynol.* 181, 34–46.

372 Galtier J., Phillips T. L., 1999. The acetate peel technique. In: Jones, T. P., Rowe, N. P.

373 (Eds.), Fossil Plants and Spores: Modern Techniques. Geological Society of
 374 London, London, pp. 67–71.

375 He, J., 2013. Fossil conifers of Late Permian age in Panxian, southwestern China.
 376 Master Degree Thesis of Institute of Botany, the Chinese Academy of Sciences.

377 He, J., Wang, S.J., Hilton, J., Shao, L.Y., 2013. *Xuanweioxylon scalariforme* gen. et sp.
 378 nov.: novel Permian coniferophyte stems with scalariform bordered pitting on
 379 secondary xylem tracheids. Rev. Palaeobot. Palynol. 197, 152–165.

380 Hilton, J., Wang, S.J., Galtier, J., Bateman, R.M. 2009a. Cordaitalean seed plants from
 381 the early Permian of north China. II. Reconstruction of *Cordaixylon tianii*. Int. J.
 382 Plant Sci. 170, 400–418.

383 Hilton, J., Wang, S.J., Galtier, J., Bateman, R.M. 2009b. Cordaitalean seed plants
 384 from the early Permian of north China. □. Reconstruction of the *Shanxioxylon*
 385 *taiyuanense* plant. Int. J. Plant Sci. 170, 951–967.

386 Li, H.Q., Taylor, D.W., 1998. *Aculeovinea yunguiensis* gen. et sp. nov.
 387 (Gigantopteridales), a new taxon of Gigantopterid stem from the Upper Permian
 388 of Guizhou Province, China. Int. J. Plant Sci. 159, 1023–1033.

389 Li, H.Q., Taylor, D.W., 1999. Vessel-bearing stems of *Vasovinea tianii* gen. et sp. nov.
 390 (Gigantopteridales) from the Upper Permian of Guizhou Province, China. Amer.
 391 J. Bot. 86, 1563–1575.

392 Neregato, R., D’Apolito, C., Glasspool, I. J., Wang, S. J., Feng, L., Windslow, P., Lu,
 393 J., Shao, L., and Hilton, J. 2016. Palynological constraints on the provenance and
 394 stratigraphic range of a Lopingian (late Permian) inter-extinction floral

395 lagerstätte from the Xuanwei Formation, Guizhou Province, China. Int. J. Coal
 396 Geol. 62, 139–150.

397 Schweingruber, F.H., 1992. Annual growth rings and growth zones in woody plants in
 398 southern Australia. Iawa Journal 13, 359–379.

399 Seyfullah, L.J., Hilton, J., Wang, S.J., Galtier, J. 2009. Anatomically preserved
 400 pteridosperm stems and rachises from Permian Floras of China. Int. J. Plant Sci.
 401 170, 814–828.

402 Shen, G.L., 1995. Permian floras, In: Li, X.X. (Ed.), Fossil Floras of China Through
 403 the Geological Ages, English edition. Guangdong Science and Technology,
 404 Guangzhou, China, pp. 127–223.

405 Shi, X., Yu, J.X., Broutin, J., Pons, D., 2015. *Junggaropitys*, a new gymnosperm stem
 406 from the Middle–Late Triassic of Junggar Basin, northwest China, and its
 407 palaeoecological and palaeoclimatic implications. Rev. Palaeobot. Palynol. 223,
 408 10–20.

409 Shi, X., Yu, J.X., Broutin, J., Pons, D., Rossignol, C., Bourquin, S., Crasquin, S., Li,
 410 Q., Shu, W.C., 2017. *Turpanopitys taoshuyuanense* gen. et sp. nov., a novel
 411 woody branch discovered in Early Triassic deposits of the Turpan Basin,
 412 northwest China, and its palaeoecological and palaeoclimate implications.
 413 Palaeogeogr., Palaeoclimatol., Palaeoecol. 468, 314–326.

414 Shi, X., Yu, J.X., Li, H., Chi, H.F., Zhang, W., 2014. *Xinjiangoxylon* gen. nov., a new
 415 gymnosperm from the latest Permian of China. Acta Geologica Sin. (English
 416 Edition) 88, 1356–1363.

417 Tian, B.L., Zhang, L.W., 1980. Fossil atlas of Wangjiazhai Mine Region in Shuicheng,
 418 Guizhou Province. China Coal Industry Publishing House, Beijing (in Chinese).
 419 Tian, B.L., Wang, S.J., 1995. Palaeozoic coal ball floras in China, In: Li, X.X. (Ed.),
 420 Fossil Floras of China through the geological ages, English edition. Guangdong
 421 Science and Technology, Guangzhou, pp. 202–211.
 422 Tian, B.L., Hu, T., Zhao, H., 1996. The first discovery of *Walchiopremnon gaoi* sp.
 423 nov. in China. In: Geological Section of Beijing Graduate School, China
 424 University of Mining and Technology (Eds.), Selected papers on coal geology.
 425 China Coal Industry Publishing House, Beijing, pp. 118–125 (in Chinese with
 426 English abstract).
 427 Wan, M.L., Yang, W., Wang, J., 2014. *Septomedullopitys szei* sp. nov., a new
 428 gymnospermous wood from Lower Wuchiapingian (Upper Permian) continental
 429 deposits of NW China, and its implication for a weakly seasonal humid climate
 430 in mid–latitude NE Pangaea. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 407,
 431 1–13.
 432 Wan, M.L., Zhou, W.M., Yang, W., Wang, J., 2016. Charred wood of *Prototaxoxylon*
 433 from the Wuchiapingian Wutonggou Formation (Permian) of Dalongkou,
 434 northern Bogda Mountains, northwestern China. *Palaeoworld* 25, 21–31.
 435 Wan, M.L., Yang, W., He, X.Z., Liu, L.J., Wang, J., 2017a. First record of fossil
 436 basidiomycete clamp connections in cordaitalean stems from the
 437 Asselian–Sakmarian (Lower Permian) of Shanxi Province, north China.
 438 *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 466, 353–360.

439 Wan, M.L., Yang, W., Liu, L.J., Wang, J., 2017b. *Ductoagathoxylon jimsarensis* gen.
 440 nov. et sp. nov., a gymnospermous stem from the Wuchiapingian (upper Permian)
 441 Wutonggou Formation in the Junggar Basin, northern Bogda Mountains,
 442 northwestern China. Rev. Palaeobot. Palynol. 241, 13–25.

443 Wan, M.L., Yang, W., He, X.Z., Zhou, W.M., Liu, L.J., Wang, J., 2017c.
 444 *Yangquanoxylon miscellum* gen. nov. et sp. nov., a gymnospermous wood from
 445 the Upper Pennsylvanian–Lower Permian Taiyuan Formation of Yangquan City,
 446 Shanxi Province, with reference to the palaeoclimate in north China.
 447 Palaeogeogr., Palaeoclimatol., Palaeoecol. 479, 115–125.

448 Wang, H., Shao, L.Y., Hao, L.M., Zhang, P.F., Glasspool, I.J., Wheelley, J.R., Wignall,
 449 P.B., Yi, T.S., Zhang, M.W., Hilton, J., 2011. Sedimentology and sequence
 450 stratigraphy of the Lopingian (Late Permian) coal measures in southwestern
 451 China. Int. J. Coal Geol. 85, 168–183.

452 Wang, S.J., Hilton, J., Tian, B.L., Galtier, J., 2003. Cordaitalean seed plants from the
 453 Early Permian of North China. I. Delimitation and reconstruction of the
 454 *Shanxioxylon sinense* plant. Int. J. Plant Sci. 164, 89–112.

455 Wang, S.J., Sun, K.Q., Cui, J.Z., Ma, S.M., 2009. Fossil plants from coal balls in
 456 China. In: Cui, J.Z. (Ed.), Fossil Flora of China. Higher Education Press, Beijing,
 457 China.

458 Wang, S.J., He, X.Y., Shao, L.Y., 2011. Cycad wood from the Lopingian (late Permian)
 459 of southern China: *Shuichengoxylon tianii* gen. et sp. nov. Int. J. Plant Sci. 172,
 460 725–734.

461 Weaver, L., McLoughlin, S., Drinnan, A.N., 1997. Fossil woods from the Upper
 462 Permian Bainmedart Coal Measures, northern Prince Charles Mountains, east
 463 Antarctica. J. Australian Geol. Geophys. 16, 655–676.

464 Wei, X.X., Zhang, X.H., Shi, G.R., Zhao, X.M., Huang, X., Luan, T.F., 2016. First
 465 report of a phytogeographically mixed (transitional) Middle–Late Permian fossil
 466 wood assemblage from the Hami area, northwest China, and implications for
 467 Permian phytogeographical, paleogeographical and paleoclimatic evolution in
 468 central Asia. Palaeogeogr., Palaeoclimatol., Palaeoecol. 448, 125–140.

469 Yao, Z.Q., Liu, L.J., Rothwell, G.W., Mapes, G., 2000. *Szecladia* new genus, a Late
 470 Permian conifer with multiveined leaves from South China. J. Paleontol. 74,
 471 524–531.

472 Zhang, W., Li, Y., Zhen, S.L., Li, N., Wang, Y.D., Yang, X.J., Yang, J.J., Yi, T.M., Fu,
 473 X.P., 2006. Chinese Fossil Wood. Chinese Forestry Press, Beijing China.

474 Zhang, W., Wang, Y.D., Zheng, S.L., Yang, X.J., Li, Y., Fu, X.P., Nan Li., 2007.
 475 Taxonomic investigations on permineralized conifer woods from the Late
 476 Paleozoic Angaran deposits of northeastern Inner Mongolia, China and their
 477 palaeoclimatic significance. Rev. Palaeobot. Palynol. 144, 261–285.

478 Zhao, H., 1990. The Permian flora of eastern Yunnan and the division and correlation
 479 of Xuanwei Formation. Master Degree Thesis of China University of Mining and
 480 Technology (Beijing)

481 Zhao, X. H., Mo, Z. G., Zhang, S. Z., Yao, Z. Q., 1980. Late Permian flora from
 482 western Guizhou and eastern Yunnan. In: Nanjing Institute of Geology and

483 Palaeontology, Academia Sinica (Ed.), Late Permian Coal Bearing Strata and
484 Biota from Western Guizhou and Eastern Yunnan. Science Press, Beijing, pp.
485 70–122 (In Chinese).

486 Zhou, Y., Jiang, X.M., 1994. Wood anatomy and ultrastructure of gymnospermous
487 woods in China. China Forestry Publishing House.

488

Figure and plate captions

Fig. 1. Left—Outline map of China showing province positions and with box showing border area of NE Yunnan Province and SW Guizhou Province. Right—enlargement of box area to show collection locality in black square (Figure modified from He et al., 2013).

Plate I. *Xuanweioxylon damogouense* sp. nov. from the Lopingian of southwestern China. **Holotype**, YNUPB11002. 1–4, 6 in cross section, 5, 7, 8 in radial section. 1. Tissue of the pith (P) and secondary xylem (SX). Area in the box is enlarged in Plate II, 5. Bar = 1 cm. Slide YH–0100. 2. Pith with **central** part (**CP**) and peripheral **zone** (**PZ**). Bar = 2 mm. Slide YH–0100. 3. Pith **central** part (**CP**) and peripheral **zone** (**PZ**). Bar = 200 µm. Slide YH–0101. 4. Secretory cells in the peripheral **zone** of pith. Bar = 200 µm. Slide YH–0101. 5. Longitudinally elongated thick-walled parenchyma cells of the peripheral **zone** of pith. Bar = 100 µm. Slide YH–0104. 6. Parenchyma cells in the **central** part of pith and the residual cell walls (arrows). Bar = 500 µm. Slide YH–0101. 7. Parenchyma cells of the **central** part of pith and the residual cells wall (arrows). Bar = 500 µm. Slide YH–0105. 8. Thick-walled parenchyma cells of the peripheral **zone** of pith with **circular** pits (arrow) on the walls. Bar = 20 µm. Slide YH–0104

Plate II. *Xuanweioxylon damogouense* sp. nov. from the Lopingian of southwestern

511 China. **Holotype** in **cross** section except 4 that is in radial section. 1. **Three indistinct**
512 **primary xylem strands** (arrows). Bar = 100µm. Slide YH-0100. 2. Indistinct mesarch
513 primary xylem **strand** (arrow). Bar = 50 µm. Slide YH-0100. 3. Mesarch primary
514 xylem **strand** (arrow). Bar = 50 µm. Slide YH-0102. 4. Helical and scalariform
515 thickenings on tracheid walls of the primary xylem. Bar = 20 µm. Slide YH-0104. 5.
516 Discontinuous false rings in the box **indicated at** Plate I, 1. Bar = 2 mm. Slide
517 YH-0100. 6. A false ring. Bar = 100 µm. Slide YH-0100.

518

519 **Plate III.** *Xuanweioxylon damogouense* sp. nov. from the Lopingian of southwestern
520 China. **Holotype**, 1, 2, 12 and 13 in **cross** section, 3-11 in radial section. 1.
521 Discontinuous and inconspicuous growth ring (arrow). Bar = 2 mm. Slide YH-0101.
522 2. Tracheids and uniseriate rays of secondary xylem. Bar = 100 µm. Slide YH-0103. 3.
523 Uniseriate scalariform bordered pits on the radial tracheid walls of secondary xylem.
524 Bar = 20 µm. Slide YH-0104. 4. SEM **image** of the secondary xylem showing
525 uniseriate scalariform bordered pits. Bar = 100 µm. 5-6. SEM photos of the secondary
526 xylem, showing uniseriate scalariform bordered pits. Bars = 50 µm. 7. Uniseriate oval
527 bordered pits on **radial** tracheid walls. Bar = 20 µm. Slide YH-0104. 8. Uniseriate to
528 multiseriate scalariform bordered pits and oval bordered pits on tracheid walls. Bar
529 =10 µm. Slide YH-0111. 9. Uniseriate to triseriate scalariform bordered pits on
530 tracheid walls. Bar = 10 µm. Slide YH-0106. 10. Uniseriate to triseriate scalariform
531 bordered pits on tracheid walls. Bar = 10 µm. Slide YH-0132. 11. Parenchyma cells
532 of the ray. Bar = 50 µm. Slide YH-0106. 12. Uniseriate rays. Bar = 200 µm. Slide

533 YH-0107. 13. Uniseriate and partly biseriate rays. Bar = 200 µm. Slide YH-0107.

534

535 **Plate IV.** *Xuanweioxylon damogouense* sp. nov. from the Lopingian of southwestern
536 China. **Holotype** in radial section. 1–6. Variation in cross-field pits. Bar = 20 µm.
537 Slide YH-0104.

538

539 **Plate V.** *Xuanweioxylon damogouense* sp. nov. from the Lopingian of southwestern
540 China. **Paratype** with 1, 2, 5–7 in **cross** section 3, 4, and 8 in radial section. 1. Tissue
541 overview including pith (P) and secondary xylem (SX). Bar = 1 mm. Slide YH-0240.
542 2. Pith cells. Bar = 1 mm. Slide YH-0240. 3. Longitudinally elongated thick-walled
543 parenchyma cells of pith's peripheral **zone**. Bar = 40µm. Slide YH-0241. 4.
544 Parenchyma cells of the pith. Bar = 40 µm. Slide YH-0241. 5. Indistinct primary
545 xylem **strand** (arrows). Bar = 200 µm. Slide YH-0240. 6. Mesarch primary xylem
546 **strand** (arrow). Bar = 50 µm. Slide YH-0240. 7. Tracheids of the secondary xylem.
547 Bar = 40 µm. Slide YH-0240. 8. Uniseriate scalariform bordered pits on radial
548 tracheid walls of secondary xylem. Bar = 20 µm. Slide YH-0241. 9. Uni- to triseriate
549 scalariform bordered pits. Bar = 20µm. Slide YH-0241. 10–12. Cross-field bordered
550 pits. Bars = 20 µm. Slide YH-0241.

551

552 **Plate VI.** *Xuanweioxylon scalariforme* He et al. from the Lopingian of southwestern
553 China. All Bars = 200 µm unless otherwise stated. 1. Primary xylem **strand** (arrows).
554 Slide WP2L-0077. 2. Primary xylem **strand** (arrows). Slide WP2L-0076. 3. Primary

555 xylem **strands** (arrows). Slide WP2–0085. 4. “Leaf gap” in the secondary xylem. Bar
556 = 100 μm . Slide WP2–0085. 5. “Leaf gap” in the secondary xylem. Slide WP2L–0076.
557 6. “Leaf gap” in the secondary xylem. Slide WP2–0085.