

KOI-3890

Kuszelewicz, James S.; North, Thomas S. H.; Chaplin, William J.; Bieryla, Allyson; Latham, David W.; Miglio, Andrea; Bell, Keaton J.; Davies, Guy R.; Hekker, Saskia; Campante, Tiago L.; Deheuvels, Sebastien; Lund, Mikkel N.

DOI:

[10.1093/mnras/stz1185](https://doi.org/10.1093/mnras/stz1185)

License:

None: All rights reserved

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Kuszelewicz, JS, North, TSH, Chaplin, WJ, Bieryla, A, Latham, DW, Miglio, A, Bell, KJ, Davies, GR, Hekker, S, Campante, TL, Deheuvels, S & Lund, MN 2019, 'KOI-3890: A high mass-ratio asteroseismic red-giant+M-dwarf eclipsing binary undergoing heartbeat tidal interactions', *Monthly Notices of the Royal Astronomical Society*, vol. 487, no. 1, pp. 14–23. <https://doi.org/10.1093/mnras/stz1185>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 11/07/2019

This article has been accepted for publication in *Monthly Notices of the Royal Astronomical Society* ©: 2019 The Author(s). Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved. doi:10.1093/mnras/stz1185

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

KOI-3890: a high-mass-ratio asteroseismic red giant+M-dwarf eclipsing binary undergoing heartbeat tidal interactions

James S. Kuszlewicz^{1,2,3★}, Thomas S. H. North^{2,3}, William J. Chaplin^{2,3}, Allyson Bieryla⁴, David W. Latham⁴, Andrea Miglio^{1,2,3}, Keaton J. Bell^{1,3}, Guy R. Davies^{2,3}, Saskia Hekker^{1,3}, Tiago L. Campante^{1,3}, Sebastien Deheuvels^{7,8} and Mikkel N. Lund^{1,2,3}

¹Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, D-37077 Göttingen, Germany

²School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

³Stellar Astrophysics Centre (SAC), Department of Physics and Astronomy, Aarhus University, Ny Muegade 120, DK-8000 Aarhus C, Denmark

⁴Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

⁵Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, Rua das Estrelas, P-4150-762 Porto, Portugal

⁶Departamento de Física e Astronomia, Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre, s/n, P-4169-007 Porto, Portugal

⁷Université de Toulouse, UPS-OMP, IRAP, F-31400 Toulouse, France

⁸CNRS, IRAP, 14, avenue Edouard Belin, F-31400 Toulouse, France

Accepted 2019 April 17. Received 2019 March 28; in original form 2018 November 26

ABSTRACT

KOI-3890 is a highly eccentric, 153-d period eclipsing, single-lined spectroscopic binary system containing a red giant star showing solar-like oscillations alongside tidal interactions. The combination of transit photometry, radial velocity observations, and asteroseismology has enabled the detailed characterization of both the red giant primary and the M-dwarf companion, along with the tidal interaction and the geometry of the system. The stellar parameters of the red giant primary are determined through the use of asteroseismology and grid-based modelling to give a mass and radius of $M_{\star} = 1.04 \pm 0.06 M_{\odot}$ and $R_{\star} = 5.8 \pm 0.2 R_{\odot}$, respectively. When combined with transit photometry, the M-dwarf companion is found to have a mass and radius of $M_c = 0.23 \pm 0.01 M_{\odot}$ and $R_c = 0.256 \pm 0.007 R_{\odot}$. Moreover, through asteroseismology we constrain the age of the system through the red giant primary to be $9.1^{+2.4}_{-1.7}$ Gyr. This provides a constraint on the age of the M-dwarf secondary, which is difficult to do for other M-dwarf binary systems. In addition, the asteroseismic analysis yields an estimate of the inclination angle of the rotation axis of the red giant star of $i = 87.6^{+2.4}_{-1.2}$ degrees. The obliquity of the system – the angle between the stellar rotation axis and the angle normal to the orbital plane – is also derived to give $\psi = 4.2^{+2.1}_{-4.2}$ degrees, showing that the system is consistent with alignment. We observe no radius inflation in the M-dwarf companion when compared to current low-mass stellar models.

Key words: asteroseismology – techniques: photometric – binaries: eclipsing – stars: evolution – stars: fundamental parameters.

1 INTRODUCTION

Over the past decade, there has been a space-based revolution courtesy of the *Kepler* (Borucki et al. 2010) and *CoRoT* (Baglin et al. 2006) missions. The study of solar-like oscillations – those excited and damped by near-surface turbulent convection – had long been confined to the Sun save a few rare cases, for example Arcturus (Smith, McMillan & Merline 1987), Procyon (Brown et al. 1991;

Arentoft et al. 2008; Bedding et al. 2010), and α CenB (Kjeldsen et al. 2005). The long-baseline, high-quality data sets available as a result of these space missions have resulted in a golden age for the field, with oscillations detected in hundreds of solar-type stars (e.g. Chaplin et al. 2014) and thousands of red giants (e.g. De Ridder et al. 2009; Hekker et al. 2009, 2011; Mathur et al. 2016; Yu et al. 2018).

Eclipsing binary systems in which the primary component is a red giant star showing solar-like oscillations are relatively rare. Asteroseismology gives an unprecedented opportunity to better characterize these systems (Hekker et al. 2010; Frandsen et al.

★ E-mail: kuszlewicz@mps.mpg.de

2013; Gaulme et al. 2013, 2014, 2016; Beck et al. 2014; Miglio et al. 2014; Brogaard et al. 2016; Rawls et al. 2016; Benbakoura et al. 2017; Brogaard et al. 2018; Themeßl et al. 2018). The detection and characterization of stellar oscillations not only allow inferences to be made about the internal processes of stars (Mosser et al. 2012; Montalbán et al. 2013; Eggenberger et al. 2017; Hekker, Elsworth & Angelou 2018), but also allow access to other properties of the stars such as ages (e.g. Soderblom 2010; Miglio et al. 2013; Silva Aguirre et al. 2018) and the stellar inclination angle with respect to our line of sight (e.g. Huber et al. 2013; Lund et al. 2014; Campante et al. 2016; Kamiaka, Benomar & Suto 2018; Kuszlewicz et al. submitted and references therein).

Amongst the many stars observed with *Kepler*, a new class of eccentric, ellipsoidal binary systems was discovered that showed photometric tidal distortions (Welsh et al. 2011; Thompson et al. 2012). The distinct ellipsoidal variations seen in the light curve are a result of the large tidal distortions of the surface of the stars during periastron, which leads to their name, ‘heartbeat’ stars. The theory behind these tidal distortions was derived by Kumar, Ao & Quataert (1995) and it was shown that the morphology of these distortions can lead to constraints on the inclination of the orbit, eccentricity, and argument of periastron without the need for an eclipse to be observed.

It is possible to find heartbeat systems where one component is an oscillating red giant. Beck et al. (2014) analysed 18 such systems where the primary star was a red giant showing solar-like oscillations, of which only three were eclipsing and two had determined mass fractions. Heartbeat systems that show solar-like oscillations are of considerable value because, when combined with possible radial velocity data, they can be characterized in great detail. Since they provide constraints on the eccentricity and inclination of the orbit, even when the system is not eclipsing these important parameters can be derived. If the systems are also eclipsing this can help improve constraints even more. This is particularly the case as there is a tendency for heartbeat stars to have low-mass companions, and so obtaining radial velocities for each component in the system is difficult. Having another means to infer the properties of the primary star, such as asteroseismology, is invaluable since this can propagate through to tighter constraints on the secondary star.

In addition to observing heartbeat systems where one component was an oscillating red giant, one star (KIC 5006817) in the sample of Beck et al. (2014) was identified as being a red giant+M-dwarf binary system. However, unlike the system in this work, it was not an eclipsing system and the orbital parameters of the system were instead inferred from the heartbeat signal. Gaulme et al. (2016) had three such red giant+M-dwarf systems in their sample, all of which were eclipsing, and used the asteroseismic scaling relations to derive the physical properties of the primary and secondary components. In this work, we are the first to apply grid-based modelling to these types of red giant+M-dwarf systems with an oscillating component that are both eclipsing and show heartbeat signals.

Precise constraints on the inclination angle of the orbit in an eclipsing (or transiting) system through the analysis of the heartbeat signal can be combined with knowledge about the inclination angle of the star (through, e.g., asteroseismology; Pesnell 1985; Gizon & Solanki 2003) to constrain the obliquity of the system.¹

The obliquity is an important parameter for understanding binary systems because it can help shed light on formation mechanisms (e.g. Bonnell et al. 1992; Bate, Lodato & Pringle 2010) and the dynamical evolution of the systems (Mazeh & Shaham 1979). The most extensive investigation into the obliquities of eclipsing binary stars has been performed by the BANANAs project (Albrecht et al. 2007, 2009, 2011, 2013, 2014). During these investigations, five systems were studied using the Rossiter–MacLaughlin effect (see Winn 2007 or Gaudi & Winn 2007 for an overview) to derive the sky-projected obliquity² of each star in the binary. All of the binaries analysed were close-in, with orbital periods in the range of ~ 6 –16 d, of which two are misaligned (CV Veturum and DI Herculis) and the others are consistent with alignment. Heartbeat systems, on the other hand, can have orbital periods larger than ~ 100 d, and their companions tend to be low-mass (and so they are likely to be single-lined spectroscopic systems), and any measurement of the Rossiter–MacLaughlin effect would be very difficult, if not unobtainable, with current instruments. Therefore, to determine the obliquity, the system needs to be eclipsing and the inclination angle of the star must have a measurement. In the case where the primary is an oscillating red giant this becomes possible through asteroseismology.

In this work we aim to derive the properties of the components of an eclipsing binary system through the use of asteroseismology, eclipse fitting, and radial velocity analysis. In addition, we aim to constrain the geometry of the system by inferring the inclination angle of the red giant primary to then obtain the obliquity of the system, which gives information as to whether the system is aligned.

2 OBSERVATIONS

KOI-3890³ (KIC 8564976) was initially identified as a possible evolved-host planetary system based on *Kepler* data using the NASA Exoplanet Archive (Akeson et al. 2013). When the transit depth (from the archive) was combined with the stellar radius, the resulting radius for the secondary clearly indicated the object is a stellar companion. The potential binary star nature of this system was first suggested in Lillo-Box et al. (2015), who collected 22 radial velocities using the CAFE spectrograph (Aceituno et al. 2013). While the phase coverage in Lillo-Box et al. (2015) is limited, the authors place constraints on the minimum eccentricity $e \geq 0.33$ and minimum radial velocity amplitude $K \geq 2.5 \text{ km s}^{-1}$, which is equivalent to a required minimum companion mass of $M > 0.0097 \pm 0.0014 M_{\odot}$. This minimum mass is equivalent to $M > 10.2 M_{\text{J}}$, and whilst this theoretically allows for a planet-mass object ($M \lesssim 13 M_{\text{J}}$; Burrows et al. 2001) this is more likely to be a binary star system.

KOI-3890 was observed near-continuously for the 4-yr duration of the *Kepler* mission in long cadence (~ 30 min) mode. *Kepler* detected a transit-like feature at a period of 152.8 d, and the system was flagged as a KOI. The long cadence (~ 30 min) *Kepler* light curve was downloaded from the Mikulski Archive for Space Telescopes (MAST)⁴ and detrended with a moving median filter with a width of 20 d to ensure the eclipses were unaffected. The light curve was also clipped at the 4σ level about the moving median.

²The sky-projected obliquity, λ , is defined as the angle between the projections of the orbital and rotation axes on the sky.

³KOI – *Kepler* object of interest.

⁴<https://archive.stsci.edu/index.html>

¹The true obliquity (as opposed to the sky-projected obliquity), ψ , is the angle between the binary orbital axis and the stellar rotation axis.




Figure 1. The left-hand panel shows the light curve phase-folded on the orbital period of 152.8 d determined from the BLS search. To aid viewing the eclipse we have limited the plot in phase to ± 0.2 . A smoothed version of the phase-folded data is given by the red dashed line and near the eclipse additional variability caused by tidal interactions can be seen between a phase of ± 0.05 . The right-hand panel shows the detrended light curve of KOI-3890, showing clear eclipses, where the different colours denote different quarters of data as indicated. The vertical black dashed lines indicate the positions of the eclipses, and their expected locations in the event of there being a gap. The second eclipse that should have been observed falls in the gap between Q3 (orange) and Q4 (pink). The eclipse at the start of Q9 (orange) is only partially observed.

2.1 Eclipses

A box least-square (BLS) period search algorithm (Kovács, Zucker & Mazeh 2002) was used to detect the eclipses. The detrended light curve is shown in Fig. 1(b), whilst the light curve folded on the orbital period obtained from the BLS (152.8 d) is shown in Fig. 1(a). In Fig. 1(b), 2 of the 10 potentially visible eclipses during the *Kepler* mission are shown to be missing or only partially observed due to falling in gaps between quarters. If the light curve is folded on the period found by the BLS algorithm, it is clear that there is additional out of eclipse variability near the phase of eclipse. The variability is shown in Fig. 1(a) and is consistent with tidal interactions, as discussed in more detail in Section 5. No secondary eclipses are detected in the data due to the geometry of the orbit (as explained in Section 7), as the impact parameter of the secondary eclipse is much greater than one.

2.2 Spectroscopy

In addition to the space-based *Kepler* observations, we acquired spectroscopic data using the Tillinghast Reflector Echelle Spectrograph (TRES) ($R = 44\,000$, $\lambda = 3900 - 9100\text{ \AA}$; Szentgyorgyi & Furész 2007) on the 1.5-m Tillinghast telescope at the Fred L. Whipple Observatory to measure radial velocities for the system. The TRES spectra were reduced and analysed using the techniques outlined in Buchhave et al. (2010). Multi-order velocities were derived by cross-correlating each spectrum, order by order, against the strongest observed spectrum (which had a signal to noise per resolution element of 29). Estimates of T_{eff} , $\log g$, and $[\text{m}/\text{H}]$ were obtained using the Spectral Parameter Classification (SPC) technique described in Buchhave et al. (2012), whereby an observed spectrum is cross-correlated against a grid of synthetic spectra based on Kurucz (1992) model atmospheres. The asteroseismic value of $\log g$ was used as a prior (such that the $\log g$ was fixed to the asteroseismic value), and $[\text{m}/\text{H}]$ (the relative metal abundance) was assumed to be equivalent to the metallicity $[\text{Fe}/\text{H}]$; the values are given Table 1. However, a reliable estimate of $v \sin i$ was not able to be made for this system. In total, 10 radial velocity measurements

Table 1. Global asteroseismic and spectroscopic parameters for the red giant KOI-3890A.

KOI-3890A	
ν_{max}	$104.3 \pm 0.3\text{ }\mu\text{Hz}$
$\Delta\nu$	$9.57 \pm 0.21\text{ }\mu\text{Hz}$
$\Delta\Pi$	$77.6 \pm 0.6\text{ s}$
T_{eff}	$4726 \pm 52\text{ K}$
$[\text{Fe}/\text{H}]$	$-0.13 \pm 0.1\text{ dex}$
$\log g_{\text{astero}}$	$2.92 \pm 0.03\text{ dex}$
$\delta\nu_{\text{rot,core}}$	$0.533 \pm 0.003\text{ }\mu\text{Hz}$
i	$87.3^{+2.7}_{-1.1}\text{ degrees}$

Table 2. Radial velocity data taken with TRES used in this work.

Time (BJD)	Velocity (ms^{-1})	Uncertainty (ms^{-1})
2457171.774856	-1597.01	16.79
2457210.763886	0.00	41.42
2457237.798983	4006.68	22.57
2457289.755597	$10\,216.62$	46.41
2457296.785214	$11\,074.15$	47.32
2457304.778182	$11\,853.78$	41.42
2457319.633079	7545.42	32.03
2457349.598870	-2987.42	30.57
2457640.650105	-7313.28	41.35
2457932.777215	4686.70	53.45

were taken over 192 d, derived from multi-order fitting to spectral templates. The data taken are given in the Table 2.

In addition to the TRES observations, Lillo-Box et al. (2015) collected 22 radial velocities using the CAFE spectrograph (Aceituno et al. 2013), a high-resolution spectrograph ($R = 59\,000 - 67\,000$) in the optical range ($4000 - 9000\text{ \AA}$) situated on the 2.2m telescope at Calar Alto Observatory. Although the phase coverage of the CAFE observations is rather limited they were also incorporated during the fitting of a model to the data since they occupy a region in phase that is not well covered by the TRES observations. The addition of

the CAFE data trebled the available radial velocity measurements, giving a total of 32 RV points (22 from CAFE and 10 from TRES).

3 ASTEROSEISMIC ANALYSIS

As a red giant star, KOI-3890A exhibits solar-like oscillations, driven by the turbulent convection in the near-surface layers. We can use these oscillations to constrain the internal and global properties of the star. In this work we used the so-called global asteroseismic parameters: ν_{\max} , the frequency of maximum oscillation power; $\Delta\nu$, the average large frequency spacing between modes of the same angular degree ℓ and subsequent radial order n ; and $\Delta\Pi$, the period spacing. These parameters were extracted from the frequency power spectrum using the method described below (see also Kallinger et al. 2014; Kiefer et al. 2015; Lund et al. 2016 for additional details). For the extraction of the asteroseismic parameters we used the frequency power spectrum of the detrended light curve (see Section 2) constructed using the Lomb–Scargle periodogram (Lomb 1976; Scargle 1982) as provided by the PYTHON package GATPSY (Vanderplas 2015; VanderPlas & Ivezić 2015). Due to there only being a few eclipses in the light curve, they were left in the time series. This did not affect the asteroseismic analysis due to the fact that the signal of the eclipses in the power spectrum is at a lower frequency ($<10 \mu\text{Hz}$) than the asteroseismic signal ($\sim 100 \mu\text{Hz}$).

3.1 Global asteroseismic parameters

The first asteroseismic parameter we extract is ν_{\max} , which is determined by fitting a model to the background of the power spectrum. We adopted model H of Kallinger et al. (2014), where the granulation background is modelled as two zero-frequency-centred ‘super-Lorentzians’, where the Lorentzian is raised to an exponent that is a free parameter rather than the standard exponent of 2 or 4 (see Kallinger et al. 2014), along with one additional ‘super-Lorentzian’ at very low frequency to account for any systematic effects. Also included in the background fit is a Gaussian component to account for the power excess due to the stellar oscillations centred on the frequency of maximum power, ν_{\max} . The fit was performed using Markov chain Monte Carlo (MCMC) and the PYTHON package EMCEE (Foreman-Mackey et al. 2013); the final fit parameters were taken as the median of the posterior distributions and the uncertainties given by the 68.3 per cent highest posterior density. The final fit to the background is shown in Fig 2. The inset in the figure shows the autocorrelation function (ACF) of the power spectrum, used to identify the average large frequency separation $\Delta\nu$, following Kiefer et al. (2015). A Lorentzian was fitted to the ACF to determine $\Delta\nu$ (the main peak) and a conservative uncertainty was adopted as the half-width at half-maximum of the fitted Lorentzian to account for any additional deviations from the asymptotic pattern.

In addition to ν_{\max} and $\Delta\nu$, the period spacing ($\Delta\Pi$) of the $\ell = 1$ mixed modes was also used to recover the stellar parameters. The mixed mode frequencies were extracted by extending the method described in Davies & Miglio (2016), where we fitted the $\ell = 1$ mixed modes in the three radial orders around ν_{\max} at the same time (rather than using the properties of the mode components we had already extracted). The priors on the parameters were all taken to be uniform with the exception of the inclination angle, which was taken to be isotropic ($p(i) \propto \sin i$). The mode identification of the star given this model is shown in Fig. 3. The period spacing was then extracted from these frequencies using the method described in Hekker et al. (2018). Vrad, Mosser & Samadi (2016) derived




Figure 2. The power spectrum of KOI-3890 is shown in black alongside the fit to the background (excluding the Gaussian component describing the power excess) in red. ν_{\max} is shown by the blue dashed line. In the inset we present the ACF showing a clear peak around $\Delta\nu$ with the red shaded region showing 1σ errorbars.

a period spacing of $\Delta\Pi_1 = 75.9 \pm 0.6$ s for this star as part of their ensemble work by taking the power spectrum of the stretched power spectrum. We obtain a slightly higher period spacing than Vrad et al. (2016), though still consistent, and both show that the star is ascending the red giant branch. To account for the difference in the methods, we adopt the conservative uncertainty from Vrad et al. (2016) on our period spacing value. The asteroseismic and spectroscopic parameters are summarized in Table 1.

3.2 Rotational splittings and stellar inclination angle

In addition to the global asteroseismic parameters, due to the high signal-to-noise ratio of the oscillations the rotational splittings and inclination angle can also be extracted for this star. The derivation of the stellar inclination angle (the angle between the rotation axis of the star and our line of sight) using asteroseismology adopts the formalism derived in Gizon & Solanki (2003). Since stars rotate, modes of the same n and l and different m are not degenerate with one another. In the case of non-radial ($l > 0$) modes the relative amplitude of each component provides information about the inclination angle of the star, whilst the frequency difference provides the rotational splitting. For red giants, the inclination angle and rotational splitting can be measured using $l = 1$ mixed modes. The inclination angle posterior probability distribution was extracted from the fit to the oscillation frequencies detailed in Section 3.1, giving a value of $i(^{\circ}) = 87.3^{+2.7}_{-1.1}$. An estimate of the core rotation can be attained from the rotational splitting of the modes (following the formulation of Mosser et al. 2018), as given in Table 1. This is broadly consistent with the value obtained by Gehan et al. (2018) of $\delta\nu_{\text{rot,core}} = 0.520 \pm 0.002 \mu\text{Hz}$, allowing for differences in formulation and method.

The frequency spacing between consecutive mixed $\ell = 1$ modes in the power spectrum of the red giant star is comparable to the rotational splitting, which complicates the mode identification. In order to check that we have the correct mode identification, we investigate the rotational splitting inferred from each radial order, which should be consistent throughout. If overlapping modes are not assumed, then the rotational splittings decrease significantly as a function of radial order, which is not physical. Therefore, the




Figure 3. A section of the KOI-3890 power spectrum showing the three radial orders used in the determination of the inclination angle. Each mixed mode is denoted by a different colour per radial order. No central $m = 0$ components are seen indicating that the inclination angle is close to 90 degrees.

interpretation of overlapping modes is needed and the first guesses in the peak-bagging have been updated accordingly.

Given the eclipsing nature of the system, it is also possible to determine the obliquity (ψ) of the system. The obliquity of the system is related to the inclination angle through the following equation (Fabrycky & Winn 2009; Morton & Winn 2014)

$$\sin \psi \cos \phi = \sin i \cos \lambda \cos i_p - \cos i \sin i_p, \quad (1)$$

where λ is the sky-projected obliquity that can be determined from the Rossiter–MacLaughlin effect and ϕ is the azimuthal angle of the system. Since the system is eclipsing, $\sin i_p \approx 1$, and so $\cos i_p \approx 0$. Following Morton & Winn (2014), equation (1) can be reduced to

$$\sin \psi \cos \phi \approx \cos i. \quad (2)$$

Since we cannot distinguish between the angles i and $\pi - i$ when inferring the inclination angle, the negative sign in equation (1) can be ignored.

The azimuthal angle varies between $-\pi$ and π , where π is defined as being along the line of sight (Fabrycky & Winn 2009). The obliquity of the system can be estimated from a Monte Carlo approach using the inclination angle posterior probability density function (PDF) extracted from the fitting. We also assume that the azimuthal angle is distributed uniformly between $-\pi$ and π since from asteroseismology we cannot make any inferences about it. The obliquity of the system was found to be $\psi(^{\circ}) = 4.2^{+2.1}_{-4.2}$, the posterior of which is shown in Fig. 4. This is consistent with alignment, meaning that the plane of the orbit of the companion star is perpendicular to the stellar rotation axis of the primary (e.g. the Earth has an obliquity of $\sim 23^{\circ}$ to its orbital plane).

4 STELLAR PROPERTIES

The global asteroseismic (including the observed $\Delta\nu$ and period spacing, $\Delta\Pi_1$) and spectroscopic parameters are used as inputs to stellar models to recover the estimated stellar properties of each

Table 3. Fundamental stellar properties from PARAM; quoted uncertainties are the 68 per cent credible interval.

KOI-3890A	
M_{\star}	$1.04 \pm 0.06 M_{\odot}$
R_{\star}	$5.8 \pm 0.2 R_{\odot}$
Age	$9.1^{+2.4}_{-1.7}$ Gyr

component in the binary system. The Bayesian code PARAM⁵ (da Silva et al. 2006; Rodrigues et al. 2014, 2017) was used to recover the fundamental stellar properties, using the MESA isochrones from Rodrigues et al. 2017 (see section 2 of Rodrigues et al. 2017 for full details regarding the input physics used in the creation of the models). In PARAM, $\Delta\nu$ was calculated from theoretical radial mode frequencies using a weighted linear fit, where the weights are inversely proportional to the distance between the given frequency and ν_{\max} . This is to ensure that the computed average $\Delta\nu$ is as close as possible to the observational value (see Rodrigues et al. 2017 for more information).

Using the above physics, we obtain the fundamental stellar properties of the red giant, given in Table 3. These will enable the characterization of the companion star from the orbital parameters derived using the radial velocity and photometric observations.

5 LIGHT-CURVE AND RADIAL VELOCITY MODELLING

The transit and radial velocity data for KOI-3890 are modelled simultaneously, using MCMC⁶ to sample the parameter space. Only phases in the folded light curve of $-0.2 < \phi < 0.2$ were retained

⁵<http://stev.oapd.inaf.it/cgi-bin/param>

⁶Using PYTHON package EMCEE (Foreman-Mackey et al. 2013)




Figure 4. Kernel density estimates of the inclination angle posterior probability distribution (left-hand panel) and obliquity (right-hand panel). The median value is shown by the solid red line, whilst the 68.3 per cent highest posterior density credible interval is encompassed by the shaded region and dashed lines.

when fitting the full unfolded transit photometry as the features caused by the tidal interactions occur near the eclipse. The eclipses with incomplete phase coverage and surrounding data were also removed from the light curve.

The radial velocity data were modelled as

$$V(t) = \gamma + K [\cos(f(t) + \omega) + e \cos \omega], \quad (3)$$

where $f(t)$ is the true anomaly, e the orbital eccentricity, ω the argument of periastron, K the radial velocity semi-amplitude, and γ the zero-point offset of the Doppler velocities. The initial guess for the period was taken from the BLS detection routine.

The PYTHON package BATMAN (Kreidberg 2015) was used to model the eclipse, based upon the transit model of Mandel & Agol (2002). We assume that the secondary is not a self-luminous body and we discuss this in more detail in Section 7. Limb-darkening parameters were taken from Sing (2010), and were fixed in the transit model, using a quadratic formulation for the limb-darkening law. Additional parameters $\sigma_{RV,CAFE}$ and $\sigma_{RV,TRES}$ are included to account for any additional variance not accounted for in the original observational uncertainties for the respective radial velocity data sets. Since the *Kepler* observations are integrated over 30 min, the transit model is supersampled by a factor 5. If this is not done, the transit duration can be underestimated (Kreidberg 2015).

The initial fit to the TRES radial velocity data indicated the orbit was highly eccentric, with the eclipses being observed near periastron. For the final fitting, the TRES data were combined with the CAFE radial velocities from Lillo-Box et al. (2015), with an additional parameter included in the fit to account for zero-point offset between the data sets.

5.1 Tidal distortion of primary

The phase-folded light curve around the time of mid-transit is shown in Fig. 1(a), where the additional flux variation just outside the transit can be seen. As discussed previously, the flux contribution from the secondary component is negligible. It can therefore be assumed that the flux variation is originating from the primary (giant) star.

This additional flux variation appears to be indicative of a heartbeat star, a type of tidally induced variation (Welsh et al. 2011; Thompson et al. 2012; Beck et al. 2014; Shporer et al.

2016; Penoyre & Stone 2019). For circular orbits ($e = 0$), this tidal effect is constant, raising a bulge on the primary star. In the case of this highly eccentric system it produces ellipsoidal variation. For eccentric orbits, the term ‘heartbeat’ reflects the passing visual similarity of the tidally induced variation to an echocardiogram.

The heartbeat is induced when the tidal distortions are largest near periastron. Whilst this system has a fairly long period, its eccentricity and the evolved nature of the primary star means that, at periastron passage, the separation between the stars is only a few times the primary radius, and so the secondary is able to raise a tidal bulge on the primary.

The heartbeat flux modulation was modelled following the prescription given in Kumar et al. (1995), adjusting for the fact that we are modelling it as a function of the true anomaly,

$$\delta F = \frac{S (1 - 3 \sin^2 i_p \cos^2 [f(t) - \omega])}{[R(t)/a]^3}. \quad (4)$$

In equation (4), S is the amplitude of the flux variation, i_p the system inclination angle (known to be $i_p \sim 90^\circ$ since the eclipse is observed), $f(t)$ the true anomaly, ω the argument of periastron, and $R(t)/a$ the distance between the two stars as a fraction of the semi-major axis a , as a function of time. $R(t)/a$ can also be expressed as equation (1) of Winn (2010),

$$\frac{R(t)}{a} = \frac{1 - e^2}{1 + e \cos f(t)}. \quad (5)$$

The addition of the flux modulation into the light-curve modelling through the inclusion of equations (4) and (5) introduces only one new parameter, the amplitude of the heartbeat modulation S . All other parameters are already included within the transit and radial velocity models.

The formulation for the heartbeat signal can produce a wide variety of possible light-curve modulations, due to the possible orientations of the system in i_p and ω (see fig. 8 of Thompson et al. 2012). In the case of KOI-3890, the presence of eclipses indicates that the system is close to edge-on ($i_p \sim 90^\circ$), and so fitting the heartbeat modulation simultaneously with the transit model offers a tighter constraint on i_p .

An additional prior can be applied to the stellar density due to the fact that the red giant primary shows solar-like oscillations. The average large frequency separation $\Delta \nu$ scales to good approximation

Table 4. Model prior distributions for the simultaneous fit to the radial velocity and photometric data. Gaussian priors are indicated by \mathcal{N} (mean, standard deviation) and uniform priors by \mathcal{U} (lower bound, upper bound). All logarithmic priors are in base 10.

Parameter	Prior
P	$\mathcal{U}(151, 153)$ (days)
$\log(R_2/R_*)$	$\mathcal{U}(-3, -1)$
$\log(a/R_*)$	$\mathcal{U}(0, 2)$
T_0	$\mathcal{U}(55025, 55035)$ (BJD)
b	$\mathcal{U}(0, 1)$
$\cos \omega$	$\mathcal{U}(-1, 1)$
$\sin \omega$	$\mathcal{U}(-1, 1)$
γ_{TRES}	$\mathcal{U}(-100, 100)$ (km s $^{-1}$)
γ_{CAFE}	$\mathcal{U}(-100, 100)$ (km s $^{-1}$)
$\log(K)$	$\mathcal{U}(-1, 2)$ (km s $^{-1}$)
$\sigma_{\text{RV,TRES}}$	$\mathcal{U}(0, 10)$ (km s $^{-1}$)
$\sigma_{\text{RV,CAFE}}$	$\mathcal{U}(0, 10)$ (km s $^{-1}$)
S	$\mathcal{U}(-100, 100)$ (ppm)
ρ_*	$\mathcal{N}(7.09, 0.31)$ (kg m $^{-3}$)

with the square-root of mean stellar density (Tassoul 1980; Ulrich 1986), i.e.

$$\frac{\Delta v}{\Delta v_{\odot}} \simeq \left(\frac{M}{M_{\odot}}\right)^{0.5} \left(\frac{R}{R_{\odot}}\right)^{-1.5} \simeq \sqrt{\frac{\rho}{\rho_{\odot}}}. \quad (6)$$

The mean stellar density can also be estimated from the light curve using equation (30) of Winn (2010),

$$\rho_* \approx \frac{3\pi}{GP^2} \left(\frac{a}{R_*}\right), \quad (7)$$

where a/R_* is taken from the transit fit.

During the fitting, the stellar density at each iteration was constructed using equation (7), and the asteroseismic density (determined through PARAM) used as a Gaussian prior. The priors used during the simultaneous fit to the radial velocity and photometric data are given in Table 4.

6 RESULTS

The results of the combined transit, tidal, and radial velocity MCMC fit are listed in Table 5. Fig. 5 shows the final fit of the model to the data. As the upper panel of Fig 5 shows, the tidal distortion of the primary would artificially boost the eclipse depth if not properly accounted for. This can be an important factor when determining the radius of the secondary since it will be overestimated if the tidal distortion is not included. In the fully convective regime – $M_* \lesssim 0.33 M_{\odot}$ (Mann et al. 2015) – it is expected that the ratio of the mass and radius of an M-dwarf is approximately unity (with respect to the Sun; e.g. Demory et al. 2009). Therefore, not accounting for tidal effects would indicate a larger radius than otherwise expected and could lead to the incorrect conclusions.


In order to calculate the secondary mass, we make use of the following expansion for K (e.g. Winn 2010)

$$K = \frac{1}{\sqrt{1-e^2}} \left(\frac{P}{2\pi G}\right)^{-1/3} \frac{M_2 \sin i_p}{(M_1 + M_2)^{2/3}}, \quad (8)$$


where M_1 is the mass of the red giant primary and M_2 is the mass of the M-dwarf companion. The secondary mass was calculated by numerically solving equation (8) with a Monte Carlo method that draws samples from the posterior distributions of each parameter. The derived mass and radius of the secondary are given in Table 6.

Table 5. Model parameter median values from the simultaneous fit to the radial velocity and photometric data, and associated uncertainties taken as the 68.3 per cent credible interval.

Parameter	Median value
P	152.826 ± 0.0002 (days)
R_2/R_*	0.0444 ± 0.0002
a/R_*	20.44 ± 0.28
T_0	55030.411 ± 0.001 (BJD)
i_p	85.3 ± 0.2 (deg)
e	0.645 ± 0.001
ω	108.7 ± 0.02 (deg)
γ_{TRES}	3.96 ± 0.20 (km s $^{-1}$)
γ_{CAFE}	-30.6 ± 0.13 (km s $^{-1}$)
K	10.1 ± 0.3 (km s $^{-1}$)
$\sigma_{\text{RV,TRES}}$	0.53 ± 0.16 (km s $^{-1}$)
$\sigma_{\text{RV,CAFE}}$	0.14 ± 0.03 (km s $^{-1}$)
S	-12.0 ± 0.1 (ppm)



(a)



(b)

Figure 5. Phase-folded final fit centred around time of mid-transit including residual plots. In the upper panel, the light curve is in black, with the final model shown dashed in red near the eclipse. The two components of the light-curve model are also shown, the transit in blue and the tidal model in green. In the lower panel the TRES radial velocity data (black) and CAFE data (orange) are shown phase folded and also centred around mid-transit, with the final model shown as dashed blue. The orange and blue shaded regions correspond to the additional variance terms (see Section 5) for the CAFE and TRES data, respectively.

Table 6. Derived mass and radius for the secondary from the fit and the asteroseismic mass and radius of the primary. Both the mass and radius of the secondary are consistent with an M-dwarf.

KOI-3890B	
M_2	$0.23 \pm 0.01 M_\odot$
R_2	$0.256 \pm 0.007 R_\odot$

7 DISCUSSION AND CONCLUSIONS

In this work we have presented analysis of the red giant/M-dwarf eclipsing binary system KOI-3890. We have determined the stellar parameters of both the primary and the secondary companion along with the eclipse parameters by including tidal distortions into the fitting process.

We have made an assumption during the analysis that it is valid to fit the eclipses as transits since we do not have spectroscopic information regarding the properties of the M-dwarf companion – to treat the M-dwarf as though it is not a self-luminous body. This assumption can be justified by considering the relative flux contribution of the two components in the system. Using the derived radius of the secondary and assuming $T_{\text{eff},2} \approx 3000$ K (see Kaltenegger & Traub 2009) we obtain a luminosity of $L_2 \approx 0.005 L_\odot$ compared to $L_1 \approx 15 L_\odot$, a factor of almost 3000 in luminosity.

There is some tension present in the combined fitting of the photometry and radial velocity data as can be seen by the large unaccounted variance parameter needed for the TRES data and by the systematic trends induced in the TRES radial velocity data (see Fig. 5b). This is due to a difference in eccentricity obtained from the photometry and radial velocity. A fit to only the radial velocity data yields $e = 0.608 \pm 0.001$ as opposed to the larger value favoured by the photometry (as seen in the combined fit). This is because the eccentricity can also be constrained in the photometry through the heartbeat signal. We have much more data in the photometric data set than in the radial velocity data set and so as a result the photometry dominates the fit, leading to the larger eccentricity. Welsh et al. (2011) also noticed this type of discrepancy between the fit to the radial velocity data alone and a combined fit to both the photometry and radial velocities.

We put forward two possible explanations for the discrepancy in the eccentricities obtained from the radial velocity data and the photometry.

(i) The observed radial velocities may not correspond to the centre of mass of the tidally distorted star. The observed data correspond to the value integrated across the distorted surface weighted by the intensity for each surface element, which could distort the apparent eccentricity inferred from the velocities.

(ii) The detrending of the data may affect the amplitude and shape of the heartbeat signal in subtle ways that cannot easily be identified, yet still lead to a different eccentricity inferred overall.

Since the heartbeat signal offers a great ability to constrain the eccentricity, any additional physical features not accounted for could potentially bias the eccentricity upwards because the amplitude of the heartbeat scales with the eccentricity. This does not change the interpretation of the system, but it means that we can only reliably say that $0.6 \leq e \leq 0.65$ (this distribution is used in all simulations requiring the eccentricity), where the lower limit is set by the radial velocity data and the upper limit by the photometry.

The mass of the companion has been computed using the eccentricity from the combined fit ($M_2 = 0.23 \pm 0.01 M_\odot$), the radial velocity data only ($M_2 = 0.23 \pm 0.01 M_\odot$) and the eccentricity range we can infer $0.6 \leq e \leq 0.65$ ($M_2 = 0.23 \pm 0.01 M_\odot$). In all of these cases the mass of the M-dwarf companion is consistent and the uncertainty in the eccentricity measurement does not affect the mass measurement. This is due to the dominant uncertainty coming from the radial velocity semi-amplitude K , which is consistent between the combined and radial-velocity-only fit. Hence, the eccentricity uncertainty does not affect the interpretation of the system and so the mass of the companion given is for the eccentricity range $0.6 \leq e \leq 0.65$. However, these systematics could also propagate through to the eclipse depth, thereby affecting the radius of the secondary.

The presence of tidally-induced distortions of the primary star is clearly seen in the light curve of KOI-3890, and these ‘heartbeat’ events have been included in the eclipse fitting. We do not observe any tidally-induced oscillations in our data, but this has been observed in other systems (e.g. Guo, Gies & Fuller 2017; Hambleton et al. 2018).

As discussed earlier in Section 2, the lack of secondary eclipses (the secondary passing behind the primary from the observer’s perspective) is due to the inclination and eccentric nature of the system. The impact parameter b for transiting or occulting objects is given in Winn (2010; equations 7,8),

$$b = \frac{a \cos i_p}{R_*} \left(\frac{(1 - e^2)}{1 \pm e \sin \omega} \right), \quad (9)$$

where eclipses are (+) and occultations (−) in the denominator. For KOI-3890, the eclipse impact parameter $b_{\text{tran}} = 0.64 \pm 0.04$; however, for the secondary eclipse $b_{\text{sec}} = 2.5 \pm 0.1$. As a result, the M-dwarf does not pass behind the primary during its orbit, and so no secondary is observed, since $b \leq 1$ is required for an eclipse or secondary eclipse to occur.

It has been shown that some M-dwarfs in binary systems (where the parameters of both components can be precisely determined) show evidence of radius inflation compared to stellar models (e.g. López-Morales & Ribas 2005; Birkby et al. 2012; and references therein). Magnetic activity inhibiting convection and causing the star to ‘puff up’ is a possible explanation to describe such behaviour (e.g. Chabrier, Gallardo & Baraffe 2007; MacDonald & Mullan 2013). Feiden & Chaboyer (2012) favour a so-called hybrid interpretation whereby they suggested that, alongside magnetic activity, radius inflation could also be a result of unseen systematics in the determination of the radius. This could come about because almost all measurements of M-dwarfs come from close binaries where the components are tidally locked, active, and rapidly rotating. However, Kesseli et al. (2018) show that neither rotation nor binarity is responsible for the observed inflated radii.

Since the majority of cases where accurate M-dwarf masses and radii are available generally come from the aforementioned eclipsing systems, there are only a small number since they must be close by to be analysed with current instrumentation. Parsons et al. (2018) showed that another useful regime for inferring M-dwarf properties is M-dwarf/white dwarf eclipsing systems. They analysed 23 such systems, finding that around 75 per cent of their sample were inflated compared to theoretical models.

KOI-3890 provides a means to look at M-dwarf properties in a very different regime to the generally close-in, short-period M-dwarf/M-dwarf eclipsing systems (e.g. Demory et al. 2009). We do, however, note that this is a special case since it is in a position that is highly influenced by the red giant primary (due to

the close proximity at periastron). The general idea of using the red giant/M-dwarf regime is still valid, since the application of asteroseismology to such eclipsing binary systems can help provide additional constraints to M-dwarf stellar models, especially in the fully convective regime where the models typically struggle to reproduce the observations. The effect of systematics is something we have already touched on regarding the inclusion of the tidal distortion in the modelling of the eclipse. We can assess whether the M-dwarf companion in our system agrees with stellar models by comparing the derived mass and radius with the models from Baraffe et al. (2015) at a similar age. The closest stellar models in age to our system are those at 8 Gyr and 10 Gyr; however, the mass and radius of the M-dwarf do not change significantly over this period and so we compare to the model at 8 Gyr. Whilst the radius for KOI-3890B is larger than the predicted value from the models, $R_*/R_{\text{model}} = 1.06 \pm 0.05$, it is consistent with the model at just over the 1σ level. The models were, however, computed for solar metallicity and since KOI-3890 is slightly metal-poor when compared to the Sun (if we assume the M-dwarf has the same metallicity as the red giant primary) this may have an additional effect.

Alongside providing good constraints on the mass and radius of the M-dwarf companion through asteroseismology, it is also possible to place constraints on the age of the system, of $9.1^{+2.4}_{-1.7}$ Gyr. This is difficult for the case of the M-dwarf/M-dwarf eclipsing systems as they are reliant on the low-mass stellar models which are known to suffer from inconsistencies in the fully convective regime. In addition, the change in luminosity and effective temperature as a function of age is very small, making it more difficult to constrain M-dwarf ages from stellar models, whereas for KOI-3890 we can take advantage of the advances made in the modelling of red giant stars using asteroseismic constraints to provide an age for the system. Even though the system is a single-lined spectroscopic binary, the asteroseismic constraints can help make up for the loss of information by yielding indirect estimates of the companion properties.

As an eccentric binary around an expanding red giant star, it is interesting to explore the potential future evolution of the system. As the primary continues to ascend the red giant branch, the models used in Section 4 predict that a $\sim 1 M_{\odot}$ star reaches a radius of $\sim 160 R_{\odot}$ (0.75 au) at the tip of the red giant branch. The current separation of the two stars at periastron is $R_{\text{peri}} = 7.25 \pm 0.10 R_*$, or 0.19 ± 0.01 au, and as such, the two stars will meet during the ascent of the primary star up the giant branch.

What configuration the orbit of the secondary will end up in is determined by the rate of tidal circularization for the system. If the rate is high, then the orbital energy (and momentum) of the M-dwarf will be dissipated in the deep convective zone of the primary, as will the orbital eccentricity. To estimate the tidal circularization time-scale for a star with a convective envelope we used equation (16) of Claret, Gimenez & Cunha (1995), i.e.

$$\tau_{\text{circ}} = (1.99 \times 10^3 \text{ yr}) M^3 \frac{(1+q)^{5/3}}{q} L^{-1/3} \lambda_2^{-1} \frac{P^{16/3}(\text{d})}{R^{22/3}}. \quad (10)$$

In the above equation M , R , and L are the total stellar mass, radius, and luminosity in solar units, and q is the mass ratio of the components (M_*/M_2). λ_2 is known as the tidal constant and is related to the internal structure of the star. We take a representative value from Claret et al. (1995) of $\lambda_2 = 0.006$ (see their fig 3).

Assuming the orbital period, P , has remained constant throughout the history of the system, τ_{circ} during the main-sequence lifetime would have been far greater than the expected main-sequence

lifetime of the red giant ($\tau_{\text{circ}} \approx 2 \times 10^{17}$ yr). The main-sequence luminosity and radius have been estimated using the relations $L \propto M^{3.9}$ and $R \propto M^{0.8}$, taking M from Table 1, neglecting any mass-loss throughout the star's lifetime thus far.

As the primary ascends the red giant branch, τ_{circ} will dramatically decrease, due to the strong dependence on R (and to a lesser extent, the dependence on L). At the present epoch $\tau_{\text{circ}} \sim 9 \times 10^{11}$ Gyr. Only when the system is significantly more evolved does the radius dependence overcome the period dependence of equation (10). For instance, assuming P does not change before $R_* = R_{\text{peri}}$ then $\tau_{\text{circ}} \sim 2 \times 10^5$ yr when the stars come into contact. As the primary continues to evolve the M-dwarf may become embedded in the expanding envelope, leading to mass transfer between the stars in a common envelope phase. Additionally the strong drag forces on the secondary in such a configuration may lead to the ejection of the common envelope, and a significant decrease in the orbital period. Since the M-dwarf will encounter the expanding envelope before the primary has reached the tip of the red giant branch, the primary will not have gone through the helium flash. In the event that the common envelope is ejected, the helium core may be exposed as an sdB star, and the binary left in a close orbit, thus providing a mechanism for the formation of sdB stars. If the drag on the M-dwarf is sufficient before the ejection of the envelope, the red giant core and the M-dwarf could collide or merge inside the common envelope.

ACKNOWLEDGEMENTS

We would like to thank the referee for their helpful comments and suggestions. The authors thank Benoît Mosser and Mathieu Vrad for providing data regarding the period spacing of the star, as well as Nathalie Themeßl for many helpful discussions. An earlier version of this work appeared in J.S.K.'s PhD thesis (<http://etheses.bham.ac.uk/id/eprint/7658>). The research leading to the presented results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007–2013) / ERC grant agreement no. 338251 (StellarAges). This research has made use of the Exoplanet Follow-up Observation Program website, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. The authors acknowledge the support of the UK Science and Technology Facilities Council (STFC). Funding for the Stellar Astrophysics Centre is provided by the Danish National Research Foundation (Grant DNR106). This research has made use of the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. T. L. Campante acknowledges support from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 792848. A.M. acknowledges support from the ERC Consolidator Grant funding scheme (project ASTEROCHRONOMETRY, G.A. n. 772293). This paper includes data collected by the *Kepler* mission. Funding for the *Kepler* mission is provided by the NASA Science Mission directorate. This research has made use of the KASOC database, operated from the Stellar Astrophysics Centre (SAC) at Aarhus University, Denmark. Funding for the Stellar Astrophysics Centre (SAC) is provided by The Danish National Research Foundation. This paper uses data products produced by the OIR Telescope Data Center, supported by the Smithsonian Astrophysical Observatory.

REFERENCES

- Aceituno J. et al., 2013, *A&A*, 552, A31
- Akeson R. L. et al., 2013, *PASP*, 125, 989
- Albrecht S., Reffert S., Snellen I. A. G., Winn J. N., 2009, *Nature*, 461, 373
- Albrecht S., Reffert S., Snellen I., Quirrenbach A., Mitchell D. S., 2007, *A&A*, 474, 565
- Albrecht S., Setiawan J., Torres G., Fabrycky D. C., Winn J. N., 2013, *ApJ*, 767, 32
- Albrecht S., Winn J. N., Carter J. A., Snellen I. A. G., de Mooij E. J. W., 2011, *ApJ*, 726, 68
- Albrecht S. et al., 2014, *ApJ*, 785, 83
- Arentoft T. et al., 2008, *ApJ*, 687, 1180
- Baglin A., Auvergne M., Barge P. et al., 2006, in Fridlund M., Baglin A., Lochard J., Conroy L., eds, *The CoRoT Mission Pre-Launch Status - Stellar Seismology and Planet Finding*, Vol. 1306, Scientific Objectives for a Minisat: CoRoT. ESA Special Publication, p. 33
- Baraffe I., Homeier D., Allard F., Chabrier G., 2015, *A&A*, 577, A42
- Bate M. R., Lodato G., Pringle J. E., 2010, *MNRAS*, 401, 1505
- Beck P. G. et al., 2014, *A&A*, 564, A36
- Bedding T. R. et al., 2010, *ApJ*, 713, 935
- Benbakoura M., Gaulme P., McKeever J., Beck P. G., Jackiewicz J., García R. A., 2017, in Reylé C., Di Matteo P., Herpin F., Lagadec E., Lançon A., Meliani Z., Royer F., eds, *SF2A-2017: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics*. p. 89
- Birkby J. et al., 2012, *MNRAS*, 426, 1507
- Bonnell I., Arcoragi J.-P., Martel H., Bastien P., 1992, *ApJ*, 400, 579
- Borucki W. J. et al., 2010, *Science*, 327, 977
- Brogaard K. et al., 2016, *Astron. Nachr.*, 337, 793
- Brogaard K. et al., 2018, *MNRAS*, 476, 3729
- Brown T. M., Gilliland R. L., Noyes R. W., Ramsey L. W., 1991, *ApJ*, 368, 599
- Buchhave L. A. et al., 2010, *ApJ*, 720, 1118
- Buchhave L. A. et al., 2012, *Nature*, 486, 375
- Burrows A., Hubbard W. B., Lunine J. I., Liebert J., 2001, *Rev. Mod. Phys.*, 73, 719
- Campante T. L. et al., 2016, *ApJ*, 819, 85
- Chabrier G., Gallardo J., Baraffe I., 2007, *A&A*, 472, L17
- Chaplin W. J. et al., 2014, *ApJS*, 210, 1
- Claret A., Gimenez A., Cunha N. C. S., 1995, *A&A*, 299, 724
- da Silva L. et al., 2006, *A&A*, 458, 609
- Davies G. R., Miglio A., 2016, *Astron. Nachr.*, 337, 774
- De Ridder J. et al., 2009, *Nature*, 459, 398
- Demory B.-O. et al., 2009, *A&A*, 505, 205
- Eggenberger P. et al., 2017, *A&A*, 599, A18
- Fabrycky D. C., Winn J. N., 2009, *ApJ*, 696, 1230
- Feiden G. A., Chaboyer B., 2012, *ApJ*, 757, 42
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Frandsen S. et al., 2013, *A&A*, 556, A138
- Gaudi B. S., Winn J. N., 2007, *ApJ*, 655, 550
- Gaulme P., Jackiewicz J., Appourchaux T., Mosser B., 2014, *ApJ*, 785, 5
- Gaulme P., McKeever J., Rawls M. L., Jackiewicz J., Mosser B., Guzik J. A., 2013, *ApJ*, 767, 82
- Gaulme P. et al., 2016, *ApJ*, 832, 121
- Gehan C., Mosser B., Michel E., Samadi R., Kallinger T., 2018, *A&A*, 616, A24
- Gizon L., Solanki S. K., 2003, *ApJ*, 589, 1009
- Guo Z., Gies D. R., Fuller J., 2017, *ApJ*, 834, 59
- Hambleton K. et al., 2018, *MNRAS*, 473, 5165
- Hekker S. et al., 2009, *A&A*, 506, 465
- Hekker S. et al., 2010, *ApJ*, 713, L187
- Hekker S. et al., 2011, *MNRAS*, 414, 2594
- Hekker S., Elsworth Y., Angelou G. C., 2018, *A&A*, 610, A80
- Huber D. et al., 2013, *Science*, 342, 331
- Kallinger T. et al., 2014, *A&A*, 570, A41
- Kaltenegger L., Traub W. A., 2009, *ApJ*, 698, 519
- Kamiaka S., Benomar O., Suto Y., 2018, *MNRAS*, 479, 391
- Kesseli A. Y., Muirhead P. S., Mann A. W., Mace G., 2018, *AJ*, 155, 225
- Kiefer R., Schad A., Herzberg W., Roth M., 2015, *A&A*, 578, A56
- Kjeldsen H. et al., 2005, *ApJ*, 635, 1281
- Kovács G., Zucker S., Mazeh T., 2002, *A&A*, 391, 369
- Kreidberg L., 2015, *PASP*, 127, 1161
- Kumar P., Ao C. O., Quataert E. J., 1995, *ApJ*, 449, 294
- Kurucz R. L., 1992, in Barbuy B., Renzini A., eds, *Proc. IAU Symp. 149, The Stellar Populations of Galaxies*. Kluwer Academic Publishers, Dordrecht, p. 225
- Lillo-Box J., Barrado D., Mancini L., Henning T., Figueira P., Ciceri S., Santos N., 2015, *A&A*, 576, A88
- Lomb N. R., 1976, *Ap&SS*, 39, 447
- López-Morales M., Ribas I., 2005, *ApJ*, 631, 1120
- Lund M. N. et al., 2014, *A&A*, 570, A54
- Lund M. N. et al., 2016, *MNRAS*, 463, 2600
- MacDonald J., Mullan D. J., 2013, *ApJ*, 765, 126
- Mandel K., Agol E., 2002, *ApJ*, 580, L171
- Mann A. W., Feiden G. A., Gaidos E., Boyajian T., von Braun K., 2015, *ApJ*, 804, 64
- Mathur S., García R. A., Huber D., Regulo C., Stello D., Beck P. G., Houmani K., Salabert D., 2016, *ApJ*, 827, 50
- Mazeh T., Shaham J., 1979, *A&A*, 77, 145
- Miglio A. et al., 2013, *MNRAS*, 429, 423
- Miglio A., Chaplin W. J., Farmer R., Kolb U., Girardi L., Elsworth Y., Appourchaux T., Handberg R., 2014, *ApJ*, 784, L3
- Montalbán J., Miglio A., Noels A., Dupret M.-A., Scuflaire R., Ventura P., 2013, *ApJ*, 766, 118
- Morton T. D., Winn J. N., 2014, *ApJ*, 796, 47
- Mosser B. et al., 2012, *A&A*, 548, A10
- Mosser B., Gehan C., Belkacem K., Samadi R., Michel E., Goupil M., 2018, *A&A*, 618, A109
- Parsons S. G. et al., 2018, *MNRAS*, 481, 1083
- Penoyre Z., Stone N. C., 2019, *AJ*, 157, 60
- Pesnell W. D., 1985, *ApJ*, 292, 238
- Rawls M. L. et al., 2016, *ApJ*, 818, 108
- Rodrigues T. S. et al., 2014, *MNRAS*, 445, 2758
- Rodrigues T. S. et al., 2017, *MNRAS*, 467, 1433
- Scargle J. D., 1982, *ApJ*, 263, 835
- Shporer A. et al., 2016, *ApJ*, 829, 34
- Silva Aguirre V. et al., 2018, *MNRAS*, 475, 5487
- Sing D. K., 2010, *A&A*, 510, A21
- Smith P. H., McMillan R. S., Merline W. J., 1987, *ApJ*, 317, L79
- Soderblom D. R., 2010, *ARA&A*, 48, 581
- Szentgyorgyi A. H., Furész G., 2007, *Rev. Mex. Astron. Astrofis. Ser. Conf.*, 28, 129
- Tassoul M., 1980, *ApJS*, 43, 469
- Themeßl N. et al., 2018, *MNRAS*, 478, 4669
- Thompson S. E. et al., 2012, *ApJ*, 753, 86
- Ulrich R. K., 1986, *ApJ*, 306, L37
- Vanderplas J., 2015, *gatspy: General tools for Astronomical Time Series in Python*. <https://doi.org/10.5281/zenodo.14833>
- VanderPlas J. T., Ivezić Ž., 2015, *ApJ*, 812, 18
- Vrard M., Mosser B., Samadi R., 2016, *A&A*, 588, A87
- Welsh W. F. et al., 2011, *ApJS*, 197, 4
- Winn J. N., 2007, in Afonso C., Weldrake D., Henning T., eds, *ASP Conf. Ser. Vol. 366, Transiting Extrapolar Planets Workshop*. Astron. Soc. Pac., San Francisco, p. 170
- Winn J. N., 2010, in Seager S., ed., *Exoplanet Transits and Occultations*. Univ. Arizona Press, p. 55
- Yu J., Huber D., Bedding T. R., Stello D., Hon M., Murphy S. J., Khanna S., 2018, *ApJS*, 236

This paper has been typeset from a \LaTeX file prepared by the author.