

From admission to discharge in mental health services: A qualitative analysis of service user involvement

Wright, N.; Rowley, E.; Chopra, A.; Gregoriou, K.; Waring, J.

DOI:
[10.1111/hex.12361](https://doi.org/10.1111/hex.12361)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Wright, N, Rowley, E, Chopra, A, Gregoriou, K & Waring, J 2015, 'From admission to discharge in mental health services: A qualitative analysis of service user involvement', *Health Expectations*, vol. 19, no. 2, pp. 367-376. <https://doi.org/10.1111/hex.12361>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 17/07/2019
Wright, N. , Rowley, E. , Chopra, A. , Gregoriou, K. and Waring, J. (2016), From admission to discharge in mental health services: a qualitative analysis of service user involvement. *Health Expect*, 19: 367-376. doi:10.1111/hex.12361

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

From admission to discharge in mental health services: a qualitative analysis of service user involvement

Nicola Wright PhD MA BN RN,* Emma Rowley PhD MSc BSc,† Arun Chopra MBBS MMedSci MRCPsych,‡ Kyriakos Gregoriou MNursSci§ and Justin Waring PhD MSc BA¶

*Assistant Professor in Mental Health, School of Health Sciences, University of Nottingham, †Senior Research Fellow Collaboration for Leadership in Applied Health Research and Care East Midlands, Deputy Director of Centre for Health Innovation, Learning and Leadership, Nottingham University Business School, University of Nottingham, ‡Consultant Psychiatrist, §Band Six Community Mental Health Nurse, Nottinghamshire Healthcare NHS Trust, and ¶Professor of Organisational Sociology, Theme Lead Collaboration for Leadership in Applied Health Research and Care East Midlands, Director of Centre for Health Innovation, Learning and Leadership, Nottingham University Business School, University of Nottingham, Nottingham, UK

Correspondence

Dr Nicola Wright, PhD MA BN RN
Assistant Professor in Mental Health
School of Health Sciences
University of Nottingham
Institute of Mental Health
Triumph Road
Nottingham
NG7 2TU
UK
E-mail: nicola.wright@nottingham.ac.uk

Accepted for publication

1 March 2015

Keywords: care transitions, mental health, service user involvement

Abstract

Background User involvement and recovery are now widely used terms within the mental health policy, research and practice discourse. However, there is a question mark about the impact these ideas have in everyday practice. Of interest is the degree of involvement in key transitions of care. In particular, admission to and discharge from acute inpatient mental health wards.

Objective To explore the nature of service user involvement in the admission and discharge process into and out of acute inpatient mental health care.

Design A qualitative study using focus groups.

Setting and Participants One acute, inpatient mental health ward was the focus of the study. Seven uniprofessional focus group interviews were conducted with ward staff, community staff and service users (total number of participants = 52). Conventional, thematic qualitative techniques were used to analyse the data.

Results The data analysed and presented in this article relate to the loss of the service user voice at the key transition points into and out of acute inpatient care. Due to the lack of resources (inpatient beds and community care follow-up), the role service users could play was diminished. In their narratives, clinical staff associated the person with the process and used language which dehumanized the individual.

Conclusion Service users experience numerous care transitions into and out of hospital. As there is the potential for these encounters to have a lasting negative effect, the importance of ensuring service users have a voice in what is happening to them is crucial.

Introduction

User involvement has become a central tenant of the design, provision and evaluation of mental health services. Understanding and recognizing the importance of an individual's experience of mental distress from their own perspective has gained increasing prominence in the past 20 years.^{1,2} Closely linked to both the consumer/survivor movement and the concept of recovery, the mantra 'no decision about me without me' has become common within the policy discourse.³

At its core, the user movement is based on ideas relating to self-help, empowerment and advocacy² and it provides a challenge to the traditional notions of professional power and expertise.⁴ Shepherd *et al.*² argue that these concepts are not new in themselves and have their roots in the American civil rights movements of the 1960s and 1970s. However, Frese and Davis⁴ argue that the history of user involvement can be traced back even further to 1845 in the UK when the Alleged Lunatics Friend Society was established and to the period immediately following the civil war in the USA (when the anti-insane asylum society was set up).

Like the concept of user involvement, Shepherd *et al.*² suggest that 'recovery is an idea whose time has come' as it incorporates five current trends in mental health. Namely, social inclusion, more responsive services matched to the needs of people, conceptual changes about the nature of mental health problems, an emphasis on individual rather than collective solutions and self-management.⁵ One of the architects of the recovery movement, Anthony (p. 527)⁶ argues that:

recovery is described as a deeply personal, unique process of changing ones attitudes, values, feelings, goals, skills and/or roles. It is a way of living a satisfying, hopeful and contributing life even with the limitations caused by illness. Recovery involves the development of new meaning and purpose in one's life as one grows beyond the catastrophic effects of mental illness.

Empirical studies such as Corrigan *et al.*, Harding *et al.* and Perry *et al.*⁷⁻⁹ provide

evidence for some of the underpinning ideas of recovery and challenge the view first articulated by Kraeplin¹⁰ that psychosis has an inevitable downward and deteriorating course. However, the concept is not without its critics.¹¹ The term may be misunderstood as being synonymous with 'cure', and therefore, it is perceived to be impossible for people with on-going mental health problems to achieve recovery.¹² Professionals may also perceive maintenance and recovery to be the same thing.¹³

Despite user involvement and recovery being in the ascendancy in local and national mental health policy, it has been questioned whether these ideals have led to more collaborative ways of working and shared decision making in the practice setting.¹⁴ It is suggested that service users with mental health problems are more likely to want involvement in decision making in comparison with those with general medical conditions.¹⁵ Peer-led initiatives such as Wellness Recovery Action Plans (WRAP) have demonstrated some positive outcomes.¹⁶ However, when initiatives have focused on collaborative planning and decision making between professionals and service users, such as facilitated joint crisis plans, limited evidence of effectiveness has been found.¹⁷ Like many health-care practice innovations, it could be argued that a gap exists between what is espoused at the policy level or in the research literature (i.e. user involvement across all strata of mental health care) and what happens in everyday care situations.

Waring *et al.*¹⁸ argue that care transitions involve a multitude of health and social care professionals working within and across different organizational boundaries. Taken within this context, the movement of service users into and out of acute inpatient mental health wards is particularly complex, given the potential for high emotion and coercive practice.¹⁹ The number of people admitted to a psychiatric hospital on a compulsory basis in England and Wales per head of population increased by over 50% in the decade to 1995 and then rose by 13% from 26, 632 to 30 092 during the decade to 2010–2011.²⁰ Given that most service

users find involuntary treatment a negative experience and describe it as unjustified even 12 months later,²¹ mechanisms for service user involvement appear to be crucial in this process.

The transition out of inpatient wards (discharge) back to the community is also challenging. Whereas strategies to reduce hospital admissions have received a large amount of research attention, including innovations for more collaborative or user-focused approaches (some of which are mentioned above), the same cannot be said in relation to hospital discharge in mental health. The first 7 days after discharge from hospital have been identified as a 'critical period' of post-discharge care when people with mental health problems are at increased risk of suicide.²² Although suicide is a devastating consequence, it is also relatively rare. In contrast, service users and their carers report a range of more mundane care problems that arise from discharge planning when their views have not been taken into account. Anecdotal reports include descriptions of medication being unavailable for collection and disruptions to social security payments.

In summary, user involvement and the concept of recovery are widely referred to within the policy, research and practice discourses. However, the extent to which this has 'trickled down' to make an impact on individual practice and care experiences is unclear. In particular, the points of transfer of care from community to inpatient and back to community are areas where the involvement of service users appears crucial and yet under reported. Therefore, this study aimed to explore the nature of service user involvement at these key transition points.

Methods

Study design

The data reported in this study originate from a study that focused on knowledge sharing at the points of transition of care into and out of inpatient mental health services.²³ Using

an improvement science methodology,^{23,24} it aimed to create a joint narrative with stakeholders (including those with lived experience) of the barriers and facilitators to knowledge sharing. The study also aimed to explore possible innovations which could be implemented to aid this process. This was a single site study which focused on one acute inpatient mental health ward and the teams and practitioners who worked both within that environment and also who supported service users prior to admission and following discharge to the community. Ethical approval for the study was granted, and NHS research governance procedures were followed (for details see Acknowledgements).

Given the complex interactions between multiple stakeholders and the multifaceted nature of the phenomenon of study, qualitative focus groups were selected as the method of choice. Focus group interviews allow for the collection of richer data as the individuals involved interact with each other as well as the interviewer to test and develop their ideas, views and opinions.²⁵ It is suggested, however, that the strength of the method is also its largest weakness. The interactive nature of the focus group interview can lead to some individuals becoming dominant and the possibility of hierarchical relationships being relocated from society or the organization into the group setting.²⁶ Within mental health care, power is a crucial aspect which needs to be considered (for example, practitioners can detain service users in hospital against their will). To attempt to redress some of these dynamics, the focus groups were organized to be uniprofessional (for example, mental health nurses), team (for example, community team staff) or service user specific.

Recruitment, sampling and participants

Due to the single site design, a purposive sampling strategy was used to identify groups and individual stakeholders who had experience of the phenomenon of study. These were as follows: consultant psychiatrists (inpatient and

community), junior (inpatient based) medical staff, inpatient mental health nurses (including nurses working in the local 136 suite¹), health-care assistants, community mental health services and service users. In total, seven focus groups were conducted and 52 participants were involved in the study.

Participants were recruited by email, phone or personal approach from a member of the research team. Full information was provided both verbally and in writing to those interested in taking part. Informed consent was obtained from all individuals. Participants were made aware that they could withdraw their consent to take part in the study at any time. The nature of a focus group means that it is not possible to assure individuals of confidentiality. However, those who took part were reassured that the raw data would be anonymized as soon as practicable after the focus group had taken place.

Data collection

Focus groups were conducted in winter 2013/14 in locations and at times convenient for the participants. They lasted for approximately 60 min and were audio-recorded and transcribed verbatim. Two members of the research team attended each focus group, one acted as the group facilitator while the other made detailed field notes to supplement the audio recording.

A semi-structured topic guide (see Table 1) was used to steer the focus group interview and was informed by themes from the literature and the experiences of members of the research team from working in mental health services. A semi-structured approach allowed for a 'conversation with purpose' to occur²⁷ but also had flexibility so that participants could discuss issues which may be particularly relevant to their experiences. Immediately prior

¹The 136 suite is a unit for the short term assessment of individuals who come into contact with the police. Following assessment individuals may be admitted to an inpatient mental health ward or discharged back to the community.

Table 1 Subject areas considered in the focus group interviews

Areas considered in relation to care transitions into hospital:
What leads to an admission
Who is involved – including the role of the service user
How long does it take
Good practice examples – what is a good admission
Challenges with the process
What can go wrong
What would make the process better
How, when and by whom is knowledge and information gained, shared and stored
Areas considered in relation to care transitions out of hospital:
When does the discharge planning process start
Who is involved – including the role of the service user
How long does it take – including delays to the discharge process and its effects
Good practice examples – what is a good discharge
Challenges with the process
What can go wrong
What would make the process better
How, when and by whom is knowledge and information gained, shared and stored

to the start of the focus groups, the process was explained to participants. It was reiterated that there were no right or wrong answers and all opinions were valued even if individuals disagreed with each other. After the main group discussion, participants were also given the opportunity to stay behind and speak with the researchers individually if they wished to do so; no one took advantage of this.

Analysis

Prior to commencing data analysis, the transcripts were checked for accuracy by comparing them to the original recordings. Any additional details from the field notes were also added to the transcripts. Familiarity with the data was ensured by reading the transcripts multiple times. Conventional qualitative methods were used to identify analytical patterns (or themes) across and within the transcripts.²⁸ Analysing data thematically, although time-consuming, provides a concise and coherent account of the story which it tells.²⁹ After obtaining an understanding of the whole narrative, meaning units

were identified within the text.³⁰ These were then grouped together, without losing the original context in which the statements were made. These grouped units were consolidated into codes, and the similarities and differences between them were compared. A further consolidation process led to the development of tentative themes which explained the data. Two members of the research team analysed the data individually and then compared the themes which they had identified. A high level of consistency was found between the two researchers in terms of the themes reported and the relevance of them to the research question. These findings were then discussed with the wider research team for verification purposes. The Nvivo computer program was used to manage the analysis process.

This article presents and discusses the theme ‘the lost service user voice’. Data in relation to the other themes identified as part of the project are reported elsewhere.

Results

Participants were asked in the focus groups to discuss and reflect on their experiences of knowledge sharing. Of particular interest were the care transition points of going into and out of acute inpatient mental health services. For all the participants (clinicians and service users), these transitions were chaotic, stressful and emotionally charged. Overall, clinical staff expressed a desire to provide the best care possible in these circumstances. Service users recognized the organizational and system constraints which impacted on how care was given and received. However, despite this desire and understanding, the service user voice was lost within the narrative of care.

Explanatory, direct quotations from the focus group transcripts are used to illustrate the analysis presented. To maintain the anonymity of the participants, the quotes are labelled only as being from inpatient, community or service user. Any identifiable details within the excerpts have been removed and replaced with *X*. Comments in square brackets have been added by the

authors to aid understanding for the reader, and ellipses denote removed sections.

The lost service user voice

The narratives from all the focus groups demonstrate a fraught and competitive environment at the transition points into and out of inpatient care. Inpatient beds are scarce resources and accessing them is extremely challenging. Given this scarcity, service users had a very limited voice in the decision-making process related to their admission and discharge from hospital. They recognized that at times, they did pose a risk to either themselves or other people and needed to be in hospital against their will. In these situations, service users stated that they valued open communication:

But you see sometimes they are right aren't they because we are a danger to ourselves and we're a danger to other people... We could go out in front of traffic and get ourselves killed... but it would be nice sometimes for it just to be dead straight. If someone could say look you're a danger to yourself and you're a danger to others. And maybe we've made an assessment and this is why. (service user)

However, if service users agreed to admission or identified that a period of respite in hospital would be helpful, facilitating this was difficult. Community staff reported that ‘informal’ admissions (where the service user requests or agrees to go into hospital voluntarily rather than being compelled by law) were virtually impossible unless there was personal contact with the inpatient consultant and you were prepared to do some ‘wheeling and dealing’:

I think it's increasingly difficult to get a bed and especially if you have a patient who is agreeing to come in for an informal bed, it's nigh on impossible to get them an informal bed... We had one person in didn't we informally. I got one person in informally but that depended on a bit of wheeling and dealing with one of the other consultants. (community team)

As well as being competitive, the environment and culture at these transition points were also conflictual and chaotic. Shared ownership of the

'problem' was identified as being non-existent, and clinical knowledge was devalued. Community staff reported that they felt their experience of working with an individual was dismissed when inpatient staff reported that they were not psychotic but personality disordered or under the influence of illicit substances:

So you'd get a patient like who've I've had before whose paranoid schizophrenia, they go into the ward and get re-diagnosed with something completely different, either a personality disorder or psychotic depression... there is no continuity of care. Despite the fact that you've got one consultant quite sure of what is going on and that affects how they decide to work with that person, it changes the kind of care that they are getting. More often than not they get discharged quite quickly because they are not psychotic at all, they have a personality disorder so they have to be discharged. (community team)

This quote raises an interesting point as to whether some service user voices are more valued and valid than others. For example, within one of the inpatient transcripts, a participant refers to 'they're all bloody PD'. This was explained by one participant as being a mechanism for being able to emotionally shut off from these individuals:

It's like PD [Personality Disorder] you know you tick the box... to that so you don't have to care for them. Emotionally you don't have to think about them afterwards you can just forget about them. (community team)

Service users also identified that some people were more likely to get lost within the system. Rather than being based on diagnosis, they suggested that it was those who were quiet and posed no challenge. Service users suggested that to achieve their goals they needed to be persistent and constantly ask for information:

I think one of the problems X for you and I'll be honest is that you come over as articulate and able to function... and I know you have considerable difficulties but for them you don't cause a huge fuss... You don't kick off big time. (service user)

The culture of care at these transition points was characterized by who had the power and

control of the resources required. At admission, this was linked to the availability of inpatient beds. At discharge, it was the availability of community staff which was perceived to be the limiting factor. Within this context, service users reported that their expertise in knowing their own mental health and identifying their needs was lost. In relation to his discharge from hospital, one service user described how his perception of health and illness was secondary to the need to free up bed space:

I was pulled in for what I thought was routine psychiatric appointment with Dr X and I was told 'I want to send you home today'. Out of nowhere... so I didn't take it well. I didn't feel ready to go out... He said he was going to be honest because I deserved it. He had pressure from above to free the beds up and I said to him 'so you don't think I am well enough to go home but it's just you need a few beds' and so I was not very happy. (service user)

It was not just service users who identified this as an issue, staff were aware that increasingly they had to discharge people when they were not ready or did not have the required aftercare to support them in the community:

We're in a position now that we're having to make some very difficult decisions and discharging people into circumstances that years ago we wouldn't have dreamed of. So we are discharging people without allocated care coordinators with just crisis as a 7 day follow up... we've had a couple of incidences where we've had to discharge people to the pavement with no accommodation... And I think as X, I feel very uncomfortable but we're having to do that increasingly more. (inpatient team)

The language participants used to describe the care transitions emphasized that the service user voice was missing. The person and the processes such as referral to other agencies were often perceived to be one and the same. Therefore, the narrative was dominated by the word 'it'. 'It' was something to be 'picked up and run with', something 'which had to be dealt with' or 'handed over verbally'. Adjectives such as 'being bounced from one pathway to the next', 'being dumped back in the home situation' or 'shipping them out' described the

movement of service users throughout the mental health-care system. In this context, service users are perceived as objects to be passed from one service to the next. However, all the participants identified that it was the system that was leading to this situation occurring rather than a deliberate motivation on the part of an individual to exclude or dehumanize service users:

We've probably been quite critical, you've asked us for our opinions really which we have given you our opinions and recognise that they do work hard but they do try very hard with people and sometimes I think it is the systems rather than the people... that let the whole thing down.(community team)

One participant took this further and stated that the system had removed the compassion from mental health care:

We all come to this job to help people and it is the system that stops us from doing that, heaven help you if you fill out a referral form wrong... the system has removed compassion from care.(inpatient team)

Service users were also aware of the constraints and pressures staff were working under:

I'm not here to in anyway...defend the system or anything else but actually it is quite hard for them. If you look at the number of staff on and the number of patients actually on the ward and you think of some of those individuals and how much time they take up, I mean there are some very unwell people on that ward.(service user)

Discussion

This study aimed to explore the nature of service user involvement at the transition points of admission and discharge to an acute inpatient mental health ward. Analysis of the data collected found that despite the current rhetoric of recovery and involvement in policy and practice discourses, the service user voice was lost at these key care delivery points. This was evident in the language used to describe what happened during care transitions and also the culture of the teams and organizations involved. While it was recognized that clinicians did not come to

mental health care with the intention of working in this way, the current organizational context had removed the compassion from clinicians' work.

The narratives described in this study identify that inpatient beds are a precious and restricted resource. Tyrer³¹ identifies that there has been a dramatic reduction in the number of beds available from the maximum in 1954 (155 000) to 27 000 in 2008. As the number of beds has reduced, this has had a 'concentrating effect' whereby the threshold for admission has increased with more service users being subjected to legal restrictions and detained in hospital against their will.³² This concentrating effect not only impacts on the inpatient environment (potentially making it a more volatile setting) but also on the quality of care in the community.³³ Quirk and Lelliott³² identify that community services rely on easy access to inpatient beds for respite periods and to manage crisis situations. However, it should be noted that the availability of beds is not the only limited resource highlighted within this study. For the care transition from inpatient ward to community to be facilitated; there also needed to be an adequate supply of community staff. These findings indicate, therefore, that the more restricted the resources the increased likelihood that the service user voice will be lost.

At the point of admission to hospital, service users identified that they may pose a risk to themselves or others and may need professionals to make decisions on their behalf. However, they were also clear that when they were able to engage with decision making, they wanted their voice heard. Initiatives such as WRAP plans¹⁶ and crisis plans¹⁷ were originally conceived to allow service users to express their preferences during a period of mental wellness so that the care plan acts as their voice and expresses their views during a crisis when they are unable to do so. However, given the current context of mental health-care provision, the implementation of this may be problematic. For example, clinicians within this study noted that should an individual request admission to hospital, it was virtually impossible to arrange.

Data from the study also identified that some voices were more likely to be heard than others. Studies such as Lauber *et al.*³⁴ demonstrate that service users can experience stigma and discrimination from the negative attitudes of mental health staff. Within the literature, particular prominence is given to those with comorbid substance misuse issues and those diagnosed with personality disorder. More negative attitudes are expressed when individuals are perceived to be 'not ill' and therefore wasting precious resources or being in some way responsible for their predicament.³⁵ As well as differential experiences based on diagnosis those service users who were deemed to be quiet were more likely to get lost at care transition points. The service user narrative demonstrates that they were aware of these issues and took active steps to increase the chances that their voices were heard, for example persistently asking the staff questions and making demands of them. Just as Quirk *et al.*³⁶ found in relation to managing risk, the data from this study highlights that service users attempt to take active steps to be heard rather than be passive recipients of staff intervention.

Goffman³⁷ conceptualized asylums as 'total institutions', namely an isolated and enclosed social system with the purpose of controlling the behaviour of individuals who live within it. Although it has been questioned whether the totality of asylums ever existed,³⁸ the 'membrane' between inpatient and community care is certainly more 'permeable' today.³⁹ Data from this study suggest that service users experience numerous transitions of care from community to inpatient services (and back again) during their 'psychiatric career'. For practitioners, care transitions at the permeable membrane involve negotiation, time and the investment of emotional energy. The findings from this study identify that these transitions are 'flash points' for conflict between services. As the expectation for shorter admissions gains traction,²⁸ the likelihood is that service users may experience more transitions in their care. Developing interventions and tools which ensure that their voice is heard and are easily

implementable in the current context appears to be crucial.

Strengths and limitations of the study

This study has illuminated some interesting and pertinent issues relating to service user involvement at the interface of community and inpatient mental health care. However, no research method is perfect and the contribution to knowledge which is made needs to be considered in the context of the methodological strengths and limitations.

Basing the study on a single research site (the inpatient ward) and the clinicians and service users who either work in it or interface with it allowed for an in-depth exploration of involvement in the admission and discharge process. However, this approach may also limit the applicability and transferability of the findings to other settings. In particular, the way mental health care is organized in respect of separate inpatient and community teams, as in this case, has been widely adopted in England following the *New Ways of Working* report,⁴⁰ but it is not universal. Similarly, there are variations in the organization and provision of mental health care internationally. This does not mean that the findings are completely irrelevant; however, consideration may need to be given when applying them to other contexts.

As well as this purposive approach, a degree of self-selection of the sample also occurred. For example, it was not possible to interview every individual who interfaced with the research site. Those who came forward to take part in the focus groups may have had particularly strong views (positive or negative), and this may have skewed the data. Only those participants who were fluent in the English language could be included in the study. It may be that service users who are non-English speakers had different experiences to those described. A further limitation is the lack of involvement of carers in the study as they may have particular issues and concerns relating to care transitions into and out of inpatient settings.

The interactive nature of focus groups is both a strength and limitation of the method. As identified previously in this article, a group setting allows participants to generate and discuss ideas between themselves. However, it is possible that the responses obtained are not the same as those which would have been generated from a one-to-one interview. For example, individuals may feel inhibited from expressing their views if these are different to the majority. Focus group interviews only provide a snapshot of experiences at a given a time point. It is not clear how consistent these views are and whether they have been influenced (positively or negatively) by recent experiences.

Conclusion

Service user involvement is a central tenant of mental health practice, policy and research. Closely related to the concept of recovery, it values expertise by experience on a par with professional knowledge. However, this study has highlighted how at key transition points in care delivery the service user voice can be lost. The current context of care is dominated by restricted resources. Within this climate, innovative solutions are needed to make sure that service users are able to influence the delivery of their care at the key points of admission and discharge to hospital.

Acknowledgements

Thanks to the individuals and services who agreed to take part in the study. We would also like to thank Bridget Roe and Ted Bradshaw who assisted with facilitating the focus groups. The study was reviewed and given a favourable opinion by a University of Nottingham Research Ethics Committee and the relevant NHS organization.

Funding

The study was supported by the SPARK fund from the Nottingham University Business School. ER and JW are part funded for their

time by NIHR CLAHRC EM, a collaboration of academic and NHS organizations across the East Midlands region. The views expressed in this article are those of the authors and not necessarily those of the NIHR, the University or NHS.

References

- 1 Davidson L, Roe D. Recovery from versus recovery in serious mental illness: one strategy for lessening confusion plaguing recovery. *Journal of Mental Health*, 2007; **16**: 459–470.
- 2 Shepherd G, Boardman J, Slade M. *Making Recovery a Reality*. London: Sainsbury Centre for Mental Health, 2008.
- 3 Department of Health. *Liberating the NHS: No Decision About Me Without Me*. London: Department of Health, 2012.
- 4 Frese FJ, Davis WW. The consumer-survivor movement, recovery and consumer professionals. *Professional Psychology: Research and Practice*, 1997; **28**: 243–245.
- 5 Davidson L. Recovery, self-management and the expert patient: changing the culture of mental health from a UK perspective. *Journal of Mental Health*, 2005; **14**: 25–35.
- 6 Anthony WA. Recovery from mental illness: the guiding vision of the mental health service system in the 1990s. *Psychosocial Rehabilitation Journal*, 1993; **16**: 521–538.
- 7 Corrigan PW, Gifford D, Rashid F, Leary M, Okeke I. Recovery as a psychological construct. *Community Mental Health Journal*, 1999; **35**: 231–239.
- 8 Harding CM, Brooks GW, Ashikanga T, Strauss JS, Breir A. The Vermont longitudinal study of persons with severe mental illness 1: methodology, study sample and overall status 32 years later. *American Journal of Psychiatry*, 1987; **144**: 718–726.
- 9 Perry BM, Taylor D, Shaw SK. “You’ve got to have a positive state of mind”: an interpretive phenomenological analysis of hope and first episode psychosis. *Journal of Mental Health*, 2007; **16**: 781–793.
- 10 Kraepelin E. *Clinical Psychiatry: A Textbook for Students and Physicians*. New York: Macmillan, 1912.
- 11 Provencher HL. Role of psychological factors in studying recovery from a transactional stress-coping approach: implications for mental health nursing practice. *International Journal of Mental Health Nursing*, 2007; **16**: 188–197.
- 12 Repper J, Perkins R. *Social Inclusion and Recovery: A Model for Mental Health Practice*. London: Bailliere Tindall, 2003.

- 13 Coleman R. *Recovery: An Alien Concept*. Gloucester: Handsell Publishing, 1999.
- 14 Storm M, Edwards A. Models of user involvement in the mental health context: intentions and implementation challenges. *Psychiatric Quarterly*, 2013; **84**: 313–327.
- 15 Deegan PE, Drake RE. Shared decision making and medication management in the recovery process. *Psychiatric Services*, 2006; **57**: 1636–1639.
- 16 Cook JA, Copeland ME, Jonikas JA, Hamilton MM, Razzano LA, Grey DD, *et al.* Results of a randomised controlled trial of mental illness self-management using wellness recovery action planning. *Schizophrenia Bulletin*, 2012; **38**: 881–891.
- 17 Thornicroft G, Farrelly S, Szmuckler G, Birchwood M, Waheed W, Flach C, *et al.* Clinical outcomes of joint crisis plans to reduce compulsory treatment for people with psychosis: a randomised controlled trial. *The Lancet*, 2013; **381**: 1634–1641.
- 18 Waring J, Marshall F, Bishop S. Understanding the occupational and organisational boundaries to safe hospital discharge. *Journal of Health Services Research and Policy*, 2015; **20** (Suppl. 1): 35–44.
- 19 Gilbert H, Rose D, Slade M. The importance of relationships in mental health care: a qualitative study of service users experiences of psychiatric hospital admission in the UK. *BMC Health Services Research*, 2008; **8**: 92.
- 20 Department of Health. *Inpatients Formally Detained in Hospitals under the Mental Health Act 1983 and Patients Subject to Supervised Community Treatment – England 2010–2011 Annual Figures*. London: Department of Health, 2011.
- 21 Priebe S, Amos T, Leese M, Morriss R, Rose D, Wykes T, *et al.* Patients views and readmissions 1 year after involuntary hospitalisation. *The British Journal of Psychiatry*, 2009; **194**: 49–54.
- 22 Meehan J, Kapur N, Hunt IM, Turnbull P, Robinson J, Bickley H, *et al.* Suicide in mental health in-patients and within 3 months of discharge. National clinical survey. *The British Journal of Psychiatry*, 2006; **188**: 129–134.
- 23 Rowley E, Wright N, Waring J, Gregoriou K, Chopra A. Protocol for an exploration of knowledge sharing for improved discharge from a mental health ward. *BMJ Open*, 2014; **4**: e005176.
- 24 The Health Foundation. *Report: Improvement Science. Research Scan*. London: The Health Foundation, 2011.
- 25 Kruegar R, Casey MA. *Focus Groups: A Practical Guide for Applied Research*, 4th edn. Thousand Oaks: Sage Publications, 2008.
- 26 Kitzinger J. The methodology of focus groups: the importance of interaction between research participants. *Sociology of Health and Illness*, 1994; **16**: 103–121.
- 27 Burgess RG. *Field Research: A Sourcebook and Field Manual*. London: Allen & Unwin, 1982.
- 28 Ritchie J, Spencer L. Qualitative data analysis for applied policy research. In: Bryman A, Burgess RG (eds) *Analyzing Qualitative Data*. London: Routledge, 1994: 173–194.
- 29 Huberman A, Miles M. Data management and analysis methods. In: Denzin NK, Lincoln YS (eds) *Collecting and Interpreting Qualitative Materials*. Thousand Oaks: Sage Publications, 1998: 179–210.
- 30 Green J, Thorogood N. *Qualitative Methods for Health Research*. London: Sage Publications, 2004.
- 31 Tyrer P. Has the closure of psychiatric beds gone too far? Yes. *British Medical Journal*, 2011; **343**: d7457.
- 32 Quirk A, Lelliott P. What do we know about life on acute psychiatric wards in the UK? A review of the research evidence. *Social Science and Medicine*, 2001; **53**: 1564–1574.
- 33 Thornicroft G, Tansella M. Components of a modern mental health service: a pragmatic balance of community and hospital care. *The British Journal of Psychiatry*, 2004; **185**: 283–290.
- 34 Lauber C, Nordt C, Braunschweig C, Rossler W. Do mental health professionals stigmatize their patients? *Acta Psychiatrica Scandinavica*, 2006; **113**: 51–59.
- 35 Bowers L. *Dangerous and Severe Personality Disorder: Response and Role of the Psychiatric Team*. London: Routledge, 2002.
- 36 Quirk A, Lelliott P, Searl C. Service users' strategies for managing risk in the volatile environment of an acute psychiatric ward. *Social Science and Medicine*, 2004; **59**: 2573–2583.
- 37 Goffman E. *Asylums: Essays on the Social Situation of Mental Patients and Other Inmates*. Garden City: Anchor Publishing, 1961.
- 38 Wright D. Getting out of the asylum: understanding the confinement of the insane in the nineteenth century. *Social History of Medicine*, 1997; **10**: 137–155.
- 39 Quirk A, Lelliott P, Searl C. The permeable institution: an ethnographic study of three acute psychiatric wards in London. *Social Science and Medicine*, 2006; **63**: 2105–2117.
- 40 Department of Health. *New Ways of Working for Everyone: Developing and Sustaining a Capable and Flexible Workforce*. London: Department of Health, 2007.