

Set families with a forbidden pattern

Karpas, Ilan; Long, Eoin

DOI:

[10.1016/j.ejc.2016.11.005](https://doi.org/10.1016/j.ejc.2016.11.005)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Karpas, I & Long, E 2017, 'Set families with a forbidden pattern', *European Journal of Combinatorics*, vol. 62, pp. 183-196. <https://doi.org/10.1016/j.ejc.2016.11.005>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 05/09/2019

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Set families with a forbidden pattern

Ilan Karpas *

Eoin Long †

Abstract

A *balanced pattern* of order $2d$ is an element $P \in \{+, -\}^{2d}$, where both signs appear d times. Two sets $A, B \subset [n]$ form a P -pattern, which we denote by $\text{pat}(A, B) = P$, if $A \triangle B = \{j_1, \dots, j_{2d}\}$ with $1 \leq j_1 < \dots < j_{2d} \leq n$ and $\{i \in [2d] : P_i = +\} = \{i \in [2d] : j_i \in A \setminus B\}$. We say $\mathcal{A} \subset \mathcal{P}[n]$ is P -free if $\text{pat}(A, B) \neq P$ for all $A, B \in \mathcal{A}$. We consider the following extremal question: how large can a family $\mathcal{A} \subset \mathcal{P}[n]$ be if \mathcal{A} is P -free?

We prove a number of results on the sizes of such families. In particular, we show that for some fixed $c > 0$, if P is a d -balanced pattern with $d < c \log \log n$ then $|\mathcal{A}| = o(2^n)$. We then give stronger bounds in the cases when (i) P consists of $d +$ signs, followed by $d -$ signs and (ii) P consists of alternating signs. In both cases, if $d = o(\sqrt{n})$ then $|\mathcal{A}| = o(2^n)$. In the case of (i), this is tight.

1 Introduction

A central goal in extremal set theory is to understand how large a set family can be subject to some restriction on the intersections of its elements. Given $\mathcal{L} \subset \mathbb{N} \cup \{0\}$, we say that a set family \mathcal{A} is \mathcal{L} -intersecting if $|A \cap B| \in \mathcal{L}$ for all distinct $A, B \in \mathcal{A}$. Taking $\mathcal{L}_t = \{s \in \mathbb{N} : s \geq t\}$, a fundamental theorem of Erdős, Ko and Rado [6] shows that \mathcal{L}_t -intersecting families $\mathcal{A} \subset \binom{[n]}{k}$ satisfy $|\mathcal{A}| \leq \binom{n-t}{k-t}$, provided $n \geq n_0(k, t)$. Another important theorem due to Frankl and Füredi [8] shows that if $\mathcal{L}_{\ell, \ell'} := \{s < \ell \text{ or } s \geq k - \ell'\}$, then any $\mathcal{L}_{\ell, \ell'}$ -intersecting family $\mathcal{A} \subset \binom{[n]}{k}$ satisfies $|\mathcal{A}| \leq cn^{\max(\ell, \ell')}$, for some constant c depending on k, ℓ and ℓ' . See [2], [3], [7], [9] for an overview of this extensive topic.

Here we are concerned with understanding the effect of restricting the *pattern* formed between elements of a set family. A *difference pattern* or *pattern* of order t is an element $P \in \{+, -\}^t$. Given such a pattern P , let $S_+(P) = \{i \in [t] : P_i = +\} \subset [t]$ and $s_+(P) = |S_+(P)|$. Define $S_-(P)$ and $s_-(P)$ analogously. Two sets $A, B \subset [n]$ form a *difference pattern* P if:

- (i) $A \triangle B = \{j_1, \dots, j_t\}$ with $j_1 < \dots < j_t$, and
- (ii) $\{i \in [t] : P_i = +\} = \{i \in [t] : j_i \in A \setminus B\}$.

We denote this by writing $\text{pat}(A, B) = P$. A family of subsets $\mathcal{A} \subset \mathcal{P}[n]$ is P -free if $\text{pat}(A, B) \neq P$ for all distinct $A, B \in \mathcal{A}$. In this paper we consider the following natural question: given a pattern P , how large can a family $\mathcal{A} \subset \mathcal{P}[n]$ be if it is P -free?

*The Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem, Israel. Email: ilan.karpas@mail.huji.ac.il.

†School of Mathematical Sciences, Tel Aviv University, Tel Aviv, Israel. E-mail: eoinlong@post.tau.ac.il.

First note the following simple observation. If $s_+(P) \neq s_-(P)$ then large P -free families exist. Indeed, if $|s_+(P) - s_-(P)| = m > 0$ then the following families are P -free:

$$\mathcal{B}_1 = \{A \subset [n] : |A| \in [0, m-1] \pmod{2m}\}; \quad \mathcal{B}_2 = \{A \subset [n] : |A| \in [m, 2m-1] \pmod{2m}\}.$$

Clearly either $|\mathcal{B}_1| \geq 2^{n-1}$ or $|\mathcal{B}_2| \geq 2^{n-1}$. We will therefore focus on the case when $s_+(P) = s_-(P) = d$. We say that such patterns are d -balanced. For a balanced pattern P it is only possible that $\text{pat}(A, B) = P$ if $|A| = |B|$. Thus, our question on balanced patterns essentially reduces to a question for uniform families. Given $0 \leq k \leq n$, define

$$f(n, k, P) := \max \left\{ |\mathcal{A}| : P\text{-free families } \mathcal{A} \subset \binom{[n]}{k} \right\}.$$

Let $f(n, k, d) = \max\{f(n, k, P) : P \text{ is } d\text{-balanced}\}$. We will also write $\delta(n, k, P)$ and $\delta(n, k, d)$ for the corresponding extremal densities, i.e. $\delta(n, k, P) := f(n, k, P) / \binom{n}{k}$, and $\delta(n, k, d) := f(n, k, d) / \binom{n}{k}$. Note also that if $\mathcal{A} \subset \binom{[n]}{k}$ is P -free then the family $\mathcal{A}^c = \{[n] \setminus A : A \in \mathcal{A}\} \subset \binom{[n]}{n-k}$ is also P -free. Therefore $f(n, k, P) = f(n, n-k, P)$ and it suffices to bound $f(n, k, P)$ for $k \leq n/2$.

Our first aim is to prove a density result for d -balanced patterns of small order. That is, we will show that for fixed d , any sequence of integers $\{k_n\}_{n=1}^\infty$ tending to infinity with n with $k_n \leq n/2$ satisfies $\lim_{n \rightarrow \infty} \delta(n, k_n, d) = 0$. The condition that k is not fixed and tends to infinity with n will be crucial. This is different from the case in the Frankl-Füredi Theorem, which tells us that we can take some fixed $k \geq 2d-1$, $\ell = k-d$ and $\ell' = d-1$, and if $\mathcal{A} \subset \binom{[n]}{k}$ with $|\mathcal{A}| = \omega(n^{k-d})$ then there are $A, B \in \mathcal{A}$ with $|A \triangle B| = 2d$, i.e. A and B form a P -pattern for *some* d -balanced pattern P . Indeed, take any fixed $k := k(d)$, and consider the family $\mathcal{A}_0 \subset \binom{[n]}{k}$ given by

$$\mathcal{A}_0 = \left\{ A \subset [n] : \left| A \cap \left(\frac{(i-1)n}{k}, \frac{in}{k} \right] \right| = 1 \text{ for all } i \in [k] \right\}.$$

Then $|\mathcal{A}_0| \geq c_k n^k$ for some absolute constant $c_k > 0$, but it is easily seen that \mathcal{A}_0 does not contain the pattern $++--$. Therefore, there does not exist a density theorem for d -balanced patterns in subsets of $\binom{[n]}{k}$ with fixed k , as in the Frankl-Füredi theorem.

Our first result shows that such a density theorem does hold for k growing with n .

Theorem 1. *Given $d, k, n \in \mathbb{N}$ with $2k \leq n$ and taking $a_d = (8d)^{5d}$ and $c_d = 6d8^{-d}$ we have*

$$\delta(n, k, d) \leq a_d k^{-c_d}.$$

By our discussion above for fixed k we see that Theorem 1 is in a sense a ‘high-dimensional’ result. Also note that Theorem 1 shows there is a constant $c > 0$ with the property that if P is a d -balanced pattern with $d \leq c \log \log n$ and $\mathcal{A} \subset \mathcal{P}[n]$ which is P -free, then $|\mathcal{A}| = o(2^n)$.

Let $\text{IP}(d)$ denote the d -balanced pattern consisting of d plus signs, followed by d minus signs. We refer to these as *interval patterns*. Given the obstruction of $\text{IP}(2)$ above, it is natural to ask for bounds on $f(n, k, \text{IP}(d))$.

Theorem 2. *Given $d, k, n \in \mathbb{N}$ with $2k \leq n$ we have*

$$\delta(n, k, \text{IP}(d)) = O(d^2 k^{-1}).$$

In particular, families $\mathcal{A} \subset \mathcal{P}[n]$ which are $\text{IP}(d)$ -free for all $d = o(\sqrt{n})$ satisfy $|\mathcal{A}| = o(2^n)$. Furthermore, this turns out to be tight – if $d \geq c\sqrt{n}$ then there are $\text{IP}(d)$ -free families $\mathcal{A} \subset \mathcal{P}[n]$ with $|\mathcal{A}| = \Omega_c(2^n)$.

Lastly, we consider the d -balanced pattern $\text{AP}(d)$ consisting of alternating plus and minus signs, e.g. $\text{AP}(2) = + - + -$. We refer to these as *alternating patterns*. Our next result proves a density result for such patterns.

Theorem 3. *Given $d, k, n \in \mathbb{N}$ with $2k \leq n$ we have*

$$\delta(n, k, \text{AP}(d)) = O\left(\log^{-1}\left(\frac{k}{d^2}\right)\right).$$

Thus again, all families $\mathcal{A} \subset \mathcal{P}[n]$ which are $\text{AP}(d)$ -free for $d = o(\sqrt{n})$ satisfy $|\mathcal{A}| = o(2^n)$. Unlike in the case of the interval patterns, we do not know if this is tight.

Before closing the introduction, we mention some further results related to this topic. A family $\mathcal{A} \subset \mathcal{P}[n]$ is said to be a *tilted Sperner family* if for all distinct $A, B \in \mathcal{A}$ we have $|B \setminus A| \neq 2|A \setminus B|$. Equivalently, \mathcal{A} is P -free for all patterns P with $|S_-(P)| = 2|S_+(P)|$. Kalai raised the question of how large a tilted Sperner family $\mathcal{A} \subset \mathcal{P}[n]$ can be. In [13], Leader and the second author proved that such families satisfy $|\mathcal{A}| \leq (1 + o(1))\binom{n}{n/2}$, which is asymptotically optimal. For sufficiently large n , the extremal families were also determined. In [14], the second author proved that this bound almost still applies if we only forbid ‘tilted pairs’ A, B with a single pattern. It was shown that if $\mathcal{A} \subset \mathcal{P}[n]$ does not contain $A, B \in \mathcal{P}[n]$ with $|B \setminus A| \neq 2|A \setminus B|$ for all distinct $A, B \in \mathcal{A}$ and satisfying $a < b$ for all $a \in A \setminus B$ and $b \in B \setminus A$ then $|\mathcal{A}| \leq C\sqrt{\log n}\binom{n}{n/2}$, for some constant $C > 0$. This condition is equivalent to \mathcal{A} being $P(d)$ -free for all patterns $P(d)$ consisting of $d +$ signs followed by $2d -$ signs. This bound was recently improved by Gerbner and Vizer in [11]. They proved that such families satisfy $|\mathcal{A}| \leq C\sqrt{\log n}\binom{n}{n/2}$. No family is known for this problem with order more than $C\binom{n}{n/2}$.

Lastly, we mention a fascinating question raised by Johnson and Talbot [12] related to Theorem 1 (similar conjectures have been raised by Bollobás, Leader and Malvenuto [4], and Bukh [5]). Our phrasing slightly differs from that in [12].

Question (Johnson–Talbot). *Is it true that for any $k \in \mathbb{N}$ and $\alpha > 0$ there is $n_0(k, \alpha) \in \mathbb{N}$ with the following property. Suppose that $n \geq n_0(k, \alpha)$ and that $\mathcal{A} \subset \binom{[n]}{n/2}$ with $|\mathcal{A}| \geq \alpha\binom{n}{n/2}$. Then there are disjoint sets $S \in \binom{[n]}{n/2 - \lfloor k/2 \rfloor}$ and $T \in \binom{[n] \setminus S}{k}$ such that the family $\mathcal{C}_{T,S} := \{S \cup U : U \in \binom{T}{\lfloor k/2 \rfloor}\}$ is contained in \mathcal{A} .*

This is true for $k = 3$, but is already open for $k = 4$. In this case it is possible to guarantee that $|\mathcal{C}_{T,S} \cap \mathcal{A}| \geq 5$ for some T, S (note $|\mathcal{C}_{T,S}| = 6$ for $k = 4$). More generally, Johnson and Talbot [12] proved that under the hypothesis above, $|\mathcal{C}_{T,S} \cap \mathcal{A}| \geq 4 \cdot 3^{(k-4)/3}$ for some T, S . We note the conclusion that dense subsets of $\mathcal{P}[n]$ contain all small patterns (from Theorem 1) would immediately follow from a positive answer to this question. Indeed, for $k = 2d$ any set $\mathcal{C}_{T,S}$ contains every d -balanced pattern. Theorem 1 may be seen as giving (weak) evidence for the question: for $k = 2d$ and any d -balanced pattern P , there is T and S and sets $A, B \in \mathcal{C}_{T,S} \cap \mathcal{A}$ with $\text{pat}(A, B) = P$.

Notation: Given a set X , we write $\mathcal{P}(X)$ for the power set of X and $\binom{X}{k} = \{A \subset X : |A| = k\}$. Given integers $m, n \in \mathbb{N}$ with $m \leq n$, we let $[n] = \{1, \dots, n\}$ and $[m, n] = \{m, m+1, \dots, n\}$. We also write $(n)_m$ for the falling factorial $(n)_m = n(n-1) \cdots (n-m+1)$.

2 Small balanced patterns

In this section we prove Theorem 1. We will find it convenient to prove many of our results restricted of the middle layer. We then simply write $f(k, P)$ for $f(2k, k, P)$, $\delta(k, P)$ for $\delta(2k, k, P)$, etc.. The following simple observation is useful to move results between different layers of the cube.

Proposition 2.1. *Let $n, m, k, l \in \mathbb{N}$ with $m \leq n$, $l \leq k$ and $k + m - l \leq n$. Let P be a pattern. Then $\delta(n, k, P) \leq \delta(m, l, P)$.*

Proof. Suppose $\mathcal{A} \subset \binom{[n]}{k}$ is P -free with $|\mathcal{A}| = \delta(n, k, P) \binom{n}{k}$. Select two disjoint sets T and U of order m and $k - l$ uniformly at random (possible as $k + m - l \leq n$). Then let $\mathcal{A}_{T,U} = \{A \in \binom{[n]}{k} : A \cup U \in \mathcal{A}\}$. As \mathcal{A} is P -free, the set $\mathcal{A}_{T,U}$ must also be P -free for all T, U , giving $|\mathcal{A}_{T,U}| \leq \delta(m, l, P) \binom{m}{l}$. However, $\mathbb{E}_{T,U} |\mathcal{A}_{T,U}| = \delta(n, k, P) \binom{m}{l}$. The result follows. \square

Our next two lemmas are the main steps in the proof of Theorem 1. Combined they will allow a recursive bound for $\delta(k, d)$ based on bounds on $\delta(k', d')$ for $k' < k$ and $d' < d$.

Lemma 2.2. *Let $d, k \in \mathbb{N}$ with $k^{1/2} \geq 16 \log k$ and let P be a d -balanced pattern with $P_1 \neq P_{2d}$. Then given any $\gamma \in [\frac{16 \log k}{k^{1/2}}, 1]$ we have*

$$\delta(k, P) \leq \max \left(\gamma, 6\sqrt{\delta(\lceil \gamma^2 k / 64 \rceil, d-1)} \right).$$

Proof. Let γ be chosen as above and let $\mathcal{A} \subset \binom{[2k]}{k}$ be P -free with $|\mathcal{A}| = \alpha \binom{2k}{k}$. If $\alpha \leq \gamma$ then there is nothing to prove, so we will assume that $\alpha > \gamma \geq \frac{16 \log k}{k^{1/2}}$. We will first show that there are many pairs $A, B \in \mathcal{A}$ with $|A \triangle B| = 2$. Indeed, given $C \in \binom{[2k]}{k+1}$ let y_C denote the number of $A \in \mathcal{A}$ with $A \subset C$. Then we have

$$\sum_{C \in \binom{[2k]}{k+1}} y_C = |\{(A, C) \in \mathcal{A} \times \binom{[2k]}{k+1} : A \subset C\}| = |\mathcal{A}|k \geq \alpha k \binom{2k}{k+1}.$$

As for every pair $A, B \in \mathcal{A}$ with $|A \triangle B| = 2$ there is a unique set $C \in \binom{[2k]}{k+1}$ with $A, B \subset C$, we obtain

$$\begin{aligned} \left| \left\{ (A, B) \in \binom{\mathcal{A}}{2} : |A \triangle B| = 2 \right\} \right| &= \sum_{C \in \binom{[2k]}{k+1}} \binom{y_C}{2} \geq \binom{2k}{k+1} \binom{\alpha k}{2} \\ &\geq \frac{\alpha^2 k^2}{4} \times \frac{2k(2k-1)}{(k+1)k} \binom{2k-2}{k-1} \geq \frac{\alpha^2 k^2}{2} \binom{2k-2}{k-1}. \end{aligned} \quad (1)$$

The first inequality holds by the convexity of $\binom{x}{2}$ and the second since $\alpha k - 1 \geq \alpha k / 2$ as $\alpha \geq 2/k$.

Now given $1 \leq i < j \leq 2k$, let $\mathcal{A}_{i,j} := \{A \in \mathcal{A} : |A \triangle B| = 2\}$. Note that from (1) we have

$$\sum_{i < j} |\mathcal{A}_{i,j}| = \left| \{(A, B) \in \mathcal{A} \times \mathcal{A} : |A \triangle B| = 2\} \right| \geq \frac{\alpha^2 k^2}{2} \binom{2k-2}{k-1}. \quad (2)$$

Also let $\alpha_{i,j}$ and $\beta_{i,j}$ be defined so that $|\mathcal{A}_{i,j}| = \alpha_{i,j} \binom{2k-2}{k-1}$ and $\beta_{i,j} = (j-i)/2k$. By (2) we find $\{i,j\}$ with $\alpha_{i,j} \geq \frac{\alpha^2}{8}$ and $\beta_{i,j} \geq \frac{\alpha^2}{16}$. Indeed, we have

$$\sum_{\{i,j\}:\alpha_{i,j} < \frac{\alpha^2}{8}} |\mathcal{A}_{i,j}| + \sum_{\{i,j\}:\beta_{i,j} < \frac{\alpha^2}{16}} |\mathcal{A}_{i,j}| < \binom{2k}{2} \frac{\alpha^2}{8} \binom{2k-2}{k-1} + 2k \times \frac{\alpha^2}{16} 2k \binom{2k-2}{k-1} \leq \frac{\alpha^2 k^2}{2} \binom{2k-2}{k-1}.$$

Combined with (2) we see that a claimed pair $\{i,j\}$ exists. Fix such a pair $\{i,j\}$ and set $\mathcal{B} = \mathcal{A}_{i,j}$. Now let $X = [i+1, j-1]$ and $Y = [n] \setminus [i,j]$ so that $\mathcal{B} \subset \binom{X \cup Y}{k-1}$. Partition elements from $\binom{X \cup Y}{k-1}$ according to how they intersect X , for each $\ell \in [0, j-i-2]$ letting

$$X_\ell = \left\{ A \in \binom{X \cup Y}{k-1} : |A \cap X| = \ell \right\}.$$

Also let $\mathcal{B}_\ell = \mathcal{B} \cap X_\ell$ and $L = \left\{ \ell : \left| \ell - \frac{|X|}{2} \right| \leq \sqrt{|X| \log \left(\frac{8}{\alpha} \right)} \right\}$. By Chernoff's inequality we have

$$\sum_{\ell \notin L} |X_\ell| \leq \frac{\alpha^2}{32} \binom{|X| + |Y|}{k-1}.$$

Using that $|\mathcal{B}| = \alpha_{i,j} \binom{|X|+|Y|}{k-1} \geq \frac{\alpha^2}{16} \binom{|X|+|Y|}{k-1}$ this shows that

$$\sum_{\ell \in L} |\mathcal{B}_\ell| \geq |\mathcal{B}| - \frac{\alpha^2}{32} \binom{|X| + |Y|}{k-1} \geq \frac{\alpha^2}{32} \binom{|X| + |Y|}{k-1} \geq \frac{\alpha^2}{32} \sum_{\ell \in L} |X_\ell|.$$

The last inequality here holds since the sets X_ℓ are disjoint subsets of $\binom{X \cup Y}{k-1}$. Thus for some $\ell \in L$ we have $|\mathcal{B}_\ell| \geq \frac{\alpha^2}{32} |X_\ell|$. By averaging, we find a set $U \subset Y$ with $|U| = k - \ell - 1$ such that the family $\mathcal{C} = \{C \in \binom{X}{\ell} : C \cup U \in \mathcal{B}_\ell\}$ satisfies $|\mathcal{C}| \geq \frac{\alpha^2}{32} \binom{|X|}{\ell}$.

To complete the proof, let Q denote the pattern obtained from P by removing P_1 and P_{2d} , i.e. $Q = P_2 \cdots P_{2d-1}$. Note that as $P_1 \neq P_{2d}$ we see that Q is $(d-1)$ -balanced. We claim that \mathcal{C} is Q -free. Indeed, suppose $C_1, C_2 \in \mathcal{C}$ with $\text{pat}(C_1, C_2) = Q$. Then by definition of \mathcal{C} and $\mathcal{B} = \mathcal{A}_{i,j}$ we have

$$\left\{ C_a \cup U \cup \{h\} : a \in \{1, 2\}, h \in \{i, j\} \right\} \subset \mathcal{A}.$$

If $P_1 = +$ we find $\text{pat}(C_1 \cup U \cup \{i\}, C_2 \cup U \cup \{j\}) = P$. If $P_1 = -$ we find $\text{pat}(C_1 \cup U \cup \{j\}, C_2 \cup U \cup \{i\}) = P$. Thus \mathcal{C} must be Q -free and

$$\frac{\alpha^2}{32} \binom{|X|}{\ell} \leq |\mathcal{C}| \leq \delta(|X|, \ell, Q) \binom{|X|}{\ell}.$$

Take $k' = \lfloor \frac{|X|}{2} - \sqrt{|X| \log \left(\frac{8}{\alpha} \right)} \rfloor$. A calculation shows that $\frac{|X|}{4} \geq \sqrt{|X| \log \left(\frac{8}{\alpha} \right)} + 2$ since $\alpha \geq \frac{16 \log k}{k^{1/2}}$ and $|X| + 2 = \beta_{i,j} 2k \geq \frac{\alpha^2 k}{8}$. This gives

$$k' \geq \frac{|X|}{2} - \sqrt{|X| \log \left(\frac{8}{\alpha} \right)} - 1 \geq \frac{|X|}{4} + 1 \geq \left\lceil \frac{\beta_{i,j} k}{2} \right\rceil \geq \left\lceil \frac{\alpha^2 k}{64} \right\rceil \geq \left\lceil \frac{\gamma^2 k}{64} \right\rceil.$$

Since $\ell \in L$ we have $k' \leq \ell \leq |X| - k'$. Using Proposition 2.1 we find that $\frac{\alpha^2}{32} \leq \delta(|X|, \ell, d-1) \leq \delta(2k', k', d-1) = \delta(k', d-1) \leq \delta(\lceil \frac{\gamma^2 k}{64} \rceil, d-1)$. Rearranging this gives $\alpha \leq 6 \sqrt{\delta(\lceil \frac{\gamma^2 k}{64} \rceil, d-1)}$. \square

Our second lemma deals with the case where P starts and ends with the same signs.

Lemma 2.3. *Let $d \in \mathbb{N}$ and let P be a d -balanced pattern with $P_1 = P_{2d}$. Then there are $d_1, d_2 \geq 1$ with $d_1 + d_2 = d$ such that the following holds. For every k_1, k_2 with $2k_1 + k_2 = k$ we have*

$$\delta(k, P) \leq \max \left(2e^{-k_1/12}, 4\delta(k_1, d_1), 4(3k_1)^{2d_1} \delta(k_2, d_2) \right).$$

Similarly for every k_1, k_2 with $k_1 + 2k_2 = k$ we have

$$\delta(k, P) \leq \max \left(2e^{-k_2/12}, 4\delta(k_2, d_2), 4(3k_2)^{2d_2} \delta(k_1, d_1) \right).$$

Proof. To begin, for each $\ell \in [0, 2d]$ let

$$c_\ell = |\{j \in [\ell] : P_j = +\}| - |\{j \in [\ell] : P_j = -\}|.$$

As P is d -balanced and $P_1 = P_{2d}$, we have $c_{2d-1} = -c_1$. Combined with the fact that $c_0 = c_{2d} = 0$ and c_ℓ changes by exactly 1 as ℓ increases, we see that $c_{2d_1} = 0$ for some $1 \leq d_1 \leq d-1$. Setting $d_2 := d - d_1$ and $Q_1 = P_1 \cdots P_{2d_1}$, $Q_2 = P_{2d_1+1} \cdots P_{2d}$ it is easy to see that these patterns are d_1 -balanced and d_2 -balanced respectively.

Now suppose that $\mathcal{A} \subset \binom{[2k]}{k}$ with $|\mathcal{A}| = \alpha \binom{2k}{k}$ and that \mathcal{A} is P -free. We will prove the first bound above as the second bound is proved identically. We will assume that $\alpha \geq 2e^{-k_1/12}$ as otherwise there is nothing to show. Partition $[2k]$ into two consecutive intervals $I_1 = [3k_1]$ and $I_2 = [3k_1 + 1, 2k]$. For each $\ell \in I_1$ let $Z_\ell := \binom{I_1}{\ell} \times \binom{I_2}{k-\ell}$. Let $L = \left\{ \ell \in I_1 : |\ell - 3k_1/2| \leq \sqrt{3k_1 \log \left(\frac{2}{\alpha} \right)} \right\}$. Note that as $|\bigcup_{\ell \notin L} Z_\ell| \leq \frac{\alpha}{2} \binom{2k}{k}$ by Chernoff's inequality, we have $|\mathcal{A} \cap Z_\ell| \geq \frac{\alpha}{2} |Z_\ell|$ for some $\ell \in L$. Fix such a choice of ℓ and set $Z := Z_\ell$ and $\mathcal{B} = \mathcal{A} \cap Z_\ell$ so that $\mathcal{B} \subset Z$ with $|\mathcal{B}| \geq \frac{\alpha}{2} |Z|$.

We will now prove that α satisfies

$$\alpha \leq \max \left(4\delta(|I_1|, \ell, Q_1), 4|I_1|^{2d_1} \delta(|I_2|, k - \ell, Q_2) \right). \quad (3)$$

To see this, we may assume that $\alpha \geq 4\delta(|I_1|, \ell, Q_1)$ as otherwise there is nothing to show. Consider the set \mathcal{P}_{Q_1} given by

$$\mathcal{P}_{Q_1} = \left\{ (A, B) \in Z \times Z : \text{pat}(A \cap I_1, B \cap I_1) = Q_1 \text{ and } A \cap I_2 = B \cap I_2 \right\}.$$

We will first show that $|\mathcal{B} \times \mathcal{B}| \cap \mathcal{P}_{Q_1} \geq \frac{\alpha}{4|I_1|^{2d_1}} |\mathcal{P}_{Q_1}|$. Indeed, for each $D \in \binom{I_2}{k-\ell}$ let

$$\mathcal{E}(D) := \left\{ C \in \binom{I_1}{\ell} : C \cup D \in \mathcal{B} \right\}; \quad \mathcal{P}_{Q_1}(D) := \{C, C' \in \mathcal{E}(D) : \text{pat}(C, C') = Q_1\}.$$

Noting that each $\mathcal{C} \subset \binom{I_1}{\ell}$ with $|\mathcal{C}| > \delta(|I_1|, \ell, Q_1) \binom{|I_1|}{\ell}$ contains C, C' with $\text{pat}(C, C') = Q_1$, we find $|\mathcal{P}_{Q_1}(D)| \geq |\mathcal{E}(D)| - \delta(|I_1|, \ell, Q_1) \binom{|I_1|}{\ell}$. Combined these give

$$|\mathcal{B} \times \mathcal{B}| \cap \mathcal{P}_{Q_1} = \sum_{D \in \binom{I_1}{\ell}} |\mathcal{P}_{Q_1}(D)| \geq \sum_{D \in \binom{I_2}{k-\ell}} \left(|\mathcal{E}(D)| - \delta(|I_1|, \ell, Q_1) \binom{|I_1|}{\ell} \right) \geq \frac{\alpha}{4} |Z|, \quad (4)$$

The final inequality here holds since $\sum_{D \in \binom{I_2}{k-\ell}} |\mathcal{E}(D)| = |\mathcal{B}| \geq \frac{\alpha}{2}|Z|$ and $\alpha \geq 4\delta(|I_1|, \ell, Q_1)$. Lastly, using that $|\mathcal{P}_{Q_1}| \leq |I_1|^{2d_1}|Z|$ together with (4), we obtain $|(\mathcal{B} \times \mathcal{B}) \cap \mathcal{P}_{Q_1}| \geq \frac{\alpha}{4|I_1|^{2d_1}}|\mathcal{P}_{Q_1}|$.

Now, from this bound we find a choice of $C, C' \in \binom{I_1}{\ell}$ with $\text{pat}(C, C') = Q_1$ such that the set

$$\mathcal{F}_{C, C'} = \left\{ D \in \binom{I_2}{k-\ell} : C \cup D, C' \cup D \in \mathcal{A} \right\}$$

satisfies $|\mathcal{F}_{C, C'}| \geq \frac{\alpha}{4|I_1|^{2d_1}} \binom{n_2}{k-\ell}$. However, if $D, D' \in \mathcal{F}$ with $\text{pat}(D, D') = Q_2$ then $C \cup D, C' \cup D' \in \mathcal{A}$ and $\text{pat}(C \cup D, C' \cup D') = Q_1 Q_2 = P$. As \mathcal{A} is P -free we see $\mathcal{F}_{C, C'} \subset \binom{I_2}{k-\ell}$ is Q_2 -free. This gives $\frac{\alpha}{4|I_1|^{d_1}} \leq \delta(|I_2|, k-\ell, Q_2)$ and proves (3).

To complete the proof, note that as $\alpha \geq 2e^{-k_1/12}$, by definition of L we have $\ell \in L \subset [k_1, 2k_1]$ and $k-\ell \in [k_2, k_2+k_1]$. As $|I_1| = 3k_1$ and $|I_2| = 2k - 3k_1 = 2k_2 + k_1$, by Proposition 2.1 we find

$$\delta(|I_1|, \ell, Q_1) \leq \delta(2k_1, k_1, Q_1) = \delta(k_1, d_1) \quad \text{and} \quad \delta(|I_2|, k-\ell, Q_1) \leq \delta(2k_2, k_2, Q_2) = \delta(k_2, d_2).$$

Combined with (3) this completes the proof. \square

Proof of Theorem 1. We prove by induction on d that with $a_d = (8d)^{5d}$ and $c_d = 6d8^{-d}$ we have

$$\delta(k, d) \leq a_d k^{-c_d}. \quad (5)$$

For $d = 1$ we have $P = +- \text{ or } P = -+$ and $\mathcal{A} \subset \binom{[2k]}{k}$ is P -free simply means that $|A \triangle B| \neq 2$ for all distinct $A, B \in \mathcal{A}$. It is well known that such families satisfy $|\mathcal{A}| \leq \frac{1}{k} \binom{2k}{k}$. Indeed, for each $C \in \binom{[2k]}{k+1}$ let y_C denote the number of $A \in \mathcal{A}$ with $A \subset C$. Then

$$\sum_{C \in \binom{[2k]}{k+1}} y_C = |\{(A, C) \in \mathcal{A} \times \binom{[2k]}{k+1} : A \subset C\}| = |\mathcal{A}| \times k.$$

However, if $|A \triangle B| \neq 2$ for all distinct $A, B \in \mathcal{A}$ we must have $y_C \leq 1$ for all C . Rearranging, we obtain the claimed upper bound on $|\mathcal{A}|$. This easily gives that (5) holds for $d = 1$.

We now prove the result for a d -balanced pattern P , assuming by induction that the theorem holds for all d' -balanced patterns with $d' < d$. We can assume that $k \geq a_d^{1/c_d} \geq 16^8$ as otherwise the statement is trivial. We will first prove this when P begins and ends with different signs, using Lemma 2.2, noting that in this range $k^{1/2} \geq 16 \log k$. To apply this, let $\gamma = 8(a_{d-1})^{1/2} k^{-\frac{c_{d-1}}{4}}$ and note that $\gamma \geq 8(a_{d-1})^{1/2} k^{-\frac{1}{4}} \geq 16(\log k) k^{-1/2}$ since $k^{1/4}/\log k \geq 1/32 \geq 2(a_{d-1})^{-1/2}$. Therefore we can apply Lemma 2.2 to find

$$\begin{aligned} \delta(k, P) &\leq \max \left(\gamma, 6 \sqrt{\delta \left(\left\lceil \frac{\gamma^2 k}{64} \right\rceil, d-1 \right)} \right) \leq \max \left(8(a_{d-1})^{1/2} k^{-\frac{c_{d-1}}{4}}, 6 \sqrt{a_{d-1} (a_{d-1} k^{1-\frac{c_{d-1}}{2}})^{-c_{d-1}}} \right) \\ &\leq 8(a_{d-1})^{1/2} k^{-\frac{c_{d-1}}{4}} \leq a_d k^{-c_d}. \end{aligned}$$

The second inequality here uses that Lemma 2.2 holds for $d-1$ by induction, the third that $(a_{d-1})^{-c_{d-1}} \leq 1$ and $1 - \frac{c_{d-1}}{2} \geq \frac{1}{2}$ and the last inequality uses that $c_d \leq \frac{c_{d-1}}{4}$.

We now move to the case where P starts and ends with the same signs. Given P let d_1 and d_2 be as in Lemma 2.3 so that $d_1 + d_2 = d$ with $d_i \geq 1$. We will assume that $d_1 \leq d_2$ as the other case follows

similarly. Let us set $k_1 = \lceil k^\beta \rceil$ where $\beta = \frac{c_{d_2}}{2d_1 + c_{d_1}}$. Set $k_2 = k - 2k_1 \geq k - 4k^\beta \geq k - 4k^{1/2} \geq \frac{k}{2}$ for $k \geq 2^6$. Then by Lemma 2.3 we have

$$\begin{aligned} \delta(k, P) &\leq \max \left(2e^{-k_1/12}, 4\delta(k_1, d_1), 4(3k_1)^{2d_1} \delta(k_2, d_2) \right) \\ &\leq \max \left(2e^{-k^\beta/12}, 4a_{d_1} r^{-\beta c_{d_1}}, 4(6k^\beta)^{2d_1} a_{d_2} \left(\frac{k}{2}\right)^{-c_{d_2}} \right) \\ &\leq \max \left(2e^{-k^\beta/12}, 4a_{d_1} k^{-\beta c_{d_1}}, 8^{2d_1+3} a_{d_2} k^{2d_1\beta - c_{d_2}} \right) \\ &\leq \max \left(2e^{-k^\beta/12}, 8^{2d_1+3} a_{d_2} k^{-\frac{c_{d_1}c_{d_2}}{2d_1+c_{d_1}}} \right) \leq a_d k^{-c_d}. \end{aligned}$$

The first part of the final inequality here uses $a_d \geq 2k^{c_d}$ for $k \leq (a_d/2)^{1/c_d}$ and that $e^{-k^\beta/12} \leq k^{-c_d}$ for $k \geq (a_d/2)^{1/c_d}$. The second part uses that $8^{2d_1+3} a_{d_2} \leq a_d$ and that since $d = d_1 + d_2$ and $d \leq 2d_2$ we have $c_d \leq 12d_2 8^{-d} \leq \frac{36d_1 d_2 8^{-(d_1+d_2)}}{2d_1+1} \leq \frac{c_{d_1} c_{d_2}}{2d_1+1} \leq \frac{c_{d_1} c_{d_2}}{2d_1+c_{d_1}}$. This completes this case and the proof of the theorem. \square

3 Interval patterns

In this section, we first prove Theorem 2. We then give several lower bounds for the case $n = 2k$ depending on value of d .

3.1 Upper Bound on $\delta(n, n/2, \text{IP}(d))$

Proof of Theorem 2. Let $m = \lfloor \frac{n}{8d^2} \rfloor$. We partition $[n]$ into m intervals, $[n] = I_1 \cup \dots \cup I_m$ with $|I_i| = \lfloor 8d^2 \rfloor$ or $|I_i| = \lceil 8d^2 \rceil$ for all $i \in [m]$.

Consider the following way of choosing elements from $\binom{[n]}{n/2}$. First select a set $T \subset \binom{[n]}{n/2-d}$ uniformly at random. Let $J = \{i \in [m] : |I_i \setminus T| \geq d\}$. As $d < |I_i|/2$, for every $i \in [m]$ we have

$$\mathbb{P}(i \in J) = \mathbb{P}(|I_i \setminus T| \geq d) > \mathbb{P}(|I_i \cap T| \leq |I_i|/2) \geq \frac{1}{2}.$$

If $i \in J$, further select a set $S_i \subset \binom{I_i \setminus T}{d}$ uniformly at random, and set $A_i = T \cup S_i$. If $i \in [m] \setminus J$ simply set $A_i = \emptyset$.

Now for every $i, j \in J$ with $i < j$, we have $\text{pat}(A_i, A_j) = \text{IP}(d)$. Also for $i \notin J$ we have $A_i \notin \mathcal{A}$, since $|A_i| = 0 \neq n/2$. We conclude that there is at most one index $i \in [m]$ with $A_i \in \mathcal{A}$. Equivalently,

$$\sum_{i=1}^m \mathbf{1}_{A_i \in \mathcal{A}} = \sum_{i=1}^m \sum_{A \in \mathcal{A}} \mathbf{1}_{A_i = A} \leq 1.$$

This is true for any choice of T and S_i 's, so in particular if we take the expectation on both sides, we have

$$\sum_{i=1}^m \sum_{A \in \mathcal{A}} \mathbb{P}(A_i = A) \leq 1. \tag{6}$$

But as $A_i \notin \mathcal{A}$ for $i \notin J$, given any $A \in \mathcal{A}$ we get that $\mathbb{P}(A_i = A) = \mathbb{P}(A_i = A | i \in J) \mathbb{P}(i \in J) > \frac{1}{2} \mathbb{P}(A_i = A | i \in J)$. Rewriting (6), this gives

$$\sum_{i=1}^m \sum_{A \in \mathcal{A}} \frac{\mathbb{P}(A_i = A | i \in J)}{2} \leq 1. \quad (7)$$

Lemma 3.1. *Let $A \in \binom{[n]}{n/2}$ be a fixed set. If $|A \cap I_i| \geq \frac{|I_i|}{2} + d$, then $\mathbb{P}(A_i = A | i \in J) \geq \frac{1}{\binom{n}{n/2}}$.*

Proof. Indeed, $\mathbb{P}(A_i = A | i \in J) = \frac{N_i(A)}{N_i}$ where

$$N_i(A) := \left| \left\{ (S_i, T) : S_i \in \binom{I_i}{d}, T \in \binom{[n] \setminus S_i}{n/2 - d}, S_i \cup T = A \right\} \right|;$$

$$N_i := \left| \left\{ (S_i, T) : S_i \in \binom{I_i}{d}, T \in \binom{[n] \setminus S_i}{n/2 - d} \right\} \right|.$$

However, we have

$$\frac{N_i(A)}{N_i} \geq \frac{\binom{4d^2+d}{d} \binom{n-8d^2}{n/2-d}}{\binom{8d^2}{d} \binom{n-d}{n/2-d}} = \frac{(4d^2+d)d \left(\frac{n}{2}-d\right)! \frac{n!}{2!}}{(8d^2)d(n-d)!} \geq \frac{\left(\frac{n}{2}-d\right)! \frac{n!}{2!}}{2^d(n-d)!} > \frac{(n/2)!(n/2)!}{(n)!} = \frac{1}{\binom{n}{n/2}}.$$

□

For a set $A \in \binom{[n]}{n/2}$, denote by $G(A) = \left| \left\{ i \in [m] : |A \cap I_i| \geq \frac{|I_i|}{2} + d \right\} \right|$. From Lemma 3.1 it follows that for any given A , we have $\sum_{i=1}^m \mathbb{P}(A_i = A | i \in J) \geq G(A) \times \frac{1}{\binom{n}{n/2}}$. Together with (7), we obtain

$$\sum_{A \in \mathcal{A}} G(A) \leq 2 \binom{n}{n/2}. \quad (8)$$

We call a set $A \in \binom{[n]}{n/2}$ *bad*, if $G(A) < m/5$. Otherwise, we say that A is *good*. Let \mathcal{B} be the family of all bad sets.

Lemma 3.2. $|\mathcal{B}| = o\left(\frac{1}{m} \binom{n}{n/2}\right)$ for sufficiently large n .

Proof. For a uniform random choice of a set $A \subseteq \binom{[n]}{n/2}$, let X_i be a random variable, with $X_i = 1$ if $|A \cap I_i| > \frac{|I_i|}{2} + d$, and $X_i = 0$ otherwise. Let $Z = X_1 + \dots + X_m$. To prove the lemma, we need to show that $\mathbb{P}(Z < m/5) = o\left(\frac{1}{m}\right)$. By linearity of expectation, $\mathbb{E}Z = m\mathbb{E}X_i = m\mathbb{P}(X_i = 1)$. Notice that for every $i \neq j$, X_i and X_j are negatively correlated, since if A has many elements in one interval, it is less likely to have many elements on another interval.

$$\mathbb{P}(X_i = 0) = \frac{\sum_{i=0}^{4d^2+d} \binom{8d^2}{i} \binom{n-8d^2}{n/2-i}}{\binom{n}{n/2}} \leq \frac{1}{2} + \frac{\sum_{i=4d^2}^{4d^2+d} \binom{8d^2}{i} \binom{n-8d^2}{n/2-i}}{\binom{n}{n/2}} \leq \frac{1}{2} + \frac{d \binom{8d^2}{4d^2} \binom{n-8d^2}{n/2-4d^2}}{\binom{n}{n/2}} < 0.79.$$

The second inequality uses Stirling's formula. Therefore $\mathbb{P}(X_i = 1) = \mathbb{E}X_i > 0.21$. Using linearity of expectation gives $\mathbb{E}Z = \sum_{i=1}^m \mathbb{E}X_i > 0.21m$.

By a version of the Chernoff-Hoeffding bound for negatively correlated variables [15], we deduce that $\mathbb{P}(A \in \mathcal{B}) = \mathbb{P}(Z < 0.2m) < \mathbb{P}(Z - \mathbb{E}Z > 0.01m) = o\left(\frac{1}{m}\right)$, finishing the proof. □

Therefore, if $|\mathcal{A}| \geq \frac{2}{m} \binom{n}{n/2}$, then $|\mathcal{A} \setminus \mathcal{B}| = (1 - o(1))|\mathcal{A}|$. Using (8), we see that

$$(1 - o(1)) \frac{m|\mathcal{A}|}{10} \leq \sum_{A \in \mathcal{A} \setminus \mathcal{B}} G(A) \leq \sum_{A \in \mathcal{A}} G(A) \leq \binom{n}{n/2}. \quad (9)$$

Equivalently $|\mathcal{A}| = O(\frac{1}{m} \binom{n}{n/2}) = O(\frac{d^2}{n} \binom{n}{n/2})$, as required. \square

3.2 Lower Bound on $\delta(n, n/2, \text{IP}(d))$

For the lower bounds, we provide different lower bounds, depending on the range of d .

Theorem 4. *The following hold:*

(i) *If $d = o(\sqrt{n})$, there is an $\text{IP}(d)$ -free family $\mathcal{A} \subseteq \binom{[n]}{n/2}$ with $|\mathcal{A}| = \Omega(\max\{\frac{1}{nd}, \frac{d^2}{n^{3/2}}\} \cdot \binom{n}{n/2})$.*

(ii) *If $d = c\sqrt{n}$, there is an $\text{IP}(d)$ -free family $\mathcal{A} \subseteq \binom{[n]}{n/2}$ with $|\mathcal{A}| = \Omega_c(\binom{n}{n/2})$.*

Proof. First we prove (i). For a set $A \in \binom{[n]}{n/2}$ let $S(A) := \sum_{i \in A} i$, the sum of the elements in A . Observe that if $\text{pat}(A, B) = \text{IP}(d)$ then $0 < |S(A) - S(B)| < nd$. Thus for any $0 \leq i \leq nd - 1$, the family $\mathcal{A}_i := \{A \in \binom{[n]}{n/2} \mid S(A) \equiv i \pmod{nd}\}$ forms an $\text{IP}(d)$ -free family. By the pigeonhole principle, we can find such i so that $|\mathcal{A}_i| \geq \frac{1}{nd} \binom{n}{n/2}$.

To obtain the second bound from (i), note that if we choose a set $A \in \binom{[n]}{n/2}$ uniformly at random,

$$\mathbb{E}[S(A)] = \frac{n(n+1)}{4}. \quad (10)$$

To calculate the variance, let

$$X_i = \begin{cases} 1 & \text{if } i \in A \\ 0 & \text{if } i \notin A. \end{cases}$$

Then $S(A) = \sum_{i=1}^n iX_i$. Now $\mathbb{E}[X_i] = \frac{1}{2}$ every $i \in [n]$ and $\mathbb{E}[X_i X_j] = \frac{1}{4}(1 - \frac{1}{n-1})$. Using this, we find

$$\begin{aligned} \text{Var}(S(A)) &= \mathbb{E}[(\sum_{i=1}^n iX_i)^2] - \mathbb{E}[\sum_{i=1}^n iX_i]^2 \leq \sum_{i \in [n]} i^2 \mathbb{E}[X_i] + \sum_{i \neq j} ij (\mathbb{E}[X_i X_j] - \mathbb{E}[X_i] \mathbb{E}[X_j]) \\ &\leq \sum_{i \in [n]} \frac{i^2}{2} \leq \frac{n^3}{2}. \end{aligned} \quad (11)$$

From (10) and (11) together, by Chebyshev's inequality we get $\mathbb{P}(|S(A) - n(n+1)/4| \leq n^{3/2}) \geq 1/2$. Equivalently, $|\{A \in \binom{[n]}{n/2} : |S(A) - n(n+1)/4| \leq n^{3/2}\}| \geq \frac{1}{2} \binom{n}{n/2}$. By an easy averaging argument, for some value $m \in [\frac{n(n+1)}{4} - \frac{n^{3/2}}{3}, \frac{n(n+1)}{4} + \frac{n^{3/2}}{2}]$.

$$|\{A \in \binom{[n]}{n/2} : S(A) \in [m - \frac{d^2}{2}, m + \frac{d^2}{2}]\}| \geq \frac{1}{2(2n^{3/2}/d^2 + 2)} \binom{n}{n/2} = \Omega\left(\frac{d^2}{n^{3/2}} \binom{n}{n/2}\right)$$

However, since two sets $A, B \in \binom{[n]}{n/2}$ with $\text{pat}(A, B) = \text{IP}(d)$ have $|S(A) - S(B)| > d^2$, this completes the proof of (i).

To prove (ii), let $c > 0$ be given and let $d = c\sqrt{n}$. Note that if $\text{pat}(A, B) = IP(d)$ then for some $i \in [n]$ we have $|A \cap [i]| \geq |B \cap [i]| + d$. This shows that $\mathcal{A} = \{A \in \binom{[n]}{n/2} : ||A \cap [i]| - i/2| < d/4 \text{ for all } i \in [n]\}$ is an $IP(d)$ -free family. We will now show that $|\mathcal{A}| = \Omega_c(\binom{n}{n/2})$.

To see this, it is convenient to identify elements of $\binom{[n]}{n/2}$ with certain walks. Let \mathcal{W}_0 denote the set of all walks $W = W_0 \cdots W_n$ of length n on \mathbb{Z} with $W_0 = W_n = 0$ and which either increase or decrease by 1 in each step (i.e. $|W_i - W_{i-1}| = 1$ for all $i \in [n]$). Note that each walk $W \in \mathcal{W}_0$ naturally corresponds to a subset of $[n]$ of size $n/2$ consisting of those steps in $[n]$ where the walk increases. Under this correspondence, the set \mathcal{A} corresponds to those walks in \mathcal{W}_0 which lie entirely in $[-d/4, d/4]$.

Now select a walk $W \in \mathcal{W}_0$ uniformly at random. Letting T denote a value to be determined, consider the following events:

$$\begin{aligned} A &= \{W_j \in [-d/4, d/4] \text{ for all } j \in [n]\} \\ B &= \{W_{in/T} \in [-d/12, d/12] \text{ for all } i \in [T-1]\} \\ C_i &= \{W_j \in [-d/4, d/4] \text{ for all } j \in [\frac{(i-1)n}{T}, \frac{in}{T}]\}, \text{ where } i \in [T]. \end{aligned}$$

Also for $i \in [T-1]$ and $a_i \in [-d/12, d/12]$, let $B_i(a_i)$ denote the event $B_i(a_i) = \{W_{in/T} = a_i\}$. We will show that

$$\mathbb{P}_{W \sim \mathcal{W}_0} \left(B \wedge \bigwedge_{i \in [T]} C_i \right) \geq c' > 0, \quad (12)$$

where c' depends only on c . Since $\bigwedge_{i \in [T]} C_i \subset A$, this will prove the result.

To begin, note that we have

$$\begin{aligned} \mathbb{P}_{W \sim \mathcal{W}_0} \left(B \wedge \bigwedge_{i \in [T]} C_i \right) &\geq \sum_{a_1, \dots, a_{T-1} \in [-d/12, d/12]} \mathbb{P}_{W \sim \mathcal{W}_0} \left(\bigwedge_{i \in [T-1]} B_i(a_i) \wedge \bigwedge_{i \in [T]} C_i \right) \\ &= \sum_{a_1, \dots, a_{T-1} \in [-d/12, d/12]} \mathbb{P}_{W \sim \mathcal{W}_0} \left(\bigwedge_{i \in [T]} C_i \mid \bigwedge_{i \in [T-1]} B_i(a_i) \right) \\ &\quad \times \mathbb{P}_{W \sim \mathcal{W}_0} \left(\bigwedge_{i \in [T-1]} B_i(a_i) \right). \end{aligned} \quad (13)$$

Let $\mathcal{W}(a, b)$ denote the collection of random walks of length n/T which start at a and end at b . Since C_i depends only on $\{W_j : j \in [(i-1)n/T, in/T]\}$, taking $a_0 = a_T = 0$ we have

$$\begin{aligned} \mathbb{P}_{W \sim \mathcal{W}_0} \left(\bigwedge_{i \in [T]} C_i \mid \bigwedge_{i \in [T-1]} B_i(a_i) \right) &= \prod_{i \in [T]} \mathbb{P}_{W \sim \mathcal{W}_0} \left(C_i \mid B_{i-1}(a_{i-1}) \wedge B_i(a_i) \right) \\ &= \prod_{i \in [T]} \mathbb{P}_{W \sim \mathcal{W}(a_{i-1}, a_i)} \left(W \text{ lies entirely in } [-d/4, d/4] \right). \end{aligned} \quad (14)$$

Claim: For every $a, b \in [-d/12, d/12]$ we have $\mathbb{P}_{W \sim \mathcal{W}(a, b)} \left(W \text{ lies entirely in } [-d/4, d/4] \right) \geq 1/2$.

Let $\mathcal{W}(a)$ denote the collection of all walks of length n/T which begin at a . Let us select W from $\mathcal{W}(a)$ uniformly at random and let $S_{n/T}$ denote the final vertex. By the reflection principle for random

walks, we have

$$\begin{aligned}
\mathbb{P}_{W \sim \mathcal{W}(a,b)}(W \text{ exceeds } d/4) &= \mathbb{P}_{W \sim \mathcal{W}(a)}(W \text{ exceeds } d/4 | S_{n/T} = b) \\
&= \frac{\mathbb{P}_{W \sim \mathcal{W}(a)}(S_{n/T} = d/2 - b)}{\mathbb{P}_{W \sim \mathcal{W}(a)}(S_{n/T} = b)} \\
&= \frac{\binom{n/T}{n/2T + (d/2 - b) - a}}{\binom{n/T}{n/2T + b - a}} \leq \frac{\binom{n/T}{n/2T + d/3}}{\binom{n/T}{n/2T + d/6}} \\
&= \frac{(n/2T - d/6)_{d/6}}{(n/2T + d/6)_{d/6}} \leq \left(1 - \frac{dT}{3n}\right)^{d/6} \leq e^{-d^2 T / 36n}.
\end{aligned}$$

Taking $T = 72/c^2$ say, we find $\mathbb{P}_{W \sim \mathcal{W}(a,b)}(W \text{ exceeds } d/4) \leq e^{-2} < 1/4$. By symmetry, this gives $\mathbb{P}_{W \sim \mathcal{W}(a,b)}(W \text{ lies entirely in } [-d/4, d/4]) \geq 1 - 2 \times (1/4) = 1/2$, as claimed.

Now by combining (14) together with the claim in (13) we find

$$\mathbb{P}_{W \sim \mathcal{W}_0}\left(B \wedge \bigwedge_{i \in [T]} C_i\right) \geq \sum_{a_1, \dots, a_{T-1} \in [-d/12, d/12]} 2^{1-T} \times \mathbb{P}_{W \sim \mathcal{W}_0}\left(\bigwedge_{i \in [T-1]} B_i(a_i)\right). \quad (15)$$

But letting $b_i := \frac{n}{2T} + a_i - a_{i-1}$ for all $i \in [T]$ where $a_0 = a_T = 0$, we have

$$\mathbb{P}_{W \sim \mathcal{W}_0}\left(\bigwedge_{i \in [T-1]} B_i(a_i)\right) = \frac{\prod_{i \in [T]} \binom{n/T}{b_i}}{\binom{n}{n/2}} = \Omega_{c,T}(d^{1-T}).$$

The final inequality follows by Stirling's approximation, using that $b_i \in [\frac{n}{2T} - \frac{d}{6}, \frac{n}{2T} + \frac{d}{6}]$ for all $i \in [T]$. Combined with (15), this gives $\mathbb{P}_{W \sim \mathcal{W}_0}(B \wedge \bigwedge_{i \in [T]} C_i) = \Omega_{c,T}(1) = \Omega_c(1)$, as required. \square

4 Alternating patterns

To begin, we prove an auxiliary lemma. Given $\mathbf{x} = (x_i)$ and $\mathbf{y} = (y_i)$ in $[m]^D$ we say that \mathbf{y} d -dominates \mathbf{x} if $|\{i \in [D] : x_i \neq y_i\}| = d$ and $x_i \leq y_i$ for all $i \in [D]$.

Lemma 4.1. *Let $d, m, D \in \mathbb{N}$ with $2md^2 \leq D$. Suppose that $\mathcal{C} \subset [m]^D$ does not contain \mathbf{x} and \mathbf{y} such that \mathbf{y} d -dominates \mathbf{x} . Then $|\mathcal{C}| \leq 2m^{D-1}$.*

Proof. To begin, choose a set $S \subset [D]$ with $|S| = d$ and a vector $\mathbf{z} \in [m]^{[D] \setminus S}$ uniformly at random. For each $i \in [m]$ let $\mathbf{z}_S(i) \in [m]^D$ denote the vector which agrees with \mathbf{z} on coordinates in $[D] \setminus S$ and equals i everywhere else. Also let $\mathcal{B}_{S,\mathbf{z}}$ denote the combinatorial line $\mathcal{B}_{S,\mathbf{z}} := \{\mathbf{z}_S(i) : i \in [m]\}$.

Now as \mathcal{C} does not contain any d -dominating pairs, for any choice of S and \mathbf{z} we have $|\mathcal{C} \cap \mathcal{B}_{S,\mathbf{z}}| \leq 1$. Letting X_i denote the indicator random variable which is 1 if $\mathbf{z}_S(i) \in \mathcal{C}$ and 0 otherwise, this gives

$$\sum_{i \in [m]} X_i \leq 1.$$

Taking expectations over all choice of S and \mathbf{z} , this gives

$$\sum_{C \in \mathcal{C}} \mathbb{P}(C \in \mathcal{B}_{S,\mathbf{z}}) = \sum_{i \in [m]} \sum_{C \in \mathcal{C}} \mathbb{P}(\mathbf{z}_S(i) = C) \leq 1. \quad (16)$$

However, an easy calculation gives that if C has k_i entries i for all $i \in [m]$, then

$$\mathbb{P}(C \in \mathcal{B}_{S,\mathbf{z}}) = \sum_{i \in [m]} \frac{\binom{k_i}{d}}{m^{D-d} \binom{D}{d}}.$$

This expression is minimized when all k_i are as equal as possible. Thus

$$\begin{aligned} \mathbb{P}(C \in \mathcal{B}_{S,\mathbf{z}}) &\geq m \frac{\binom{D/m}{d}}{m^{D-d} \binom{D}{d}} = m \frac{(D/m)_d}{m^{D-d} D_d} = \frac{m}{m^D} \prod_{l \in [0, d-1]} \left(1 - \frac{l(m-1)}{D-l}\right) \\ &\geq \frac{1}{m^{D-1}} \left(1 - \sum_{l \in [0, d-1]} \frac{l(m-1)}{D/2}\right) \\ &\geq \frac{1}{m^{D-1}} \left(1 - \frac{md^2}{D}\right) \geq \frac{1}{2m^{D-1}}. \end{aligned}$$

The final line here used $2md^2 \leq D$. Combined with (16) this gives $|\mathcal{C}|/2m^{D-1} \leq 1$, as required. \square

We are now ready for the proof of Theorem 3.

Proof of Theorem 3. By Proposition 2.1 it suffices to prove the theorem for $n = 2k$. Let $m = \lfloor \frac{\log_2(n/d^2)}{2} \rfloor$. For convenience we assume that n is divisible by m , with $Km = n$. Let $[n] = \bigcup_{i=1}^K I_i$ be a partition of $[n]$ where $I_i = \{(i-1)m+1, \dots, im\}$ for all $i \in [K]$. Given a set $T \subset [K]$, let $T^c = [K] \setminus T$ and let

$$\mathcal{B}_T := \{A \subset \bigcup_{i \in T^c} I_i : |A \cap I_i| \neq 1 \text{ for all } i \in T^c\}.$$

Given $B \in \mathcal{B}_T$ and $\mathbf{x} \in [m]^T$ we also let $B(\mathbf{x}) := B \cup \{(i-1)m+j-1 : i \in T, x_i = j\}$ and

$$\mathcal{C}_B := \{B(\mathbf{x}) : \mathbf{x} \in [m]^T\}.$$

Note that for every $A \subset [n]$ there is a unique $T \subset [K]$, $B \in \mathcal{B}_T$ and $\mathbf{x} \in [m]^T$ such that $A = B(\mathbf{x})$. Thus we have the disjoint union

$$\binom{[n]}{n/2} = \bigcup_{T \subset [K]} \bigcup_{\substack{B \in \mathcal{B}_T \\ |B| = \frac{n}{2} - |T|}} \mathcal{C}_B. \quad (17)$$

We will first show that almost all sets A in $\binom{[n]}{n/2}$ are of the form $A = B(\mathbf{x})$ where $T \subset [K]$ and $B \in \mathcal{B}_T$ with $|T| \geq mK/2^{m+1} = n/2^{m+1}$. To see this, given a set $A \subset [n]$, let $A_i = A \cap I_i$ for all $i \in [K]$. We will say that $A \subset [n]$ is *bad* if $T(A) = \{i \in [K] : |A_i| = 1\}$ satisfies $|T(A)| \leq \frac{m}{2^{m+1}} K$. We claim that there are at most $O(e^{-n^{1/2}/2} 2^n)$ sets are bad. Indeed, if we select $A \subset [n]$ uniformly at random, we have $\mathbb{P}(|A_i| = 1) = m/2^m$, which gives $\mathbb{E}(|T(A)|) = \frac{mK}{2^m} = \frac{n}{2^m}$. As $|A_i| = 1$ for each $i \in [K]$ independently, by Chernoff's inequality, we find that $\mathbb{P}(|T(A)| - \frac{n}{2^m} \leq -\frac{n}{2^{m+1}}) \leq e^{-\frac{n}{2^{m+1}}}$. As $m \leq \log_2(n/d^2)/2 \leq \frac{\log_2 n}{2}$ we find that $\mathbb{P}(A \text{ is bad}) \leq e^{-n^{1/2}/2}$. Equivalently, $|\{A \subset [n] : A \text{ is bad}\}| = O(e^{-n^{1/2}/2} 2^n)$.

Now suppose that $T \subset [K]$ with $|T| \geq n/2^{m+1}$ and $B \in \mathcal{B}_T$. Note that given $\mathbf{x}, \mathbf{y} \in [m]^T$, if \mathbf{y} d -dominates \mathbf{x} then $\text{pat}(B(\mathbf{x}), B(\mathbf{y})) = \text{AP}(d)$. Noting that as $m = \lfloor \log_2(n/d^2)/2 \rfloor$ we have $|T| \geq n/2^{m+1} \geq 2^m d^2 \geq 2md^2$. Setting $D = |T|$, Lemma 4.1 therefore shows that any $\mathcal{A} \subset \binom{[n]}{n/2}$ which is $\text{AP}(d)$ -free satisfies

$$|\mathcal{A} \cap \mathcal{C}_B| \leq 2m^{|T|-1} = \frac{2}{m} |\mathcal{C}_B|. \quad (18)$$

Summing over all $T \subset [K]$ and $B \in \mathcal{C}_T$, combined with (17) and (18), this gives

$$\begin{aligned} |\mathcal{A}| &\leq \sum_{T \subset [K]} |\mathcal{A} \cap \bigcup_{\substack{B \in \mathcal{B}_T \\ |B|=n/2-|T|}} \mathcal{C}_B| \leq |\{A \subset [n] : A \text{ bad}\}| + \sum_{\substack{T \subset [K]: \\ |T| \geq 2md^2}} \sum_{B \in \mathcal{B}_T} |\mathcal{A} \cap \mathcal{C}_B| \\ &\leq O\left(\frac{2^n}{e^{\sqrt{n}/2}}\right) + \sum_{\substack{T \subset [K]: \\ |T| \geq 2md^2}} \sum_{\substack{B \in \mathcal{B}_T \\ |B|=n/2-|T|}} \frac{2}{m} |\mathcal{C}_B| \\ &\leq \frac{2+o(1)}{m} \binom{n}{n/2}. \end{aligned}$$

This completes the proof of the theorem. \square

5 Concluding remarks and open problems

In this paper we proved bounds on the size of families $\mathcal{A} \subset \mathcal{P}[n]$ which avoid a d -balanced pattern P . Our proof shows that such families satisfy

$$|\mathcal{A}| = O(a_d n^{-c_d} 2^n),$$

where $a_k = (8d)^{5d}$ and $c_d = 6d8^{-d}$. In particular, families \mathcal{A} which avoid a d -balanced pattern with $d < c \log \log n$ satisfy $|\mathcal{A}| = o(2^n)$ for some absolute constant $c > 0$. It would be interesting to improve the density bound here and/or extend the range of d for which this zero density property holds.

Another interesting question is the following: which balanced pattern P has the strongest effect on the density of P -free families $\mathcal{A} \subset \mathcal{P}[n]$? That is, what is $\min_P \delta(n, k, P)$, where the minimum is taken over all balanced patterns P ? If instead of patterns we only forbid intersection sizes (as discussed in the Introduction) then there are a number of very strong density results for subsets of $\mathcal{P}[n]$. For example, the Frankl-Rödl [10] theorem shows that given $\epsilon > 0$, if $\mathcal{A} \subset \mathcal{P}[n]$ and $|A \cap B| \neq t$ for some $\epsilon n \leq t \leq (1/2 - \epsilon)n$ then $|\mathcal{A}| \leq (2 - \delta)^n$, where $\epsilon = \epsilon(\delta) > 0$. It would be very interesting to know if there exists a pattern which forces a superpolynomial density in n . That is, does there an increasing sequence of naturals $(n_k)_{k \in \mathbb{N}}$ and balanced patterns $(P_k)_k$ with $\delta(n_k, n_k/2, P_k) = n_k^{-\omega_k(1)}$ for some function $\omega_k(1)$ tending to infinity with k ?

Lastly, how large can d be (as a function of n) while still giving $\delta(n, n/2, \text{AP}(d)) \rightarrow 0$ as $n \rightarrow \infty$. Theorem 3 proves that this holds for any $d = o(\sqrt{n})$.

References

- [1] N. Alon and J. Spencer: **The Probabilistic Method**, Wiley, 3rd ed., 2008.

- [2] L. Babai and P. Frankl: **Linear algebra methods in combinatorics**, Department of Computer Science, University of Chicago, preliminary version, second edition, September 1992.
- [3] B. Bollobás: **Combinatorics. Set systems, hypergraphs, families of vectors and combinatorial probability**, Cambridge University Press, 1986.
- [4] B. Bollobás, I. Leader and C. Malvenuto: Daisies and other Turán problems. *Combin. Probab. Comput.* **20** (5) (2011), 743-747.
- [5] B. Bukh: Set families with a forbidden subposet, *Electron. J. Combin.* **16**(1) (2009), Paper 142.
- [6] P. Erdős, C. Ko and R. Rado: Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford Ser. (2)* **12** (1961), 313-320.
- [7] P. Frankl: Intersection theorems for finite sets and geometric applications, *Proceedings of the International Congress of Mathematicians*, Berkeley, USA, 1986.
- [8] P. Frankl and Z. Füredi: Forbidding just one intersection, *J. Combin. Theory Ser. A*, **39**(2) (1985), 160-176.
- [9] Z. Füredi: Turán type problems, *Surveys in Combinatorics*, London Math. Soc. Lecture Note Ser. 166, Cambridge Univ. Press, Cambridge, 1991, 253-300.
- [10] P. Frankl and V. Rödl: Forbidden intersections, *Trans. Amer. Math. Soc.*, **300**(1) (1987), 259-286.
- [11] D. Gerbner and M. Vizer: A note on tilted Sperner families with patterns, <http://arxiv.org/abs/1507.02242>
- [12] J.R. Johnson and J. Talbot: Vertex Turán problems in the hypercube, *J. Combin. Theory Ser. A* **117**(4) (2010), 454-465.
- [13] I. Leader and E. Long, Tilted Sperner families, *Discrete Appl. Math.*, **163** (2014), 194-198.
- [14] E. Long, Forbidding intersection patterns between layers of the cube, *J. Combin. Theory Ser. A*, **134** (2015), 103-120.
- [15] A. Panconesi and A. Srinivisan, Randomized distributed edge coloring via an extension of the Chernoff-Hoeffding bounds, *SIAM Journal on Computing*, **26**(2) (1997), 350-368.