

Environmental reconstruction of a later prehistoric palaeochannel record from Burrs Countryside Park, Bury, Greater Manchester

Smith, David; Fletcher, Mark; Head, Katie; Smith, Wendy; Howard, Andy

DOI:

[10.1179/146141010X12640787648658](https://doi.org/10.1179/146141010X12640787648658)
[10.1179/146141010X12640787648658](https://doi.org/10.1179/146141010X12640787648658)

Document Version

Early version, also known as pre-print

Citation for published version (Harvard):

Smith, D, Fletcher, M, Head, K, Smith, W & Howard, A 2010, 'Environmental reconstruction of a later prehistoric palaeochannel record from Burrs Countryside Park, Bury, Greater Manchester', *Environmental Archaeology*, vol. 15, no. 1, pp. 16-31. <https://doi.org/10.1179/146141010X12640787648658>,
<https://doi.org/10.1179/146141010X12640787648658>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Environmental reconstruction of a later prehistoric palaeochannel record from Burrs Countryside Park, Bury, Greater Manchester

David N. Smith, Mark Fletcher, Katie Head, Wendy Smith and Andy J. Howard

The results from an analysis of pollen, insect and plant macrofossil remains from the shallow wood peat fill of a palaeochannel associated with hummocky (deglacial) sediments infilling the valley of the River Irwell at Walmersley known locally as Burrs Countryside Park, Bury, Greater Manchester (NGR SD 796 127) are presented. Radiocarbon dates from the deposit provide an age range from 1310 to 1040 cal. BC to 920 to 800 cal. BC. The environmental record indicates that during the Middle and Late Bronze Age the valley floor was covered by carr woodland consisting mainly of alder and hazel, including areas of open water. Insect and pollen remains indicate that mixed deciduous woodland (birch, elm, beech and ash) was present on the surrounding slopes. Although non-arboreal pollen remains low and may merely indicate woodland edge grassland communities, increases in the percentage of oak and birch pollen over time may reflect small-scale clearance of the alder carr in the valley bottom. Alternatively, if the pollen diagram is merely recording a decline in alder pollen at the top of the profile, this could also explain the increase in the relative proportion of other arboreal taxa. There are no taxa present directly indicative of human activities associated with the nearby Castle Steads hillfort. The palaeoenvironmental results from this site appear to match those for the Greater Manchester area and wider north-west region in general, suggesting that there was a mosaic of environments present in the Bronze Age, with some areas cleared of woodland earlier than others.

Keywords: Greater Manchester, Bronze Age, palaeoenvironments, insects, pollen

Introduction

The late prehistoric archaeology of the north-west of England, particularly the area around Greater Manchester, is relatively under-explored when compared to other areas of Great Britain. This paucity of data is particularly acute for the Late Bronze Age and Iron Age in the region. This is despite clear evidence for Bronze Age settlement from the Manchester Airport site, and the occurrence of a number of large hillforts in the area during the Late Iron Age, for

example, Mellor at Stockport, Castle Steads at Bury, Beeston Castle, Cheshire and Eddisbury, Cheshire.

This pattern is particularly true for palaeoenvironmental studies in this region. Hall and Huntley (2007, 51) and Kenward (2009) in the English Heritage regional reviews for this area stress the rarity of palaeoenvironmental sites of Bronze Age date east of the Pennines and in greater Manchester specifically. Major studies are limited to the English Heritage-funded North West Wetlands Survey for Greater Manchester, centred on a study of pollen at Chat Moss Mire as well a number of smaller mires (Wells *et al.* 1993; Hall *et al.* 1995). Despite this activity, the number of Bronze Age and Iron Age sites with palaeoenvironmental investigation in the region still remains relatively small and only provides broad and disparate data with which to attempt landscape

David Smith (corresponding author) and A. J. Howard, Institute of Archaeology and Antiquity, The University of Birmingham, Edgbaston, Birmingham B15 2TT, UK; e-mail: D.N.SMITH@bham.ac.uk; M. Fletcher, Matrix Archaeology Ltd, Kenwood House, 10 Kenwood Rd, Stretford, Manchester M32 8PT, UK; K. Head, School of Geography, University of Plymouth, Davy Building, Drake Circus, Plymouth, Devon PL4 8AA, UK; W. Smith, Oxford Archaeology South, Janus House, Osney Mead, Oxford OX2 0ES, UK




Figure 1 Location of Bury, Walmersley, Greater Manchester, UK

reconstruction. As a result of this paucity of data in the region, Hall and Huntley (2007, 254) suggest that all wetland sites west of the Pennines and dating to the earlier prehistoric warrant specific attention.

Taken out of context, the site described here could be seen as ‘parochial’ and the research undertaken as somewhat limited since only a relatively small period of time is considered. The results are somewhat unspectacular, largely confirming our views of the Late Bronze Age/Early Iron Age environment of this region. However, given the discussion above, it is clear that even such a limited study as this is of regional importance for Greater Manchester and helps to fill what are in fact previously ‘blank’ time periods.

This paper describes the results of an environmental analysis of a shallow deposit of Middle to Late Bronze Age wood peat associated with hummocky (deglacial) sediments infilling the valley of the River Irwell at Walmersley, Bury, Greater Manchester (NGR SD 796 127) (Fig. 1). In addition to the opportunity to explore natural vegetation

succession and landscape change, the position of the sampling site, directly below the small promontory hillfort at Castle Steads, potentially provided the opportunity to reconstruct the environmental setting of this area in the period prior to Late Iron Age to Romano-British human occupation.

Background

The work results from commercial trial trenching undertaken (under PPG16) by Matrix Archaeology, prior to groundworks related to the Burrs Caravan Park, and funded by the Caravan Club. The site was located in an area of pastureland 2 km to the north of Bury town centre, and lies at the base of the slope below the Castle Stead Hillfort. A series of radiocarbon dates on charcoal from the fort ditches has produced a combined age range of 550 cal. BC to 247 cal. AD (Beta-56798; Beta-58075; Beta-58076; Beta-58077: 1-sigma – Fletcher 1986; 1992). The peat-filled palaeochannel appeared to have developed as an oxbow and, until the 1990s, was regularly flooded after heavy rain. At c. 91 m OD, the palaeochannel



Figure 2 The excavation in progress at Bury showing the depth of the deposit excavated

was elevated about 6 m above the level of the present riverbed. The trial-trench was located just 100 m to the south of the Castle Steads promontory, which rises to a height of 114.50 m OD.

An organic-rich 'wood peat' infilled the palaeo-channel hollow associated with the hummocky terrace sediments aggraded during deglaciation at the end of the last cold stage (the Devensian). These Quaternary sediments rest upon shales and fine sandstones of the Carboniferous Coal Measures. The soils developed upon these parent materials comprise either groundwater gleys or freely draining brown-earths (Mackney *et al.* 1983; Fletcher 1986).

Environmental assessment of the sedimentary sequence (pollen, insect and macroscopic plant remains) confirmed the potential of this deposit to yield proxy records for climate and land-use (Head 2005; Smith and Smith 2005). Radiocarbon dating of the uppermost palaeochannel sediments (0–10 cm) yielded an age estimate of 920 to 800 cal. BC (Beta-210417: 2 sigma) and dating of the basal deposits (40–50 cm) yielded an age estimate of 1310 to 1040 cal. BC (Beta-210418: 2 sigma). This indicated that the organic sequence extended from the Middle well into the Late Bronze Age.

Methodology and results

As the organic-rich sequence was relatively thin (0.5 m) (see Fig. 2), environmental samples were removed in 10 cm spits. Since the dates from the upper and lower units were chronologically close, it was decided to concentrate full environmental

analysis on three of the five samples (# 1, 0–10 cm; # 3, 20–30 cm; # 5, 40–50 cm).

Pollen

Sediment samples of 2 cm³ were measured volumetrically. The samples were washed in 10% hydrochloric acid and then digested by 10% potassium hydroxide for 20 minutes in a boiling water bath to break up the soil matrix and dissolve any humic material. As the samples contained a large amount of organic matter, they were acetolysed for three minutes to break down the cellulose material. Finally, the pollen pellet was stained with safranin, washed in alcohol to dehydrate the sample, and preserved in silicon oil.

Pollen grains were counted to a total of 500 land pollen grains (TLP) and analysis was undertaken on a GS binocular polarising microscope at 400× magnification. Identification was aided by using the pollen reference collection maintained by Worcestershire County Council Historic Environment and Archaeology Service and the reference manual by Moore *et al.* (1991). Nomenclature follows Stace (1997) and Bennett (1994). The pollen diagram (Fig. 3) was constructed using TILIA, TILIA.GRAPH and TGView 2.0.2 software (Grimm 1990; 2004). The diagram was not divided into pollen assemblage zones in this case, as only three samples were analysed.

Insects

The samples were processed using the standard method of paraffin flotation as outlined in Kenward




Figure 3 Pollen percentage diagram for Bury

et al. (1980). Each processed sample was of 20 litres. The insect remains present were sorted from the flots and stored in ethanol. The Coleoptera (beetles) present were identified under a low-power binocular microscope at magnifications between $\times 10$ – $\times 45$ and by direct comparison to the Gorham and Girling Collections of British Coleoptera held at the University of Birmingham. The records are presented in Table 1, where the taxonomy for the Coleoptera (beetles) follows that of Lucht (1987). Where applicable, each taxon has been assigned to one or more ecological grouping(s) and these are indicated in the second column of Table 1. These groupings are derived from the preliminary classifications outlined by Robinson (1981; 1983) and are described at the end of Table 1. The various proportions of these groups, expressed as percentages of the total Coleoptera present in the faunas, are shown in Table 2 and Fig. 4. The dung/fowl, tree, grassland and moorland groupings are calculated as a proportion of the terrestrial taxa recovered rather than as a proportion of the minimum number of individuals for the whole fauna (effectively excluding the dominant water beetles from this statistic). Column 6 in Table 1 indicates modern rarity of certain of the recovered taxa. The scheme used follows the Red Data Book (RDB) classifications of Hyman and Parsons (1992; 1994). Column 7 in Table 1 presents the host plants for the various species of phytophage (plant feeding) beetles recovered. The information included is primarily taken from Koch (1992). The plant nomenclature follows that of Stace (1997).

Macroscopic plant remains

Sub-samples of 500 ml were washed over a 0.3 mm sieve and all of the material retained was sorted at

$\times 12$ magnification. Identifications were made at magnifications up to $\times 40$, in comparison with the Institute of Archaeology and Antiquity reference material, as well as in consultation with standard keys (Beijerinck 1976; Schoch *et al.* 1988; Cappers *et al.* 2006). Nomenclature follows Stace (1997). A list of taxa recovered from all three samples and a list of taxa with a specific habitat(s) are illustrated in Tables 3 and 4 respectively. A summary interpretation of the waterlogged plant macrofossil data is provided in Table 5.

The reconstructed landscape present at Bury during the Middle and Late Bronze age

The three environmental proxies studied (pollen, insects and plant macrofossil remains) all suggest that a similar environment existed through out the sequence

Indicators for woodland

All three proxy indicators clearly suggest that woodland formed a dominant aspect of the landscape. Arboreal pollen (trees and shrubs) accounts for 85% of the terrestrial land pollen (TLP) in the basal sample with a slight decline to 74% by the upper sample (Fig. 3). Woodland is represented by ecological group '1' for the Coleoptera data presented in Tables 1 and 2 and Fig. 4 and accounts for approximately 25% of the terrestrial fauna recorded in all three samples. Values such as this for woodland indicators in the archaeoentomological record often are used to suggest the presence of a closed canopy (Robinson 1981; 1983; Whitehouse and Smith 2004; Smith and Whitehouse 2005).

The pollen suggests that, at least locally, this woodland mainly consisted of alder (*Alnus glutinosa*

Table 1 The insects recovered from Bury

	1	3	5	RDB status	Phytophage plant hosts(from Koch 1992)
HEMIPTERA					
Family, gen. et spp. indet.	+	++	++		
COLEOPTERA					
Carabidae					
<i>Nebria brevicollis</i> (F.)	1	-	-		
<i>Clivina fossor</i> (L.)	1	-	-		
<i>Trechus quadristriatus</i>	1	-	-		
(Schrk.) or <i>T. obtusus</i> Er.					
<i>Trechoblemus micros</i> (Hbst.)	1	-	-		
<i>Bembidion</i> spp.	1	-	-		
<i>Patrobis septentrionis</i> Dej.	1	-	-		
<i>Pterostichus diligens</i> (Sturm)	ws	2	1		
<i>Pterostichus nigrita</i> (Payk.)	ws	2	-		
<i>Pterostichus anthracinus</i> (Ill.)	ws	-	-		
<i>Pterostichus oblongopunctatus</i> (F.)	t	-	-	nb	
<i>Pterostichus aethiops</i> (Panz.)	t	-	-	nb	
<i>Abax parallelepipedus</i> (Pill. Mitt)	t	-	-		
<i>Agonum fuliginosum</i> (Panz.)	ws	2	-		
<i>Platynus assimilis</i> (Payk.)	t	-	1		
Halipidae					
<i>Halipilus</i> sp.	1	-	1		
Dytiscidae					
<i>Coelambus impressopunctatus</i>	1	-	-		
(Schall.)					
<i>Hydroporus palustris</i> (L.)	a	-	5		
<i>Hydroporus</i> spp.	a	-	3		
<i>Stictotarsus duodecimpustulatus</i> (F.)	a	-	1		
<i>Agabus bipustulatus</i> (L.)	3	1	-		
<i>Agabus sturmi</i> (Gyll.)	6	-	-		
<i>Agabus</i> spp.	a	4	1		
<i>Ilybius ater</i> (Geer)	a	-	2		
<i>Ilybius fuliginosus</i> (F.)	a	-	4		
<i>Dytiscus</i> spp.	a	1	-		
Hydraenidae					
<i>Hydraena britteni</i> Joy	4	-	-		
<i>H.</i> spp.	48	4	2		
<i>Ochthebius</i> spp.	a	-	2		
<i>Limnebius</i> spp.	a	-	-		
<i>L.</i> spp.	a	1	5		
<i>Helophorus</i> spp.	a	-	-		
Hydrophilidae					
<i>Coelostoma orbiculare</i> (F.)	a	2	1		
<i>Megasternum boletophagum</i> (Marsh.)	a	1	1		
<i>Laccobius</i> spp.	a	-	1		
<i>Enochrus</i> sp.	a	-	1		
<i>Hydrobius fuscipes</i> (L.)	a	2	4		

Table 1 Continued

		1	3	5	RDB status	Phytophage plant hosts(from Koch 1992)
Histeridae						
<i>Abraeus globosus</i> (Hoffm.)	df	1	–	–		
Silphidae						
<i>Silpha</i> spp.		1	–	–		
Ptiliidae						
<i>Acrotrichis</i> spp.		1	1	1		
Liodidae						
<i>Agathidium</i> spp.	t	1	–	–		
Scydmaenidae						
<i>Scydmaenidae</i> gen. et spp. indet.		2	–	–		
Staphylinidae						
<i>Proteinus</i> spp.		5	–	1		
<i>Eusphalerum</i> spp.		3	1	3		
<i>Phyllodrepa loptera</i> (Steph.)		4	–	2		
<i>Omalius</i> spp.		1	–	1		
<i>Lathrimaemum unicolor</i> (Marsh.)	ws	1	1	–		
<i>Olophrum piceum</i> (Gyll.)	ws	4	–	3		
<i>Acidota crenata</i> (F.)	ws	–	1	1		
<i>Lesteva heeri</i> (Fauv.)	ws	6	2	–		
<i>Lesteva longelytrata</i> (Goeze)	ws	1	–	1		
<i>Lesteva pubescens</i> Mannh.	ws	1	–	–		
<i>Lesteva</i> spp.		–	1	1		
<i>Trogophloeus</i> spp.	ws	1	–	–		
<i>Oxytelus rugosus</i> (F.)	df	2	–	–		
<i>Stenus</i> spp.		20	1	1		
<i>Lathrobium</i> spp.		4	1	–		
<i>Gyrophynus fracticornis</i> (Müll)		1	–	–		
<i>Xantholinus</i> spp.		2	–	–		
<i>Philonthus</i> spp.		5	2	1		
<i>Bolitobius</i> spp.		1	–	–		
<i>Tachyporus</i> sp.		1	–	–		
<i>Tachinus rufipes</i> (Geer)		1	–	–		
<i>Aleocharinidae</i> gen. et spp. indet.		4	2	1		
Pselaphidae						
<i>Rybaxis</i> spp.		–	1	–		
<i>Brachygluta</i> spp.		1	–	–		
Cantharidae						
<i>Cantharis rufa</i> L.		1	–	–		
<i>Cantharis pallida</i> Goeze		1	–	1	nb	
Elatridae						
<i>Melanotus rufipes</i> (Hbst.)	t	1	–	–		
<i>Athous haemorrhoidalis</i> (F.)	g	1	–	–		
Eucnemidae						
<i>Melasis buprestoides</i> (L.)	t	–	1	–	nb	Dead timber in old broadleaf woodland
Throscidae						

Table 1 Continued

	1	3	5	RDB status	Phytophage plant hosts(from Koch 1992)
<i>Throscus</i> spp.					
Dascillidae	g	–	1		
<i>Dascillus cervinus</i> (L.)	g	–	1		
Helodidae					
Helodidae gen. & spp. indet. (? <i>Cyphon</i> spp.)	ws	2	32		
Dryopidae					
<i>Elmis aenea</i> (Müll)	a	–	–		
<i>Esolus parallelepipedus</i> (Müll)	a	–	1		
<i>Limnius volckmari</i> (Panz.)	a	–	1		
Nitidulidae					
<i>Epuraea</i> sp.	g	–	1		
Rhizophagidae					
<i>Rhizophagus bipustulatus</i> (F.)	t	–	1		Usually under bark of a range of broadleaf trees
Cryptophagidae					
<i>Cryptophagus</i> spp.	1	–	–		
Lathridiidae					
<i>Lathridius minutus</i> (Group)	1	–	–		
<i>Corticaria</i> sp.	–	1	–		
Colydiidae					
<i>Cerylon</i> spp.	t	–	–		
Coccinellidae					
<i>Chilocorus renipustulatus</i> (L.)	t	–	1		normally on willow (<i>Salix</i> spp.)
Cisidae					
<i>Cis</i> sp.	t	–	2		
Anobiidae					
<i>Grynobius planus</i> (F.)	t	1	–		
<i>Anobium punctatum</i> (Geer)	t	–	–		
Ptinidae					
<i>Ptinus</i> spp.	1	3	–		
Mordellidae					
<i>Anaspis</i> spp.	2	–	2		
Scarabaeidae					
<i>Geotrupes</i> spp.	d	1	–		
<i>Aegialia sabuleti</i> (Panz.)	d	–	–	nb	
<i>Aphodius fimetarius</i> (L.)	d	–	–		
Lucanidae					
<i>Sinodendron cylindricum</i> (L.)	t	–	–		Dead wood of a range of broadleaf trees
Cerambycidae					
<i>Grammoptera</i> spp.	t	–	–		
Chrysomelidae					
<i>Donacia versicolore</i> (Brahm.)	ws	–	–		On pond weed (<i>Potamogeton</i> spp.)
<i>Donacia vulgaris</i> Zschach.	ws	–	1		On bur-reed (<i>Sparganium</i> spp.), sedges (<i>Carex</i> spp.) and bullreed (<i>Typhaea</i> spp.)
<i>Plateumaris sericea</i> (L.)	ws	–	2		Usually on sedges (<i>Carex</i> spp.)
<i>Lema</i> spp.	g	–	–		

Table 1 Continued

		1	3	5	RDB status	Phytophage plant hosts(from Koch 1992)
<i>Chrysomela</i> sp.	g	1	-	-		
<i>Phaedon</i> spp.	g	2	-	-		
<i>Prasocuris phellandrii</i> (L.)	ws	3	-	3		On aquatic cow parsleys (APIACEAE)
<i>Gastroidea viridula</i> (Geer)	g	-	-	1		Usually on dock (<i>Rumex</i> spp.)
Scolytidae						
<i>Scolytus scolytus</i> (F.)	t	1	-	-		Usually on elm (<i>Ulmus</i> spp.)
<i>Ieperisinus varius</i> (F.)	t	1	-	-		Usually associated with ash (<i>Fraxinus excelsior</i> L.)
<i>Dryocoetes alni</i> (Georg)	t	1	-	-		Usually on alder (<i>Alnus</i> spp.)
<i>Xyloterus domesticus</i> (L.)	t	1	-	-		Usually associated with beech (<i>Fagus sylvatica</i> L.)
Curculionidae						
<i>Deporaus betulae</i> (L.)	t	3	1	-		Usually birch (<i>Betula</i> spp.)
<i>Phyllobius</i> spp.	t	2	-	1		
<i>Strophosoma</i> spp.		-	-	1		
<i>Tanyssphyrus lemnae</i> (Payk.)	ws	1	1	-		On duck weed (<i>Lemna</i> spp.)
<i>Notaris</i> spp.	ws	-	1	-		
<i>Micrelus ericae</i> (Gyll.)	m	1	-	-		Usually on heather (<i>Erica</i> spp./ <i>Calluna</i> spp.)
<i>Rhynchaenus</i> spp.	t	1	-	1		
<i>Rhamphus pulicarius</i> (Hbst.)	t	2	-	-		On willow (<i>Salix</i> spp.)
DIPTERA						
<i>Cyclorhapha</i> gen. et spp. indet.		+++				
TRICOPTERA						
Gen. et spp. Indet.		+++	+++	+++		

Ecological groupings

a – aquatic species
ws – waterside species either from muddy riverbanks/shores or from waterside vegetation
df – species associated with dung and foul matter
d – species associated with dung
g – species associated with grassland and pasture
t – species either associated with trees or with woodland in general – species associated with moorland and acid bogs
For non Coleoptera the numbers present have been estimated using the following scale:
+ =>10 individuals
++ =<10 individuals
+++ =<20 individuals
++++ =<50 individuals




Figure 4 The proportions of the ecological groups of Coleoptera recovered from Bury

(L.) Gaertn) and hazel (*Corylus avellana*-type) (Fig. 2). There is an increase in the representation of alder in the middle of the sequence. The overall dominance of alder is also evident in both the insect and plant macrofossils recovered from this middle deposit. The presence of the beetle *Dryocoetes alni*, which is normally associated with alder (Koch 1992) is considered significant. Normally, alder is under-represented in the archaeoentomological record (Girling 1985; Smith et al. 2000; Smith and Whitehouse 2005). The presence of other trees, most likely as part of the alder carr, is indicated by the insects *Chilocorus renipustulatus* and *Rhamphus pulicarius*, which are associated with willow (*Salix* spp. — Koch 1992), and by the seeds of downy birch (*Betula pubescens* Ehr.). The insect remains suggest that deadwood (such as from fallen trunks and tree limbs) was present. This is represented by taxa such as *Agathidium*, *Melanotus rufipes*, *Rhizophagus bipustulatus*, *Grynobius planus*, *Anobium punctatum*, *Sinodendron cylindricum* and *Grammoptera* sp. Several of the ground beetles recovered are associated with leaf litter and forest soils, including *Pterostichus*

oblongopunctatus, *P. aethiops*, *Abax parallelepipedus* and *Platynus assimilis* (Lindroth 1974). Wood-sorrel (*Oxalis acetosella* L.), a well-known plant of shaded and/or woodland environments (Packham 1978), was recovered in the plant macrofossil assemblage; however, this taxon can occur elsewhere, such as along hedgerbanks (e.g. Stace 1997, 475) or in rough grazing, especially on hill slopes (Packham 1978, 680).

More mixed woodland probably grew on the higher valley sides. Limited indicators for elm (*Ulmus* spp.), oak (*Quercus* spp) and silver birch (*Betula pendula* Roth), as well as occasional examples of ash (*Fraxinus excelsior* L.), lime (*Tilia cordata* Mill.) ivy (*Hedera helix* L.) and pine (*Pinus* spp.) are all found in the pollen, plant macrofossil and insect records. The pollen record suggests that there is a minor decline in the occurrence of alder and elm pollen towards the upper part of the sequence.

Indicators for open areas and cultivated ground

There is little evidence for the presence of substantial clearance or cultivated ground in the pollen record; both herb and heath pollen only account for 15% to 20% of the terrestrial land pollen in each sample. This is a minor aspect of the pollen profiles, which is dominated by grass (*Poaceae* undiff.) with only the occasional presence of other herbs primarily meadow buttercup (*Ranunculus acris*-type), common sorrel (*Rumex acetosa* L.) and meadowsweet (*Filipendula* spp.). There is a slight increase in the proportions of these species towards the top of the sequence, perhaps suggesting an increase in clearings in the area, although this simply could indicate localised

Table 2 The relative proportions of the ecological groups of Coleoptera recovered from Burrs Countryside Park, based on MNI

Sample number	1	3	5
Number of individuals	293	48	107
Number of species	95	31	47
% aquatic	37.2%	33.3%	33.6%
% waterside	24.9%	27.1%	40.2%
% dung and dung foul (of terrestrial)	6.3%	5.3%	0.0%
% grassland (of terrestrial)	6.3%	0.0%	14.3%
% trees (of terrestrial)	25.2%	15.8%	25.0%
% moorland (of terrestrial)	0.0%	0.0%	3.6%

damp/wet ground. Classic indicators for clearance such as ribwort plantain (*Plantago lanceolata* L.) and self-heal (*Prunella vulgaris*-type), however, appear only in the upper sample, supporting an open grassland interpretation; either cleared by humans or reflecting woodland-edge grassland communities.

Conversely, the presence of areas of cleared ground, meadow or pasture is only indicated in the insect faunas from the lower two samples. The evidence for this consists of a few individuals of *Geotrupes* and *Aphodius* ‘dung beetles’, the ‘click

beetle’ *Athous haemorrhoidalis*, which is associated with grassland, and the ‘leaf beetle’ *Gastroidea viridula*, which is associated with docks. However, this ecological group only represents a very small proportion of this insect fauna (see ecological group ‘p’ in Table 2 and Fig. 4) and probably suggests that clearings (not necessarily created through human agency) were limited at this time. One possible explanation for the discrepancy between the archaeoentomological and palynological data may be a decline in alder pollen in the upper sample; this would

Table 3 Waterlogged plant remains from Bronze Age palaeochannel deposits at Bury

Sample number	1	3	5	
Sample location in column	0–10 cm	20–30 cm	40–50 cm	
Latin Binomial				English Common Name
<i>Urtica dioica</i> L.	–	1	–	Common Nettle
cf. <i>Urtica dioica</i> L.	1	–	–	?Common Nettle
<i>Betula pendula</i> Roth	–	1	5	Silver Birch
<i>Betula pubescens</i> Ehr.	–	5	12	Downy Birch
<i>Betula</i> spp. – indeterminate	18	64	37	Birch
<i>Betula</i> spp. – catkin	–	–	1	Birch
cf. <i>Betula</i> spp. – buds	–	2	–	Birch
<i>Alnus glutinosa</i> (L.) Gaertn.	9	481	317	Alder
<i>Alnus glutinosa</i> (L.) Gaertn. – infrustructure	–	1	4	Alder
cf. <i>Alnus glutinosa</i> (L.) Gaertn.	–	10	–	Possible Alder
<i>Stellaria media</i> (L.) Vill. agg.	–	185	–	Common Chickweed
<i>Cerastium</i> spp.	3	–	–	Mouse-ear
CARYOPHYLLACEAE – <i>Silene</i> type	–	–	1	Pink Family
cf. <i>Salix</i> sp. – bud	2	1	–	Possible Willow Bud
<i>Filipendula ulmaria</i> (L.) Maxim.	–	–	2	Meadowsweet
<i>Rubus</i> spp.	1	6	–	Bramble/Blackberry
cf. <i>Sanguisorba minor</i> Scop. ssp. minor	2	2	–	Possible Salad Burnet
<i>Melilotus/Medicago/Trifolium</i> sp.	–	1	–	Melilot/Medick/Clover
<i>Medicago</i> sp. – pod fragment	–	10	–	Medick
<i>Myriophyllum</i> sp.	–	1	–	Water-milfoil
<i>Oxalis acetosella</i> L.	2	11	1	Wood-sorrel
<i>Galeopsis</i> sp.	–	–	1	Hemp-nettle
<i>Lycopus europaeus</i> L.	1	101	2	Gypsywort
LABIACEAE – unidentified	1	–	1	Mint Family
<i>Callitriche</i> spp.	96	132	123	Water-starwort
<i>Campanula</i> sp.	2	–	–	Bellflower
<i>Galium</i> sp.	–	1	–	Bedstraw
ASTERACEAE – unidentified	–	–	1	Daisy Family
<i>Alisma</i> cf. <i>plantago-aquatica</i> L.	1	50	452	Possible Water-plantain
<i>Potamogeton</i> spp.	–	–	3	Pondweed
<i>Juncus</i> spp.	25	289	5	Rush
<i>Carex</i> spp. – 2-sided urticale	38	386	8	Sedge
<i>Carex</i> spp. – small 3-sided urticale	–	60	–	Sedge
POACEAE – indeterminate large grass caryopsis	1	110	–	Large-seeded Grass
POACEAE – indeterminate medium grass caryopsis	33	–	7	Medium-seeded Grass
POACEAE – indeterminate small grass caryopsis	12	25	3	Small-seeded Grass
Unidentified – anther	–	–	3	Anther
Unidentified – buds	1	2	27	Bud
Unidentified – large, elongated bud	–	3	5	Large, Elongated Bud
Unidentified – bud scars	–	4	1	Bud Scar
Unidentified – fruit/bud	–	4	–	Fruit/Bud
Unidentified – leaf fragment	1	–	–	Leaf
UNIDENTIFIED	3	6	3	Unidentified
TOTAL IDENTIFICATIONS	253	1955	1025	
OTHER REMAINS OBSERVED				
Bryophyte – unidentified fragments	–	–	+	Moss
<i>Daphnia</i> spp.	+	++	++	Waterflea

Nomenclature follows Stace (1997). Unless otherwise stated, counts are for seeds (in the widest sense). Key: + =<5 items, ++ =5–20 items, +++ >20 items

Table 4 Habitats of waterlogged plant macrofossils from Bronze Age deposits at Burrs Countryside Park (Habitat information follows Stace 1997)

HABITAT	WATER SIDE	WATER- DAMP GROUND	FENS/ MOORS	HEATH- LAND	GRASS- LAND	CULTIV- ATED GROUND	ROUGH GROUND	WASTE/ WOOD- LAND	HEDGE- BANKS	WOOD- LAND	ROCKY PLACES	OPEN GROUND (UN- SHADED)	SHADED PLACES	ACID SOILS	CALCA- REOUS/ BASE- RICH SOIL	NITR- OGEN RICH SOIL	NEU- TRAL SOIL	English Common Name
Latin Binomial																		
AQUATIC TAXA																		
<i>Myriophyllum</i> spp.																		Water-milfoil
<i>Callitriche</i> spp.																		Water-starwort
<i>Alisma</i> cf. <i>plantago-aquatica</i> L.																		Possible
<i>Potamogeton</i> spp.																		Water-plantain
WATERSIDE TAXA																		Pondweed
<i>Alnus glutinosa</i> (L.) Gaertn.																		Alder
<i>Lycopus europaeus</i> L.																		Gypsywort
<i>Carex</i> spp.																		Sedge
WET/DAMP GROUND TAXA																		
<i>Filipendula ulmaria</i> (L.) Maxim.																		Meadowsweet
<i>Juncus</i> sp.																		Rush
WOODLAND TAXA																		
<i>Urtica dioica</i> L.																		Common Nettle
<i>Betula pendula</i> Roth																		Silver Birch
<i>Betula pubescens</i> Ehr.																		Downy Birch
<i>Oxalis acetosella</i> L.																		Wood-sorrel
TAXA OF VARIED ENVIRONMENTS/ SOILS																		
<i>Galeopsis</i> spp.																		Hemp-nettle
<i>Stellaria media</i> s.l.																		Common Chickweed
cf. <i>Sanguisorba minor</i> Scop. ssp. minor																		Possible Salad Burnet

automatically result in a relative increase in the proportion of herb/heath pollen, even if the actual number of pollen grains was constant.

There is also a suggestion that some areas of heath or bog may have been locally present. Small traces of heather (*Calluna vulgaris* (L.) Hull) pollen were found along with two individuals of the weevil *Micrelus ericae*, which feeds on heather. Small numbers of the spores of horse tail (*Equisetum* spp.), polypody fern (*Polypodium* spp.) and bracken (*Pteridium aquilinum* agg.) were also recorded on the pollen slides. A single individual of the scarabid beetle *Aegialia sabuleti* was also recovered. This species is normally associated with decaying matter on sandy ground (Jessop 1986).

Wetlands and local bodies of water

There is clear evidence in all of the environmental proxies that this wood carr peat developed in an area which contained pools of still or stagnant water surrounded by waterside vegetation, damp ground and carr woodland. The insect fauna was dominated by aquatic and waterside species (see groups 'a' and 'ws' in Table 2 and Fig. 4). The vast majority of the species recovered are 'diving water beetles' (Dytiscidae) such as *Agabus bipustulatus*, *Ilybius ater*, *I. fuliginosus* and *Coelambus impressopunctatus*. These species are all associated with small bodies of slow-flowing or still water, often with dense stands of waterside vegetation (Nilsson and Holmen 1995). *Agabus sturmi* and the hydraenid *Hydraena britteni* usually favour similar bodies of water, particularly if decaying tree leaves are present (Hansen 1987; Nilsson and Holmen 1995). Other water beetles present such as the *Ochthebius*, *Limnebius*, and *Laccobius* species, and *Coelostoma orbiculare*, also indicate similar conditions (Hansen 1987). Damp, muddy and possibly also waterside conditions

are also suggested by several of the species of plant macrofossils recovered, including gypsywort (*Lycopus europaeus* L.), rush (*Juncus* spp.) and sedge (*Carex* spp.). Evidence for slow-flowing, deep water is strongly supported by abundant macrofossil remains of water starwort (*Callitriche* sp.) and water plantain (*Alisma* cf. *plantago-aquatica* L.), which account for 40%–56% of all seed identifications in the samples (Table 5). It is probable, therefore, that this deposit represents alder carr developing around a pool.

Large numbers of the 'reed beetle' *Donacia vulgaris* suggest that either sedges or bur-reeds surrounded the open areas of water. The presence of these species of plant is also recorded in both the pollen and plant macrofossils assemblages. Open areas of water also seem to have supported pondweed (*Potamogeton* spp.) and duckweed (*Lemna* sp.), since these plants are the hosts of *Donacia versicolorea* and the weevil *Tanysphyrus lemnae* respectively. The striking yellow and blue-striped 'leaf beetle' *Prasocuris phellandrii* suggests that waterside 'cow parsleys' also occurred in the area (Koch 1992). The presence of dense stands of waterside vegetation growing in soft silt and mud is also suggested by the ecology of the range of the Carabidae, 'ground beetles', recovered. These included *Patrobus septentrionis*, *Pterostichus diligens*, *P. nigrita*, *P. anthracinus* and *Agonum fuliginosum* (Lindroth 1974). Similar conditions are often favoured by several of the species of staphylinid 'rove beetles' recorded such as the *Lesteva* species, *Lathrimaeum unicolor* and *Olophrum piceum* (Tottenham 1954). The helodid genus *Cyphon*, present in large numbers, usually is associated with shallow detritus and leaf filled waters in small ponds (Harde 1984).

Table 5 Summary of macroscopic plant remains analysis. Numbers in [] = relative proportion of all identifications

Sample	Water conditions	Surrounding Vegetation
Sample 1 (0–10 cm)	Deep water – water starwort (<i>Callitriche</i> spp.) abundant [38%]	Some evidence for damp to wet ground (<i>Juncus</i> spp./ <i>Carex</i> spp.) [25%]. Only limited evidence for woodland, with a small quantity of indeterminate birch (<i>Betula</i> spp.) seed present [7%]
Sample 3 (20–30 cm)	Deep water – water starwort (<i>Callitriche</i> spp.) and water plantain (<i>Alisma</i> cf. <i>plantago-aquatica</i> L.) present, but not accounting for a large proportion of the overall assemblage [9%]	Strong evidence for damp to wet ground (<i>Lycopus europaeus</i> L., <i>Juncus</i> spp. and <i>Carex</i> spp. seeds abundant) [40%]. Also supported by substantial quantities of alder (<i>Alnus glutinosa</i> (L.) Gaertn.) seeds. Evidence for woodland (most likely carr) with alder and birch seeds abundant [37%]
Sample 5 (40–50 cm)	Deep water – water starwort (<i>Callitriche</i> spp.) and water plantain (<i>Alisma</i> cf. <i>plantago-aquatica</i> L.) abundant [56%]	Evidence for damp ground very limited, with only small quantities of <i>Lycopus europaeus</i> L., <i>Juncus</i> spp. and <i>Carex</i> spp. seeds recovered [2%]. Alder and birch seeds still abundant – suggesting woodland (most likely carr with birch in drier areas) present [29%]

Discussion: the use and change in the landscape at Bury and its comparison to regional patterns

This multi-proxy analysis suggests that, at least on the valley floor, the landscape was dominated by alder carr wetland during the Middle and Late Bronze Age. Though alder carr dominates throughout the whole period of the sedimentary record, there does appear to be a change in the vegetational sequence from one dominated by alder/hazel carr woodland to one including a more substantial proportion of oak and birch. It is possible that this represents a change in how pollen from the valley sides was 'filtered' by the surrounding vegetation and the alder carr during this period (Waller 1994); alternately reductions in local prolific pollen producers (such as alder) may have resulted in increased proportional representation of the dryland taxa (Fyfe 2006). Indeed, it is clear from all the proxy environmental indicators from these three deposits that there are changes to both the surface and marginal waterside vegetation, which suggests that the area may have gone through drier and wetter phases. It is also possible that this change in the pollen spectra simply may represent a 'recolonisation' of the landscape by oak and birch during the Late Bronze Age. This may suggest that the area had been cleared to a limited extent by local populations in the Early or Middle Bronze Age. This is perhaps supported by Barnes' (1993) work on the pollen from Castle Steads itself which showed a decline and subsequent recovery in elm pollen at this time. In addition, in the later or uppermost deposit, there are occasional clearance indicators in the pollen data such as ribwort plantain (*Plantago lanceolata* L.) and self-heal (*Prunella vulgaris* L.), although these herbs never appear to have contributed significantly to the overall vegetation and may merely highlight woodland edge communities or a general opening up of the landscape by means other than human agency. The presence of small clearings in earlier periods is also suggested by the recovery of a small number of dung beetles. It is possible that these changes in vegetation indicate clearings created through human activity such as grazing livestock and farming. However, the role of wild, as well as domesticated, animals in forming and maintaining such small clearances should not be forgotten (Buckland and Edwards 1984; Robinson 2000; Vera 2000).

A single undated pollen sample from the 1992 excavations from the ditch sequence in Castle Steads hillfort suggests that many aspects of the landscape

seen in the Bronze Age at this site continue into later periods (Barnes 1993). Perhaps the main differences are that arboreal pollen had declined to 50% TLP and that there were higher values for open ground indicators, such as grasses and ribwort plantain (*Plantago lanceolata*). Barnes (1993) suggested that this may relate to the opening the area around the plateau of the Hillfort, but that the valley of the Irewell probably remained thickly covered with alder even in this later period. Similarly, Tallis and McGuire (1971) found that material from below the Roman Road at nearby Ainsworth also gave very high values for alder.

Several sites within the immediate Greater Manchester area have produced similar results to those from Bury. At nearby Hyde, pollen evidence from two peat deposits at Brook House Meadow and Godley Hall Brook (Ogle *et al.* 1997) indicated an alder carr landscape existed during the Bronze Age, with evidence for major clearance during the Late Bronze Age and Early Iron Age (c. 810–415 cal. BC). Similarly, to the south-west of Manchester, pollen from the (first) River Bollin site (Garner 2001) suggested that during the Iron Age the area comprised marginal carr by the river, with the surrounding slopes colonised by hazel and ash (Shimwell and Downhill 1998). Other pollen spectra in the region appear to indicate that clearance and farming, unlike the situation at Bury, were of importance from the Early Bronze Age onwards. This is clearly seen at the excavations at Mellor, Stockport (Thompson *et al.* 2005), where Bronze Age round houses and possible animal enclosures predate the large Iron Age settlement. At the second site from the River Bollin (Oversley Farm), the pollen record for the Bronze Age indicated a dominance of open heathland (with hazel/alder/birch scrub), which would have been grazed by stock. It is also believed that crops were cultivated around the site, and this is borne out by the recovery of charred emmer (*Triticum dicoccum* Schübl.) and barley (*Hordeum* sp.) grain, as well as the presence of crop weeds (Garner 2001). In nearby Cheshire, palaeoenvironmental evidence of clearance and regeneration also has been recorded at a number of Bronze Age and Iron Age sites (Wells *et al.* 1993; Hall *et al.* 1995). Just south of Manchester, at Lindow Moss, the clearance and regeneration of Early Bronze Age woodland was noted in the pollen record, with evidence of increased disturbance during the Iron Age (Branch and Scaife 1995). A notable highly wooded phase and increased period of wetness was also recorded at this site

between 770 and 400 cal. BC and is correlated with climatic deterioration (Branch and Scaife 1995; Mullin 2003; Leah *et al.* 1997). Analysis of pollen samples from around Bar Mere Hillfort in Cheshire also suggest woodland clearance, cultivation and regeneration between 2000 and 1500 cal. BC, followed possibly by the selective clearance of oak until cal. 1200 BC (Schoenwetter 1982). On a more regional scale, the pollen diagrams from Deep Clough, Central Rossendale, Lancashire, also suggest that clearance of woodland was relatively late, with a Late Bronze Age/Early Iron Age date suggested (Tallis and McGuire 1971).

It is clear from this brief review of regional environmental data that the Bronze Age landscape of this part of north-west England was a complex mosaic of different environments. Overall, there is a trend for carr woodland to dominate the valley floors before extensive clearance for agriculture during the Iron Age. The environmental record from Burrs broadly agrees with this general model, indicating a large, alder/hazel carr woodland adjacent to the channel, which eventually was cut off and infilled as a slow-flowing or still pool. However, in contrast to other sites in the north-west, no taxa indicative of intense human activities, such as cereal pollen, charred plant remains or charcoal, were recorded, despite the site's location immediately below the Iron Age Castle Steads hillfort. This suggests that the area of the hillfort was not occupied until after the date of this peat formation (e.g. post 920 to 800 cal. BC). However, as outlined above, it seems highly probable that Bronze Age settlements did exist within the Irwell valley between Bury and Ramsbottom. Notably the evidence includes the discovery in 1908 of two urns and a bronze dagger by Bury parish church, 1.8 km south of Burrs (SD 805110), the 1960s excavation of a ring bank cemetery at Whitelaw Hillock, 3 km north of Burrs, and the circular cairn at Bank Lane (SD 805172).

There has been considerably less work on the entomological and plant macrofossil record for the region. Palaeoentomological studies are restricted to those associated with the Lindow (raised mire) bog bodies in Cheshire (Dinnin, pers. comm.; Girling 1986; Skidmore 1986) and limited work at Brooks Farm, North Lancashire (Osborne 1995). The English Heritage Environmental Archaeology Database only records a handful of published reports for Bronze Age waterlogged plant macrofossils in the region (http://ads.ahds.ac.uk/catalogue/specColl/eab_eh_2004; consulted Sept. 2008), with the major-

ity of work carried out on mires under the auspices of the North West Wetlands Survey (Hall *et al.* 1995; Leah *et al.* 1997; Middleton *et al.* 1995; Wells *et al.* 1993); the results from Burrs appear to be the first published data directly from a river channel in the Greater Manchester area.

Nationally, the insect fauna recovered compares well with a number of other carr sites which have been examined. There is an obvious correspondence to faunas from several of the later trackways from the Somerset Levels that ran through similar alder woodlands (Girling 1977; 1979; 1980; 1985). Several deposits from the floodplains of the River Thames and Trent have produced similar faunas indicative of dense stands of alder carr (Dinnin 1997; Robinson 1991; 1993; 2000; Smith *et al.* 2005).

Conclusions

The organic-rich sediments infilling the small palaeo-channel at Burrs Countryside Park, Bury, Greater Manchester, have provided a useful insight into the Middle to Late Bronze Age landscape of this area. Although extensive palaeoenvironmental studies have been undertaken in this part of north-west England, they have tended to focus primarily on the wetland mires which form such notable features of the contemporary landscape. Therefore, this record from a palaeochannel forms a useful comparative dataset, describing vegetation patterns in a more confined area of the valley floor. It demonstrates that this part of the Irwell Valley was densely wooded throughout the Bronze Age, and there are few signs of human activity, such as clearance and farming, even in the Late Bronze Age. This is particularly notable since the Iron Age hillfort at Castle Steads is adjacent to the site. This perhaps suggests that occupation of this promontory did not occur until the Iron Age.

This work has highlighted two methodological issues, which have wider implications. First, using pollen results from three sub-samples in this 50 cm sequence clearly only glosses over possible environmental changes during approximately 400 years, at most. As an anonymous reviewer pointed out, it would have been preferable to examine 10–12 pollen samples over this sequence. However, this work was carried out as part of commercial trial-trenching and without English Heritage or curatorial support, and most clients would be unsympathetic to expend so much resources on what is in essence a non-archaeological problem.

Although only speculative at this stage, we have put forward the possibility that a decline in one element of the pollen taxa (in this case arboreal pollen) could

result in a relative increase in the proportions of other plant communities. This suggests that a multi-proxy approach to environmental reconstruction is essential to ensure that changes in arboreal pollen do not cloud our reconstruction of past environments.

Acknowledgements

The authors would like to thank Gill Campbell (English Heritage) for providing information from the as yet unpublished English Heritage regional reviews for the area. We would also like to thank Harry Kenward (University of York) for providing useful comments on an earlier draft. We are indebted to both anonymous referees for their many useful comments. Ralph Fyfe also looked over a late version of the paper and made many useful suggestions. Fig. 1 was prepared by Louise Buglass.

References

- Barnes, B. 1993. A palaeoenvironmental investigation of submitted samples originating from Castle Steads, Walmersley, Bury, pp. 20–27 in Fletcher, M. (ed.), *Castle Steads, Bury*. Unpublished evaluation report, North West Archaeological Surveys.
- Bennett, K. D. 1994. Annotated catalogue of pollen and pteridophyte spore types of the British Isles. Unpublished report, Department of Plant Sciences, University of Cambridge.
- Beijerinck, W. 1976. *Zadenatlas der Nederlandsche Flora: Ten Behoeve van de Botanische, Palaeontologische, Bodemcultuur en Warenkennis*. Amsterdam: Backhuys and Meesters.
- Branch, N. and Scaife, R. 1995. The stratigraphy and pollen analysis of peat sequences associated with Lindow III bog body, pp. 19–31 in Turner, R. and Scaife, R. (eds.), *Bog Bodies. New Discoveries and New Perspectives*. London: British Museum.
- Buckland, P. C. and Edwards, K. 1984. The longevity of pastoral episodes of clearance activity in pollen diagrams: the role of post-occupation grazing. *Journal of Biogeography* **1**, 243–49.
- Cappers, R. T. J., Bekker, R. M. and Jans, J. E. A. 2006. *Digital Seed Atlas of the Netherlands*. Groningen: Barkhuis Publishing and Groningen University Library.
- Dinnin, M. 1997. Holocene beetle assemblages from the Lower Trent floodplain at Bole Ings, Nottinghamshire, U.K. *Quaternary Proceedings* **5**, 83–104.
- Fletcher, M. 1986. A fortified site at Castle Steads, Walmersley, Bury. *Greater Manchester Archaeological Journal* **2**, 31–34.
- Fletcher, M. 1992. Castle Steads, Bury. Unpublished evaluation report, North West Archaeological Surveys.
- Fyfe, R. M. 2006. GIS and the application of a model of pollen deposition and dispersal: a new approach to testing landscape hypotheses using the POLLANDCAL models. *Journal of Archaeological Science* **33**, 483–93.
- Garner, D. 2001. The Bronze Age of Manchester Airport: Runway 2, pp. 41–56 in Bruck, J. (ed.), *Bronze Age Landscapes Tradition and Transformation*. Oxford: Oxbow Books.
- Girling, M. A. 1977. Fossil insect assemblages from Rowland's track. *Somerset Levels Papers* **3**, 51–60.
- Girling, M. A. 1979. Fossil insects from the Sweet track. *Somerset Levels Papers* **5**, 84–93.
- Girling, M. A. 1980. The fossil insect assemblage from the Baker Site. *Somerset Levels Papers* **6**, 36–42.
- Girling, M. A. 1985. An 'old forest' beetle fauna from a Neolithic and Bronze Age peat deposit at Stileway. *Somerset Levels Papers* **11**, 80–85.
- Girling, M. A. 1986. The insects associated with Lindow Man, pp. 90–91 in Stead, I. M., Bourke, J. B. and Brothwell, D. (eds.), *Lindow Man. The Body in the Bog*. London: British Museum.
- Grimm, E. C. 1990. TILIA and TILIA.GRAPH. PC spreadsheet and graphics software for pollen data. *INQUA Working Group on Data-handling Methods, Newsletter* **4**, 5–7.
- Grimm, E. C. 2004. TGVView 2.0-2.
- Hall, A. R. and Huntley, J. P. 2007. *A Review of the Evidence for Macrofossil Plant Remains from Archaeological Deposits in Northern England* (Research Department Report Series 87/2007). London: English Heritage.
- Hall, D., Wells, C. E. and Huckerby, E. 1995. *The Wetlands of Greater Manchester* (North West Wetlands Survey 2, Lancaster Imprints 3). Lancaster: Lancaster University Archaeology Unit.
- Hansen, M. 1987. *The Hydrophilidae (Coleoptera) of Fennoscandia and Denmark Fauna* (Fauna Entomologica Scandinavica 18). Leiden: Scandinavian Science Press.
- Harde, K. W. 1984. *A Field Guide in Colour to Beetles*. London: Octopus Books.
- Head, K. 2005. Pollen remains from Bury Caravan Park. Unpublished report, Historic Environment and Archaeology Service, Worcestershire County Council.
- Hyman, P. and Parsons, M. S. 1992. *A Review of the Scarce and Threatened Coleoptera of Great Britain* (UK Nature Conservation Volume 3/1). Peterborough: UK Joint Nature Conservation Committee.
- Hyman, P. and Parsons, M. S. 1994. *A Review of the Scarce and Threatened Coleoptera of Great Britain* (UK Nature Conservation Volume 3/2). Peterborough: UK Joint Nature Conservation Committee.
- Jessop, L. 1986. *Coleoptera: Scarabaeidae*. (Handbooks for the Identification of British Insects 5/11). London: Royal Entomological Society of London.
- Kenward, H. K. 2009. *Northern Regional Review of Environmental Archaeology: Invertebrates in Archaeology in the North of England* (Research Department Report Series 12–2009). London: English Heritage.
- Kenward, H. K., Hall, A. R. and Jones, A. K. G. 1980. A tested set of techniques for the extraction of plant and animal macrofossils from waterlogged archaeological deposits. *Science and Archaeology* **22**, 3–15.
- Koch, K. 1992. *Die Käfer Mitteleuropas* (Ökologie Band 3). Krefeld: Goecke and Evers.
- Leah, M. D., Wells, C. E., Appleby, C. and Huckerby, E. 1997. *The Wetlands of Cheshire* (North West Wetlands Survey 4: Lancaster Imprints 5). Lancaster: Lancaster University Archaeology Unit.
- Lindroth, C. H. 1974. *Coleoptera: Carabidae* (Handbooks for the Identification of British Insects 4/2). London: Royal Entomological Society of London.
- Lucht, W. H. 1987. *Die Käfer Mitteleuropas* (Katalog). Krefeld: Goecke and Evers.
- Mackney, D., Hodgson, J. M., Hollis, J. M. and Staines, S. J. 1983. *Soils of England and Wales: Sheet 3 Midland and Western England*. Harpenden: Soil Survey of England and Wales.
- Middleton, R., Wells, C. E. and Huckerby, E. 1995. *The Wetlands of North Lancashire*. (North West Wetlands Survey 3: Lancaster Imprints 4). Lancaster: Lancaster University Archaeology Unit.
- Moore, P. D., Webb, J. A. and Collinson, M. E. 1991. *Pollen analysis* (2nd edn.). Oxford: Blackwell Scientific Publications.
- Mullin, D. 2003. *The Bronze Age Landscape of the Northern English Midlands* (British Archaeological Reports British Series 351). Oxford: British Archaeological Reports.
- Nilsson, A. N. and Holmen, M. 1995. *The Aquatic Adephaga (Coleoptera) of Fennoscandia and Denmark II. Dytiscidae* (Fauna Entomologica Scandinavica 35). Leiden: E. J. Brill.
- Ogle, M. I., Robinson, M. E. and Shimwell, D. W. 1997. *Field Survey and Laboratory Analysis of Peat Deposits at Brook House Meadow, Godley Hall Brook, Hyde*. Manchester: University of Manchester.
- Osborne, P. J. 1995. Arthropod analysis, pp. 157–59 in Middleton, R., Wells, C. E. and Huckerby, E. (eds.), *The Wetlands of North Lancashire* (North Wetlands Survey 3: Lancaster Imprints 4). Lancaster: Lancaster University Archaeological Unit.
- Packham, J. R. 1978. *Oxalis acetosella* L. (Biological Flora of the British Isles List Br. Vasc. Pl. (1958) No. 170, 1). *Journal of Ecology* **66**, 669–93.
- Robinson, M. A. 1981. The use of ecological groupings of Coleoptera for comparing sites, pp. 251–86 in Jones, M. and Dimbleby, G. (eds.), *The Environment of Man: The Iron Age to the Anglo-Saxon Period* (British Archaeological Reports, British Series 87). Oxford: British Archaeological Reports.

- Robinson, M. A. 1983. Arable/pastoral ratios from insects?, pp. 19–47 in Jones, M. (ed.), *Integrating the Subsistence Economy* (British Archaeological Reports, International Series 181). Oxford: British Archaeological Reports.
- Robinson, M. A. 1991. The Neolithic and Late Bronze Age insect assemblages, pp. 277–325 in Needham, S. (ed.), *Excavation and Salvage at Runnymede Bridge, 1978: the Late Bronze Age Waterfront Site*. London: British Museum.
- Robinson, M. 1993. The Iron Age environmental evidence, pp. 101–20 in Allen, T. G. and Robinson, M. A. (eds.), *The Prehistoric Landscape and Iron Age Enclosed Settlement at Mingies Ditch, Hardwick-with-Yelford, Oxon* (Thames Valley Landscapes: The Windrush Valley Volume 2). Oxford: Oxford Archaeological Unit, Oxford.
- Robinson, M. A. 2000. Coleopteran evidence for the elm decline, Neolithic activity in woodland, clearance and the use of the landscape, pp. 27–36 in Fairbairn, A. S. (ed.), *Plants in Neolithic Britain and Beyond* (Neolithic Studies Group Seminar Papers 5). Oxford: Oxbow Books.
- Schoch, W. H., Pawlik, B. and Schweingruber, F. H. 1988. *Botanical Macro-remains: An Atlas for the Determination of Frequently Encountered and Ecologically Important Plant Seeds*. Berne and Stuttgart: Paul Haupt Publishers.
- Schoenwetter, J. 1982. Environmental archaeology of the Peckforton Hills. *Cheshire Archaeological Bulletin* 8, 10–11.
- Shimwell, D. and Downhill, S. B. 1998. The evaluation of the environmental archaeological potential of selected contexts from the Runway Two Development at Manchester Airport. Chester: Gifford and Partners unpublished report.
- Skidmore, P. 1986. The Dipterous remains, p. 92 in Stead, I. M., Bourke, J. B. and Brothwell, D. (eds.), *Lindow Man. The Body in the Bog*. London: British Museum.
- Smith, D., Osborne, P. and Barratt, J. 2000. Beetles and evidence of past environments at Goldcliff, pp. 245–60 in Bell, M., Caseldine, A. and Neumann, H. (eds.), *Prehistoric Intertidal Archaeology in the Welsh Severn Estuary* (CBA Research Report 120) London: Council for British Archaeology.
- Smith, D. N., Roseff, R., Bevan, L., Brown, A. G., Butler, S., Hughes, G. and Monckton, A. 2005. Archaeological and environmental investigations of a Late Glacial and Holocene river valley sequence on the River Soar, at Croft, Leicestershire. *The Holocene* 15, 353–77.
- Smith, D. N. and Smith, W. 2005. An assessment of the insect and plant macrofossil remains from the Palaeochannel deposits at the Burrs Caravan Park, Bury, Greater Manchester. Unpublished report, IAA, University of Birmingham.
- Smith, D. N. and Whitehouse, N. 2005. Not seeing the trees for the woods: a palaeontomological perspective on Holocene woodland composition, pp. 136–61 in Smith, D. N., Brickley, M. B. and Smith, W. (eds.), *Fertile Ground: Papers in Honour of Professor Susan Limbrey* (AEA Symposia No. 22). Oxford: Oxbow Books.
- Stace, C. 1997. *New Flora of the British Isles* (2nd edn.). Cambridge: Cambridge University Press.
- Tallis, J. and McGuire, J. 1971. Central Rossendale — the evolution of an upland vegetation. I. The clearance of woodland. *Journal of Ecology* 60, 721–37.
- Thompson, A., Noble, P. and Reid, D. 2005. Excavation report for the 2004 and 2005 seasons at Mellor, Stockport. Manchester: University of Manchester Archaeological Unit.
- Tottenham, C. E. 1954. *Coleoptera. Staphylinidae, Section (a) Piestinae to Euaesthetinae* (Handbooks for the Identification of British Insects IV, 8a). London: Royal Entomological Society.
- Vera, F. W. M. 2000. *Grazing Ecology and Forest History*. Wallingford: CABI Publishing.
- Waller, M. P. 1994. Paludification and pollen representation: the influence of wetland size on *Tilia* representation in pollen diagrams. *The Holocene* 4, 430–34.
- Wells, C., Huckerby, E. and Hall, D. 1993. Archaeological and palaeoecological survey in Greater Manchester, pp. 29–33 in *North West Wetlands Survey Annual Report*. Lancaster: Lancaster University Archaeology Unit.
- Whitehouse, N. J. and Smith, D. N. 2004. 'Islands' in Holocene forests: implications for forest openness, landscape clearance and 'culture steppe' species. *Environmental Archaeology* 9, 203–12.