

Utilising functional imaging to predict survival in paediatric brain tumours

Grist, James; Withey, Stephanie; Powell, Stephen; Peet, AC

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Grist, J, Withey, S, Powell, S & Peet, AC 2019, 'Utilising functional imaging to predict survival in paediatric brain tumours', Paper presented at The British Neuro-Oncology Society Annual Meeting, London, United Kingdom, 3/07/19 - 5/07/19.

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BNOS abstract – 250 words. No references, images, or tables.

Title: Utilising functional imaging to predict survival in paediatric brain tumours.

Authors: James T. Grist¹, Stephanie Withey^{1,2,3}, Lesley MacPherson⁴, Adam Oates⁴, Stephen Powell¹, Jan Novak^{2,5}, Laurence Abernethy⁶, Barry Pizer⁷, Richard Grundy⁸, Simon Bailey⁹, Dipayan Mitra¹⁰, Theodoros N. Arvanitis^{1,2,11}, Dorothee P. Auer^{12,13}, Shivaram Avula⁶, Andrew C Peet^{1,2}.

1. Institute of Cancer and Genomic Sciences, School of Medical and Dental Sciences, University of Birmingham, Birmingham, UK.
2. Oncology, Birmingham Women's and Children's NHS foundation trust, Birmingham, United Kingdom.
3. RRPPS, University Hospitals Birmingham NHS foundation trust, Birmingham, United Kingdom.
4. Radiology, Birmingham Women's and Children's NHS foundation trust, Birmingham, United Kingdom.
5. Department of Psychology, School of Life and Health sciences, Aston University, Birmingham, United Kingdom.
6. Radiology, Alder Hey Children's NHS foundation trust, Liverpool, United Kingdom.
7. Oncology, Alder Hey Children's NHS foundation trust, Liverpool, United Kingdom.
8. The Children's Brain Tumour Research Centre, University of Nottingham, Nottingham, United Kingdom.
9. Sir James Spence Institute of Child Health, Royal Victoria Infirmary, Newcastle upon Tyne, United Kingdom
10. Neuroradiology, Royal Victoria Infirmary, Newcastle Upon Tyne, United Kingdom.
11. Institute of Digital Healthcare, WMG, University of Warwick, Coventry, United Kingdom.
12. Sir Peter Mansfield Imaging Centre, University of Nottingham Biomedical Research Centre, Nottingham, United Kingdom.
13. NIHR Nottingham Biomedical Research Centre, Nottingham, United Kingdom.

Intro

Brain tumours are a common cause of death in the paediatric population. We have previously shown that MR imaging and spectroscopy can be used to non-invasively differentiate between tumour types. Here, we demonstrate that functional imaging can be highly predictive of survival and grade in a paediatric cohort.

Methods

Perfusion (PWI) and diffusion weighted imaging (DWI) were performed in a multi-site (Birmingham Children's Hospital, Royal Victoria Infirmary, Alder Hey, Nottingham) cohort ([grade, 5-year survival alive:dead number] = [I,15:1],[II, 5:1],[III,2:3],[IV,8:11]). ROIs were drawn on T₂ imaging and functional imaging features (mean, standard deviation, skewness, and kurtosis) were derived. Supervised machine learning was used to predict 5-year survival and tumour grade from features. ANOVA and post-hoc tests were used to assess differences in features between grade and 5-year survival status.

Results

5-year survival was predicted with 89%, 85%, and 87% accuracy with all imaging, perfusion, or diffusion features, respectively.

A significant difference in perfusion was found between surviving and diseased participants (1.71 ± 0.82 vs 2.62 ± 1 mL/100g/min, respectively, $p < 0.05$). A significant difference in ADC ($\text{mm}^2 \text{s}^{-1}$) between tumour grades was found (1 vs 4 (1533 ± 458 vs 857 ± 239), 4 vs 3 (857 ± 239 vs 1197 ± 137), 4 vs 2 (857 ± 239 vs 1440 ± 557), corrected $p < 0.05$).

Conclusion

We have shown that perfusion and diffusion imaging features can be used to non-invasively assess tumour grade and estimate 5-year survival status in a cohort of paediatric brain tumours.