

## The Intriguing Role of Interleukin 13 in the Pathophysiology of Asthma

Marone, Giancarlo; Granata, Francesco Paolo; Pucino, Valentina; Pecoraro, Antonio; Enrico, Heffler; Stefania, Loffredo; Guy W, Scadding; Varricchi, Gilda

DOI:  
[10.3389/fphar.2019.01387](https://doi.org/10.3389/fphar.2019.01387)

License:  
Creative Commons: Attribution (CC BY)

*Document Version*  
Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*  
Marone, G, Granata, FP, Pucino, V, Pecoraro, A, Enrico, H, Stefania, L, Guy W, S & Varricchi, G 2019, 'The Intriguing Role of Interleukin 13 in the Pathophysiology of Asthma', *Frontiers in Pharmacology*, vol. 10, 1387. <https://doi.org/10.3389/fphar.2019.01387>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



OPEN ACCESS

# The Intriguing Role of Interleukin 13 in the Pathophysiology of Asthma

**Edited by:**

George Grant,  
University of Aberdeen,  
United Kingdom

**Reviewed by:**

Koshika Yadava,  
University of Oxford,  
United Kingdom  
Laurent Pierre Nicod,  
University of Lausanne, Switzerland  
Katherina Sewald,  
Fraunhofer Institute for Toxicology  
and Experimental Medicine (FHG),  
Germany

**\*Correspondence:**

Gilda Varricchi  
gildanet@gmail.com

**\*ORCID:**

Giancarlo Marone  
orcid.org/0000-0003-0233-7830  
Francescopaolo Granata  
orcid.org/0000-0002-2793-8889  
Valentina Pucino  
orcid.org/0000-0002-7683-4568  
Enrico Heffler  
orcid.org/0000-0002-0492-5663  
Antonio Pecoraro  
orcid.org/0000-0003-3617-2819  
Stefania Loffredo  
orcid.org/0000-0002-5871-1898  
Guy Scadding  
orcid.org/0000-0002-6458-6771  
Gilda Varricchi  
orcid.org/0000-0002-9285-4657

**Specialty section:**

This article was submitted to  
Inflammation Pharmacology,  
a section of the journal  
Frontiers in Pharmacology

**Received:** 22 July 2019

**Accepted:** 31 October 2019

**Published:** 06 December 2019

**Citation:**

Marone G, Granata F, Pucino V,  
Pecoraro A, Heffler E, Loffredo S,  
Scadding GW and Varricchi G (2019)  
The Intriguing Role of Interleukin 13 in  
the Pathophysiology of Asthma.  
*Front. Pharmacol.* 10:1387.  
doi: 10.3389/fphar.2019.01387

**Giancarlo Marone**<sup>1,2†</sup>, **Francescopaolo Granata**<sup>3,4†</sup>, **Valentina Pucino**<sup>5†</sup>,  
**Antonio Pecoraro**<sup>6†</sup>, **Enrico Heffler**<sup>7,8†</sup>, **Stefania Loffredo**<sup>3,4,9†</sup>, **Guy W. Scadding**<sup>10†</sup>  
and **Gilda Varricchi**<sup>3,4,9\*†</sup>

<sup>1</sup> Department of Public Health, University of Naples Federico II, Naples, Italy, <sup>2</sup> Azienda Ospedaliera Ospedali dei Colli, Monaldi Hospital Pharmacy, Naples, Italy, <sup>3</sup> Department of Translational Medical Sciences and Center for Basic and Clinical Immunology Research (CISI), University of Naples Federico II, Naples, Italy, <sup>4</sup> WAO Center of Excellence, University of Naples Federico II, Naples, Italy, <sup>5</sup> College of Medical and Dental Sciences, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, United Kingdom, <sup>6</sup> Immunodeficiency Centre for Wales, University Hospital of Wales, Cardiff, United Kingdom, <sup>7</sup> Personalized Medicine, Asthma, and Allergy, Humanitas Clinical and Research Center, IRCCS, Milan, Italy, <sup>8</sup> Department of Biomedical Sciences, Humanitas University, Milan, Italy, <sup>9</sup> Institute of Experimental Endocrinology and Oncology "G. Salvatore" (IEOS), National Research Council (CNR), Naples, Italy, <sup>10</sup> Allergy and Clinical Immunology, Imperial College, National Heart and Lung Institute, London, United Kingdom

Approximately 5–10% of asthmatic patients worldwide suffer from severe asthma. Experimental and clinical studies have demonstrated that IL-13 is an important cytokine in chronic airways inflammation. IL-13 is involved in Th2 inflammation and has been identified as a possible therapeutic target in the treatment of asthma. Two different human monoclonal antibodies (mAbs) anti-IL-13 (tralokinumab and lebrikizumab) block binding and signaling of IL-13 to its receptors, IL-13R $\alpha$ 1 and IL-13R $\alpha$ 2. Several randomized, double-blind, placebo-controlled multicenter studies have evaluated the safety and efficacy of tralokinumab and lebrikizumab in the treatment of adult patients with severe asthma, but all have failed to meet their primary endpoints. No serious adverse events related to the treatment with these anti-IL-13 mAbs have been reported in these studies. These negative clinical results contrast with positive findings from blocking IL-13 signaling in experimental models of asthma, raising doubts about the transferrable value of some models. Interestingly, dupilumab, a mAb which blocks both IL-4 and IL-13 signaling reduces exacerbation rates and improves lung function in severe asthmatics. These results suggest that IL-4 and IL-13 share some, but not all functional activities in airway inflammation. Tralokinumab might show efficacy in a highly selected cohort of asthmatics characterized by overexpression of IL-13.

**Keywords:** asthma, biologics, chronic rhinosinusitis, interleukin 4, interleukin 13, nasal polyposis, tralokinumab

## INTRODUCTION

Bronchial asthma is a chronic inflammatory disorder of the airways characterized by reversible airflow obstruction, bronchial hyperreactivity (BHR), mucus overproduction, angiogenesis, and airway remodeling (Detoraki et al., 2010; Holgate et al., 2015). Asthma is a common disorder resulting in substantial morbidity, healthcare expenditure, and death (Lang and Polansky 1994; Lange et al., 1996). Worldwide, up to 300 million people are affected by asthma, making it one of the most common chronic diseases (World Health Organization. Global surveillance, prevention,

and control of chronic respiratory diseases. <http://www.who.int/gard/publications/GARD/2007>) and approximately 250,000 people die from asthma each year (Lozano et al., 2012). In the majority of patients, asthma can be controlled by combinations of inhaled glucocorticoids (ICS), short- or long-acting  $\beta_2$ -adrenergic agonists (LABA), long-acting muscarinic antagonists, and leukotriene receptor antagonists, according to the Global Initiative for Asthma (GINA) guidelines (GINA <http://www.ginasthma.org/pdf>). However, in approximately 15% of patients, the disease is refractory to conventional treatments (Chachi et al., 2013; Chachi et al., 2017) and results in hospital admissions due to severe exacerbations (Sekiya et al., 2016; Kerkhof et al., 2018).

As with many chronic inflammatory diseases, clinicians now realize that the traditional classification of asthma has been an oversimplification (Marone et al., 2005). Different asthma phenotypes, each with distinct pathophysiology, are now being defined as asthma endotypes (Wenzel 2012; Fahy 2015; Hinks et al., 2016; Terl et al., 2017). The heterogeneity of different forms of asthma reflects the involvement of different immune cell populations and inflammatory mediators (Bagnasco et al., 2017; Varricchi et al., 2017; Marone et al., 2019). Asthma can be classified according to two major endotypes: “T2-high” asthma is characterized by increased levels of type 2 inflammation mainly mediated by mast cells, eosinophils, basophils, T-helper 2 cells, group 2 innate lymphoid cells (ILC2s), and immunoglobulin E (IgE)-producing B cells (Fahy, 2015). Patients with T2-high asthma have eosinophilia and other signs of type 2 inflammation including high levels of IL-4 and IL-13 (Fahy, 2015; Kaur and Chupp, 2019). Increased blood and sputum levels of eosinophils, serum IgE, and the fraction of exhaled nitric oxide (FeNO) have been associated with the mechanisms of T2-high asthma (Gandhi et al., 2017; Robinson et al., 2017). Another feature is the high expression of the prostaglandin D<sub>2</sub> receptor chemoattractant receptor-homologous molecule (CRTH<sub>2</sub>) on Th2 lymphocytes (Cosmi et al., 2000; Marone et al., 2019). “T2-low” asthma is less well characterized and may include several different endotypes (Wenzel, 2012; Choy et al., 2015; Fahy, 2015; Pepper et al., 2017). It has been suggested that Th1 and Th17 pathways and neurogenic inflammation may be involved (Ricciardolo et al., 2017; Samitas et al., 2017).

Levels of IgE, blood eosinophils and FeNO can be useful to guide the selection of monoclonal antibodies (mAbs) in the treatment of different forms of severe uncontrolled asthma. For example, several studies have demonstrated the safety and efficacy of mAbs blocking IL-5 (i.e., mepolizumab and reslizumab) for treatment of patients with severe eosinophilic asthma (Bel et al., 2014; Ortega et al., 2014;

Bjermer et al., 2016; Khatri et al., 2019). Benralizumab, a mAb against IL-5R $\alpha$ , expressed on the surface of human eosinophils (Varricchi et al., 2018a), is particularly effective in patients with severe asthma with high blood eosinophils (FitzGerald et al., 2016; FitzGerald et al., 2018). Because IgE and the high affinity receptor for IgE (Fc $\epsilon$ RI) play a central role in atopic asthma (Varricchi et al., 2018d; Borriello et al., 2019; Varricchi et al., 2019), a mAb anti-IgE (omalizumab) is indicated for the treatment of patients (aged  $\geq 12$  years) with moderate-to-severe uncontrolled allergic asthma (Samitas et al., 2015; Hew et al., 2016). Serum concentrations of IgE are used to guide this treatment.

IL-4 and IL-13 were among the first identified cytokines orchestrating Th2 inflammation (Macchia et al., 2015; McCormick and Heller 2015; Bagnasco et al., 2016). IL-4 and IL-13 are potent mediators of type 2 inflammation with both overlapping and distinct functions. Pascolizumab, a mAb selectively blocking IL-4, failed to produce positive effects (Hart et al., 2002). Two mAbs blocking IL-13 (i.e., anrukinzumab, lebrikizumab) have shown marginal effects in the treatment of asthmatic patients (Corren et al., 2011; Gauvreau et al., 2011; Hanania et al., 2015; Bagnasco et al., 2016; Hanania et al., 2016). Another mAb anti-IL-13 (i.e., tralokinumab, AstraZeneca) failed to reduce asthma exacerbation rate in severe uncontrolled asthmatics (Piper et al., 2013; Brightling et al., 2015; Panettieri et al., 2018; Russell et al., 2018). Considering the individual relevance of IL-4 and IL-13 in the pathogenesis of asthma, these results are surprising and intriguing. A simplistic explanation is that individual blockade of IL-4 or IL-13 is insufficient to inhibit the complex orchestration of allergic inflammation and clinical consequences in severe asthma. This hypothesis is indirectly supported by the efficacy of dupilumab, which inhibits both IL-4 and IL-13 signaling mediated by IL-4R $\alpha$ , in patients with severe uncontrolled asthma (Castro et al., 2018; Rabe et al., 2018).

In this review we analyze the biological and immunological effects of IL-13 in the context of experimental models of asthma and in asthmatic patients. Despite promising findings in several experimental models of allergic inflammation, the results of multicenter studies evaluating the efficacy of anti-IL-13 mAbs in patients with asthma were surprisingly negative. Possible explanations of these discrepancies are discussed.

## Biological and Immunological Effects of Interleukin 13


IL-13 is a pleiotropic cytokine originally cloned from activated human T-lymphocytes (Minty et al., 1993). The human *IL13* gene is located on chromosome 5q31-33 in the cluster of genes encoding IL-4, IL-3, IL-5, IL-9, and granulocyte-macrophage colony-stimulating factor (GM-CSF). The gene encoding IL-13 is upstream of the *IL4* gene, leading to the speculation that these genes arose as a duplication event during evolution. However, IL-13 has only 25% homology with IL-4 thus explaining why these cytokines share some, but not all functional properties. IL-13 can be produced by stimulated Th2 cells (de Vries 1998), B lymphocytes (Hajoui et al., 2004), CD8<sup>+</sup> cells (Dakhama et al., 2013), type 2 ILCs (Jia et al., 2016), alveolar macrophages

**Abbreviations:** ACQ, asthma control questionnaire; AHR, airway hypersensitivity; BAL, bronchoalveolar lavage; CRSwNP, chronic sinusitis with nasal polyps; DC, dendritic cell; ERK, extracellular signal-regulated kinase; DPP-4, dipeptidyl peptidase-4; Fc $\epsilon$ RI, high affinity receptor for IgE; FeNO, fraction of exhaled nitric oxide; FEV1, forced expiratory volume in the first second; ILC, innate lymphoid cell; JAK, Janus kinase; LABA, long-acting  $\beta_2$ -agonist; ICS, inhaled glucocorticoid; mAb, monoclonal antibody; OVA, ovalbumin; PEF, peak expiratory flow; PGD2, prostaglandin D2; PRO, patient-reported outcomes; SABA, short-acting  $\beta_2$ -agonist; STAT6, signal transducer and activator of transcription; TSLP, thymic stromal lymphopoietin; TYK2, tyrosine kinase 2; VCAM, vascular cell adhesion molecule.

(Hancock et al., 1998), human mast cells (Fushimi et al., 1998), and basophils (Ochensberger et al., 1996; Redrup et al., 1998; Borriello et al., 2015).

**Figure 1** schematically illustrates the complex receptor system which mediates the signaling of IL-4 and IL-13. The IL-4R $\alpha$  subunit is a component of both the type I and type II receptors. Type I receptors are composed of the IL-4R $\alpha$  subunit complexed with common  $\gamma$  chain ( $\gamma$ C); this receptor binds to IL-4 and is expressed on cells of hematopoietic stem cell origin. The type II receptor complex consists of IL-4R $\alpha$  partnering with IL-13R $\alpha$ 1 and is found on many non-hematopoietic cells, such as bronchial epithelial cells, smooth muscle cells, fibroblasts, and keratinocytes (Akaiwa et al., 2001). IL-4 signals through both the type I and type II receptor complexes whereas IL-13 signals only through the type II complex, because IL-13 binds to IL-13R $\alpha$ 1, whereas IL-4 primarily binds to IL-4R $\alpha$  (McKenzie et al., 1999). In addition, the two cytokines have different functions and signaling. IL-4R $\alpha$ ,  $\gamma$ C, and IL-13R $\alpha$ 1 all contain proline rich

regions that can bind the Janus kinases JAK1, JAK2, JAK3, and TYK2. In hematopoietic cells that express  $\gamma$ C and the associated JAK3, IL-4 binding to type I receptor results in the activation of JAK1, JAK2, and JAK3 (Hershey, 2003; Bhattacharjee et al., 2013). IL-4 and IL-13 binding to type II receptor activate JAK1, JAK2, and TYK2. Activation of JAKs results in phosphorylation of cytoplasmic tyrosines leading to the recruitment of STAT6 to the receptor, followed by its phosphorylation and activation. The activation of STAT6 is the primary signaling event in the response to IL-4 or IL-13 (Cao et al., 2016). In certain experimental conditions STAT1 and STAT3 can also be activated by both IL-4 and IL-13 (Wang et al., 2004; Bhattacharjee et al., 2013; Pham et al., 2019). The cytoplasmic domain of human IL-13R $\alpha$ 1 contains two tyrosine residues, which might serve as docking sites for STAT3 (Hershey, 2003). Phosphorylated STAT6 and STAT3 monomers dimerize and then translocate to the nucleus, bind to specific DNA elements to regulate transcription (Bhattacharjee et al., 2013).




**FIGURE 1** | Schematic representation of the three receptors that bind IL-4, IL-13, or both. Type I receptor is composed of the IL-4R $\alpha$  subunit complexed with common  $\gamma$ C. This receptor, expressed on hematopoietic cells, binds to IL-4. Ligand binding by type I receptor complex leads to activation of Janus family kinases (JAK1, JAK2, and JAK3) and subsequent phosphorylation of signal transducer and activator transcription 6 (STAT6). Type II receptor consists of IL-4R $\alpha$  complexed with IL-13R $\alpha$ 1 and is found in many non-hematopoietic cells (e.g., bronchial epithelial cells, smooth muscle cells, fibroblasts, keratinocytes). Ligand binding type II receptor complex leads to activation of JAK1, JAK2, and tyrosine kinase 2 (TYK2) and subsequent phosphorylation of STAT6 and STAT3. Activation of JAKs leads to the recruitment of STATs to the receptors, followed by STAT phosphorylation and dimerization. Activated STAT dimers translocate to the nucleus, bind specific DNA elements, and initiate activation of downstream genes. IL-4 signals through both type I and type II receptors, whereas IL-13 signals only through type II receptor. IL-13 also binds to a third IL-13R $\alpha$ 2 receptor whose functions are largely unknown. Under certain circumstances, IL-13 signaling through IL-13R $\alpha$ 2 results in phosphorylation of ERK1/2 in a STAT6-independent manner and the formation of the dimeric transcription factor AP-1. Phosphorylated AP-1 translocates to the nucleus and bind to specific DNA elements.



LaPorte and collaborators have examined in detail the molecular and structural basis of the IL-4/IL-13 receptor system (LaPorte et al., 2008). They demonstrated that IL-4 first binds to IL-4R $\alpha$  to form a binary complex which then binds to the  $\gamma$ c to form the functional ternary complex (type I receptor). IL-4 and IL-13 can also bind with high affinity to IL-4R $\alpha$  and IL-13R $\alpha$ 1, respectively (type II receptor). The authors also compared the kinetics and potency of IL-4 and IL-13 signaling. IL-4 induced tyrosine phosphorylation of STAT6 more rapidly and more potently than IL-13. The latter observation was supported by previous experiments on cultured airway smooth muscle cells (Laporte et al., 2001). IL-4 is a central mediator for Th2 cell polarization, initiation of IgE synthesis, and recruitment of eosinophils (Chatila, 2004; Wynn, 2015). Although IL-13 has some redundancy in these effects, this cytokine has additional roles in mediating goblet cell hyperplasia, airway smooth muscle contractility, collagen deposition, and fibrosis (Gour and Wills-Karp 2015).

There is a distinct IL-13R $\alpha$ 2 subunit, to which only IL-13 binds (Chen et al., 2009). Initially this receptor was considered a decoy receptor (Ingram and Kraft, 2012) involved in removing IL-13 by internalization (Wood et al., 2003; Lupardus et al., 2010; Kasaian et al., 2011). Although the IL-13R $\alpha$ 2 lacks canonical JAK-STAT signaling activity (Kawakami et al., 2001), this hypothesis has come into question because several studies have shown that, under certain circumstances, IL-13R $\alpha$ 2 can mediate IL-13 signaling (Fichtner-Feigl et al., 2006; Fujisawa et al., 2009; He et al., 2013). In human airways, it was found that IL-13R $\alpha$ 2 is involved in IL-13 signaling through the transcription factor activator protein-1 (AP-1) to induce the activation of TGF- $\beta$  (Fichtner-Feigl et al., 2006). Recently, it has been demonstrated that IL-13 induced phosphorylation of ERK1/2 and the downstream activation of AP-1-related genes in human nasal epithelial cells (Liu et al., 2018). The authors proposed that engagement of IL-13R $\alpha$ 2 by IL-13 activates



mitogen-activated protein ERK1/2 pathway and downstream AP-1-related gene *C-JUN*.

**Figure 2** shows schematically that IL-13 is produced by several immune cells and has many diverse functions on a wide variety of cell types relevant to the pathogenesis of allergic disorders. IL-13 can be produced by activated ILC2 (Shimokawa et al., 2017; Wallrapp et al., 2018), Th2 cells (Finkelman et al., 2004; Wynn, 2015), mast cells (Burd et al., 1995; Fushimi et al., 1998; Varricchi et al., 2019), macrophages (Hancock et al., 1998), basophils (Gibbs et al., 1996; Ochensberger et al., 1996; Redrup et al., 1998; Patella et al., 2000; Genovese et al., 2003), eosinophils (Schmid-Grendelmeier et al., 2002; Varricchi et al., 2018a), and B cells (Hajoui et al., 2004). In human B cells IL-13 has similar effects as IL-4, including promoting B cell proliferation and inducing class switching to IgE and IgG<sub>4</sub> in combination with CD40/CD40L (Oettgen and Geha, 2001) and inducing expression of the low-affinity IgE receptor CD23 (Punnonen et al., 1993; Gould and Sutton, 2008). In macrophages IL-13 favors the M2 polarization (Martinez-Nunez et al., 2011; Bhattacharjee et al., 2013). IL-13 promotes survival, activation, and recruitment of eosinophils (Luttmann et al., 1996; Horie et al., 1997; Pope et al., 2001). In addition, IL-13 stimulates eosinophil trafficking from the peripheral blood to the site of inflammation by inducing the production of IL-5 and eosinophil chemokines such as eotaxins (Webb et al., 2000; Rosenberg et al., 2007) IL-13 promotes FcεRI expression and proliferation of human mast (Nilsson and Nilsson, 1995; Kaur et al., 2006).

IL-13 also has important effects on non-hematopoietic cells, including endothelial cells, smooth muscle cells, fibroblasts, epithelial cells, and sensory neurons. IL-13 is a potent inducer of vascular cell adhesion molecule-1 (VCAM-1) on endothelial cells, an important aspect in the recruitment of eosinophils (Bochner et al., 1995). In addition, IL-13 increases the expression of β<sub>1</sub> integrin and VCAM-1 on human lung fibroblasts (Doucet et al., 1998) and increases muscular contraction in response to acetylcholine (Laporte et al., 2001; Grunstein et al., 2002). Furthermore, IL-13 enhances proliferation and cholinergic-induced contraction of smooth muscle cells (Wills-Karp, 2001) and induces collagen synthesis in human fibroblasts contributing to airways remodeling. In epithelial cells, IL-13 is a potent inducer of eotaxin (Li et al., 1999). Moreover, IL-13 induces mucus overproduction and goblet cell metaplasia (Kuperman et al., 2002; Kondo et al., 2006). IL-13 induces vascular endothelial growth factors (VEGFs) (Corne et al., 2000) which are pro-angiogenic factors relevant in bronchial asthma (Detoraki et al., 2009; Detoraki et al., 2010; Varricchi et al., 2018b). Recently, it has been demonstrated that IL-13 and IL-4 directly activate mouse and human sensory neurons which express IL-13Ra1 and IL-4Ra (Oetjen et al., 2017).

## Interleukin 13 in Experimental Models of Asthma

Seminal studies in animal models of allergic asthma demonstrated that selective neutralization of IL-13 reduced airway hypersensitivity (AHR), bronchoalveolar lavage (BAL) eosinophils, and mucus overproduction (Grunig et al., 1998; Wills-Karp et al., 1998). Furthermore, IL-13 delivery to the

airways caused all of these effects (Grunig et al., 1998; Wills-Karp et al., 1998). Overexpression of IL-13 in the lung of mice caused mucus hypersecretion, subepithelial fibrosis, eotaxin production, and eosinophilic infiltration (Zhu et al., 1999). Interestingly, mice with targeted deletion of IL-13 failed to develop allergen-induced AHR, despite the presence of eosinophilic pulmonary infiltration (Walter et al., 2001). Mice lacking STAT6 were protected from pulmonary effects of IL-13 (Kuperman et al., 2002). Importantly, reconstitution of STAT6 in epithelial cells only was sufficient for IL-13-induced AHR and mucus production in the absence of inflammation and fibrosis. Administration of anti-IL-13 in a mouse model of chronic asthma inhibited eosinophil recruitment in the airways, goblet cell hyperplasia, and subepithelial fibrosis, but only marginally inhibited AHR (Kumar et al., 2004). In another study, blockade of IL-13 with sIL-13Ra2-human IgG fusion protein inhibited AHR associated with brief allergen exposure, but did not modify AHR associated with chronic airway remodeling (Leigh et al., 2004).

An IL-13 vaccine prepared by inserting a murine IL-13 peptide into a viral carrier protein, induced sustained and intense anti-IL-13 IgG antibodies (Ma et al., 2007), associated with inhibition of ovalbumin (OVA)-induced acute airway allergic responses. In a more recent study, the same experimental approach suppressed BAL IL-13 concentration and eosinophils, subepithelial collagen deposition, and mucus hyperproduction (Ma et al., 2013). Interestingly, while IL-13 vaccine inhibited AHR development, it did not revert AHR. The latter findings suggest that IL-13 may be crucial in the development, but not in the maintenance of airway hyperresponsiveness.

*Il13ra2* gene silencing or blockade of IL-13Ra2 signaling led to marked downregulation of TGF-β1 production and collagen deposition in a model of lung fibrosis (Fichtner-Feigl et al., 2006). In IL-13Ra2-deficient mice, AHR and airway inflammation (i.e., mucus production and BAL eosinophils) were attenuated compared to wild type mice following house dust challenge (Chen et al., 2013). More recently, it was reported that, in IL-4Ra-deficient mice, IL-13, but not IL-4, was required for development of OVA-mediated AHR and goblet cell hyperplasia (Kirstein et al., 2016). Munitz and collaborators studying *Il13ra1*<sup>-/-</sup> mice found that IL-13Ra1 is critical for baseline IgE production, AHR, mucus production, and eotaxin production. By contrast, Th2 and IgE responses to antigen were IL-13Ra1-independent (Munitz et al., 2008).

In a novel mouse model of non-allergic asthma overexpression of the activator protein-1 (AP-1) subunit Fra2, caused airway inflammation with IL-13 overexpression, BAL eosinophilia, mucus hyperproduction, AHR, and peribronchial collagen deposition (Gungl et al., 2018). Administration of anti-IL-13 antibody markedly decreased STAT6 phosphorylation in the lung, BAL eosinophilia, and goblet cell hyperplasia. However, peribronchial collagen deposition and bronchial smooth muscle width were not affected by anti-IL-13 administration. The interesting results obtained in this experimental model may be more reflective of severe asthma which exhibit poor response to mAb anti-IL-13.

Genetic deletion of the IL-33 receptor in a mouse model of experimental asthma increased TSLP production, which

stimulated the emergence of IL-13<sup>+</sup> ILC2s and lung mast cells leading to airway hyperresponsiveness (Verma et al., 2018).

Collectively, the results derived from different experimental models indicate that IL-13 plays an important role in the development of several aspects of asthma. However, it is becoming evident that IL-13 neutralization is not sufficient to reverse certain aspects of airway inflammation once they are established.

## Interleukin 13 Expression in Asthmatic Patients

Increased concentration of IL-13 have been found in the blood (Alasandagutti et al., 2017), sputum, bronchial mucosa (Berry et al., 2004), and BAL fluid (Prieto et al., 2000) of asthmatic patients compared to healthy individuals. Increased IL-13 expression in asthma was confirmed by IL-13 mRNA overexpression (Truyen et al., 2006) and by *ex vivo* stimulation of sputum T cells (Boniface et al., 2003). Following allergen challenge, IL-13 is increased in BAL (Huang et al., 1995; Kroegel et al., 1996). *Ex vivo* BAL T cells express IL-13 mRNA (Bodey et al., 1999); expression is inversely related to forced expiratory volume in 1 s (FEV<sub>1</sub>) (Barcelo et al., 2006). Several studies have shown that IL-13 is expressed in bronchial biopsies in patients with asthma (Kroegel et al., 1996; Naseer et al., 1997; Berry et al., 2004; Saha et al., 2008). IL-13<sup>+</sup> ILC2 were increased in the circulation of asthmatic patients with levels correlating with asthma severity (Jia et al., 2016). High production of IL-13 by cord blood CD4<sup>+</sup> T cells is a predictor of development of atopic disorders (Martinez 2002).

## Polymorphisms in the Interleukin 13/Interleukin 4 Receptor Complex Associated With Asthma

Polymorphisms have been identified both in the IL-13 promoter (IL-13-1112 T) and the coding region (IL-13+Arg 130Gen, IL-13+2044G > A) in asthmatic patients (Hershey et al., 1997; Graves et al., 2000; Heinzmann et al., 2000; Arima et al., 2002; Cameron et al., 2006). One of the polymorphisms (Arg 130 Gln) (A/G) identified occur in the region critical for receptor-ligand interactions. Multiple polymorphism in IL-4R gene have also been identified and associated with asthma (Hershey et al., 1997; Kruse et al., 1999; Ober et al., 2000). Recently, three meta-analyses of IL-13 polymorphisms in adults and children suggested that the IL-13+1923 C/T polymorphism is associated with increased risk of asthma (Liu et al., 2014; Mei and Qu, 2017; Xu et al., 2017).

## Tralokinumab and Lebrikizumab for the Treatment of Severe Uncontrolled Asthma

Several anti-IL-13 mAbs (anrakinumab, lebrikizumab, tralokinumab) have provided an opportunity to investigate the role of this cytokine in the pathophysiology of severe asthma, as well as assessing treatment response. A phase 1 study evaluated the pharmacokinetics, safety, and tolerability of tralokinumab in asthmatic patients receiving three different i.v. doses: 1 mg/kg, 5 mg/kg, or 10 mg/kg (Singh et al., 2010). Despite the small sample

size, pharmacokinetics were linear over the dose range studied. The half-life was found to be 2–3 weeks and tralokinumab exhibited an acceptable safety profile.

A phase 2a, randomized, double-blind, placebo-controlled, parallel-group, multicenter study investigated the effects of different dose regimens of tralokinumab in 194 adults with moderate-to-severe asthma inadequately controlled with standard therapy (Piper et al., 2013). Three dose regimens were evaluated: 47 patients received s.c. tralokinumab 150 mg, 51 patients received tralokinumab 300 mg, 48 subjects received tralokinumab 600 mg, and 48 patients received placebo. The primary endpoint was the change from baseline in mean Asthma Control Questionnaire score (ACQ-6) at week 13. Secondary endpoints were change in FEV<sub>1</sub>, pre-bronchodilator lung function, patient-reported outcomes (PROs), rescue  $\beta_2$ -agonist use, and safety outcomes. Mean ACQ-6 score improved in all treatment groups, from baseline to week 13. These changes in ACQ-6 persisted through week 24 and were greater in active patients with higher IL-13 sputum concentrations compared with subjects with lower IL-13 sputum concentrations or subjects receiving placebo. Improvement in FEV<sub>1</sub> were higher in patients with peripheral eosinophil counts  $\geq 300$  cells/ml. Pulmonary function improvements were higher in tralokinumab patients with higher sputum IL-13 ( $\geq 10$  pg/ml<sup>-1</sup>) compared to tralokinumab patients with lower sputum IL-13 ( $\leq 10$  pg/ml<sup>-1</sup>) and patients receiving placebo. However, there was no significant difference in asthma exacerbation rate. The authors defined asthma exacerbations as either a progressive increase of asthma symptoms (cough, wheeze, chest tightness, and/or shortness of breath) or a reduction of >20% in peak expiratory flow or FEV<sub>1</sub> from baseline that did not resolve after the initiation of rescue medications and resulted in an administration of systemic glucocorticoids.

The phase 2b clinical trial run by Brightling and colleagues evaluated safety and tolerability profile and the reduction in exacerbation rate and FEV<sub>1</sub> improvement (Brightling et al., 2015). The authors randomized 452 severe asthmatic patients, all with two-to-six asthma exacerbations in the previous year, to receive tralokinumab (300 mg s.c. either every 2 weeks or every 2 weeks for 3 months and then every 4 weeks) or placebo as add-on therapy for 1 year. At the end of the study, they reported no changes in the annual exacerbation rate at week 52 (primary endpoint) in patients treated with tralokinumab, either every 2 or 4 weeks, *versus* placebo or in time to first exacerbation. The authors defined asthma exacerbation as an increase in asthma symptoms resulting in use or increase in dose of systemic glucocorticoids for three or more consecutive days. Similarly, secondary endpoints such as improvements in prebronchodilator FEV<sub>1</sub>, ACQ-6, and AQLQ(S) were not significant in patients treated with tralokinumab compared with placebo. They found a significant improvement in FEV<sub>1</sub> in patients treated with tralokinumab every 2 weeks. In this study measurement of serum dipeptidyl peptidase-4 (DPP-4) and periostin concentrations were included as a predictive candidate biomarkers before the study was unmasked. Subgroup analyses of patients receiving tralokinumab every 2 weeks with airway reversibility at baseline, but not receiving oral glucocorticoids, showed some clinical improvements in the subgroup who had raised serum DPP-4



and periostin (Brightling et al., 2015). These preliminary results suggested that certain subpopulations of patients with severe asthma might respond to tralokinumab treatment.

A phase 2 multicenter, double-blind, randomized, placebo-controlled trial evaluated the effects of tralokinumab on eosinophilic airway inflammation in uncontrolled moderate-to-severe asthma (MESOS) (Russell et al., 2018). In this study, participants aged 18–75 years were randomly assigned to receive tralokinumab (300 mg s.c. every 2 weeks) or placebo. The primary outcome measure was change from baseline to week 12 in bronchial biopsy eosinophil count. Secondary outcome measures included change in blood and sputum eosinophil counts. Exploratory outcomes included FeNO and blood IgE concentrations. Tralokinumab did not affect bronchial, peripheral blood, or eosinophil counts compared to placebo at week 12. FeNO concentrations and total blood IgE were significantly reduced. The authors concluded that IL-13 is not crucial for eosinophilic airway inflammation control in patients with moderate-to-severe asthma.

Two large phase 3 clinical trials, STRATOS 1 and STRATOS 2, explored the use of periostin and DPP-4 as biomarkers of IL-13-driven inflammatory patterns in patients aged 12–75 years with severe uncontrolled asthma treated with tralokinumab (300 mg s.c. every 2 weeks for 52 weeks) or placebo (Panettieri et al., 2018). The primary endpoint was the annualized asthma exacerbation rate reduction at week 52 in the all-comers population for STRATOS 1 and in the biomarker-positive population for STRATOS 2. The results of both trials confirmed that tralokinumab did not improve the annual asthma exacerbation rate in the all-comers population with severe asthma. In contrast to the preliminary results of the phase 2 trial (Brightling et al., 2015), periostin and DPP-4 were not shown to predict response to tralokinumab. In both trials, tralokinumab-treated participants had a small increase in blood eosinophil counts from baseline, whereas placebo-treated participants did not. A recent meta-analysis of six randomized clinical trials suggested that tralokinumab was well tolerated and modestly improved FEV<sub>1</sub> but did not reduce asthma exacerbations in severe uncontrolled asthma (Zhang et al., 2019).

Given the need to reduce oral glucocorticoid administration in patients with severe asthma, treatments that may allow tapering of glucocorticoids without loss of disease control are needed. A 40-week, randomized, double-blind trial (TROPOS) evaluated the oral glucocorticoid-sparing potential of tralokinumab in patients with severe, uncontrolled asthma requiring maintenance glucocorticoid treatment plus ICS/LABA (Busse et al., 2019). One hundred forty patients were randomized to tralokinumab (300 mg s.c. every 2 weeks) or placebo. The primary endpoint was percentage change from baseline in average glucocorticoid dose at week 40, while maintaining asthma control. Secondary endpoints included patients with a prescribed maintenance glucocorticoid dose of  $\leq 5$  mg, those with greater than 50% reduction in prescribed maintenance glucocorticoids dose and annual asthma exacerbation rate. An asthma exacerbation was defined as worsening of asthma that required a temporary increase in systemic glucocorticoids for  $\geq 3$  days or that resulted in an emergency-room or urgent-care visit that led to a temporary increase in systemic glucocorticoids for  $\geq 3$  days to treat symptoms or an inpatient hospitalization due

to asthma. There were no significant between-group differences for primary and secondary endpoints. Reporting of adverse events and serious adverse events were similar for the tralokinumab and placebo groups.

A randomized, double-blind, placebo controlled study examined the effects of another anti-IL-13 mAb, lebrikizumab (250 mg s.c. once monthly for 6 months), on change in prebronchodilator FEV<sub>1</sub> from baseline to week 12 in 219 adults with uncontrolled asthma (Corren et al., 2011). Lebrikizumab treatment was associated with greater improvement in percent change in FEV<sub>1</sub> in patients with high pretreatment levels of serum periostin compared to patients with low periostin levels. In two replicate studies (LUTE and VERSE) in patients with moderate-to-severe uncontrolled asthma, lebrikizumab (37.5, 125 or 250 mg s.c. every 4 weeks) reduced asthma exacerbation rate by 60% compared to placebo in periostin-high patients and by 5% in periostin-low patients (Hanania et al., 2015). In these studies the authors defined asthma exacerbation as new or increased asthma symptoms that led to treatment with systemic glucocorticoids or to hospitalization. Two replicate, phase 3 trials (LAVOLTA1 and LAVOLTA2) explored the use of periostin and eosinophilia ( $\geq 300$  cells/ $\mu$ l) as biomarkers of IL-13-driven inflammatory patterns in patients with severe uncontrolled asthma treated with lebrikizumab (37.5 mg or 125 mg s.c. once every 4 weeks for 52 weeks or placebo) (Hanania et al., 2016). The primary endpoint was the reduction in the rate of asthma exacerbations over 52 weeks in biomarker-high patients (periostin  $\geq 50$  ng/ml or blood eosinophils  $\geq 300$  cells/ $\mu$ l). In contrast to the preliminary results of phase 2 trials (Corren et al., 2011; Hanania et al., 2015), lebrikizumab did not consistently show significant reduction in asthma exacerbations in biomarker-high patients.

## Anti-Interleukin 13 in Nasal Polyposis

Chronic rhinosinusitis with nasal polyps (CRSwNP) is a common and relevant comorbidity for severe asthma (Heffler et al., 2013). The prevalence of CRSwNP is greater in asthmatics compared to the general population (Settipane and Chafee, 1977) and it increases with the severity of asthma (Pearlman et al., 2009; Lin, 2011), with the highest prevalence rates in non-atopic, late-onset, severe asthmatics (Amelink et al., 2013). More than 60% of patients with CRSwNP have asthma (Ragab et al., 2004; Guida et al., 2010). Moreover, CRSwNP is one of the significant determinants of poor asthma control (Heffler et al., 2013) and a hallmark of refractory eosinophilic asthma (Amelink et al., 2013). The clinical evidence of a strong relationship between CRSwNP and severe asthma raises the possibility of a shared pathogenesis. In particular both diseases seem to be dependent on eosinophilic inflammation mediated by epithelial cytokines such as thymic stromal lymphopoietin (TSLP), IL-25, and IL-33 secreted as a consequence of epithelial damage (Boita et al., 2016; Metha 2016; Varricchi et al., 2018c). The latter activation is mediated by ILC2 which produce several type 2 cytokines such as IL-13 and IL-5 (Aron and Akbari 2017; Poposki et al., 2017). Recently, it has been reported that there were significant increases in IL-13, IL-13R $\alpha$ 1, and IL-13R $\alpha$ 2



mRNA and protein concentrations in nasal polyp epithelium (Liu et al., 2018). Moreover, IL-13 treatment resulted in mucus overproduction and impairment of ciliary function of human nasal epithelial cells. Therefore, an anti-IL-13 strategy, such as tralokinumab, may be helpful in the management of patients with nasal polyps. Unfortunately, there are no clinical trials of treatment of CRSwNP with anti-IL13 mAbs. Interestingly, patients treated with dupilumab, a mAb against IL-4R $\alpha$ , the common receptor for both IL-4 and IL-13 had a significant reduction in polyp size and improvement in symptoms and nasal and olfactory function (Bachert et al., 2016).

## CONCLUDING REMARKS AND PERSPECTIVES

Three phase 2 clinical trials (Piper et al., 2013; Brightling et al., 2015; Russell et al., 2018) and three phase 3 clinical trials (Panettieri et al., 2018; Busse et al., 2019) have shown that tralokinumab did not lower the annual exacerbation rate and did not improve ACQ-6 scores compared to placebo in severe uncontrolled asthmatic patients. These negative findings parallel the results of two replicate, phase 3 trials with lebrikizumab (Hanania et al., 2016). These negative results are surprising given the wide spectrum of pro-inflammatory and pro-fibrogenic activities of IL-13 in experimental models of asthma (Chu et al., 1998; Laporte et al., 2001; Komai et al., 2003; Kanoh et al., 2011) and in asthmatic patients (Huang et al., 1995; Kroegel et al., 1996; Boniface et al., 2003; Truyen et al., 2006; Saha et al., 2008; Jia et al., 2016).

There are several possible explanations of these negative findings. First, it is likely that IL-13 is not the main cytokine involved in the complex network of severe asthma pathogenesis, meaning blocking it alone is ineffective. Second, as shown in several trials (Brightling et al., 2015; Hanania et al., 2016; Panettieri et al., 2018), the biomarkers (e.g., periostin, DPP-4, peripheral eosinophil count) used to identify responders to anti-IL-13 therapy are not optimal. Third, in both experimental models (Walter et al., 2001) and clinical studies (Brightling et al., 2015; Hanania et al., 2016; Panettieri et al., 2018) blocking IL-13 appears to have no effect on reducing tissue or blood eosinophilia, the pathophysiological feature most closely linked to asthma exacerbations. Fourth, perhaps the route of administration (i.e., s.c.) and the size of the anti-IL-13 mAb particles are not ideal. For instance, it has been reported the preliminary efficacy of a nebulized inhaled anti-IL-13 mAb antigen-binding fragment in macaque model of asthma (Lightwood et al., 2018). Finally, we cannot exclude the possibility that some of these negative findings could be due to the inclusion in clinical trials of patients with Th2-low asthma.

Interestingly, the initial attempts to develop cytokine therapies for asthma focusing on antagonizing IL-4 were also unsuccessful (Hart et al., 2002; Steinke 2004), as was an attempt to block the combined receptor (Wenzel et al., 2007; Burmeister Getz et al., 2009). By contrast, dupilumab, which binds to IL-4R $\alpha$  and consequently blocks both IL-4 and IL-13 signaling, decreases asthma exacerbations and improves respiratory symptoms in patients with persistent asthma (Castro et al., 2018; Rabe et al., 2018). These observations suggest that only the effective simultaneous blockade of signaling from two

main cytokines (i.e., IL-4 and IL-13) is effective in the treatment of severe asthma. Using allergic preclinical models, it has been demonstrated that the combined blockade of the IL-13 and IL-33 pathways leads to a greater inhibition of type 2 inflammation over inhibition of either pathway alone (Ramirez-Carrozzi et al., 2017). Similarly, co-blockade of IL-13 and IL-25 attenuated AHR, eosinophil infiltration in the lung, and mucus hyperproduction in a mouse model of OVA-induced asthma (Zhang et al., 2017). Recently, a novel dual antagonist anti-TSLP/IL-13 bispecific antibody has been described (Venkataramani et al., 2018). It will be interesting to see whether combinatorial blockade of multiple cytokines, including IL-13, may yield additional efficacy over single-axis therapies alone.

It is intriguing that IL-13 blockade modulates several aspects of different experimental models of allergic asthma (Laporte et al., 2001; Walter et al., 2001; Komai et al., 2003; Munitz et al., 2008; Chen et al., 2013; Ma et al., 2013; Chachi et al., 2017). Moreover, recent results demonstrate that anti-IL-13 antibody improves bronchial hyperresponsiveness and mucus production in a mouse model of non-allergic asthma (Gungl et al., 2018). These positive experimental results contrast with negative results in the treatment of asthmatic patients with different anti-IL-13 mAbs (tralokinumab and lebrikizumab) (Piper et al., 2013; Brightling et al., 2015; Hanania et al., 2016; Panettieri et al., 2018; Russell et al., 2018; Busse et al., 2019). These findings highlight that the results from murine studies do not always predict clinical effectiveness.

In conclusion, despite several efforts, attempts to demonstrate a benefit of anti-IL-13 in patients with severe asthma remain unproven. While this may be because the right predictive biomarkers or patient phenotypes have not yet been identified, it is perhaps more likely that there is enough redundancy in the pathophysiology of severe asthma to persist without IL-13. This is not to say, at this stage, that another clinical indication might not be found for IL-13-blocking drugs in future.

## AUTHOR CONTRIBUTIONS

All authors contributed to reviewing the current literature and writing of the manuscript and approved the final version of the paper. Conceptualization: GM, GS, GV. Original draft preparation: GM, GS, GV. Final editing: GM, FG, VP, AP, EH, SL, GS, GV.

## FUNDING

This work was supported in part by grants from CISI-Lab Project (University of Naples Federico II) and TIMING Project (Regione Campania).

## ACKNOWLEDGMENTS

The authors thank the members of CISI Laboratory-WAO Center of Excellence and our colleagues for their contributions to some of the work reviewed herein and we apologize to the authors who have contributed importantly to this field and whose work was not cited because of space restrictions. The authors thank Dr. Gjada Criscuolo for a critical reading of the manuscript and the medical graphic artist Fabrizio Fiorbianco for the elaboration of figures.

## REFERENCES

- Akaiwa, M., Yu, B., Umeshita-Suyama, R., Terada, N., Suto, H., Koga, T., et al. (2001). Localization of human interleukin 13 receptor in non-haematopoietic cells. *Cytokine* 13, 75–84. doi: 10.1006/cyto.2000.0814
- Alasandagutti, M. L., Ansari, M. S., Sagurthi, S. R., Valluri, V., and Gaddam, S. (2017). Role of IL-13 genetic variants in signalling of asthma. *Inflammation* 40, 566–577. doi: 10.1007/s10753-016-0503-3
- Amelink, M., de Groot, J. C., de Nijs, S. B., Lutter, R., Zwinderman, A. H., Sterk, P. J., et al. (2013). Severe adult-onset asthma: a distinct phenotype. *J. Allergy Clin. Immunol.* 132, 336–341. doi: 10.1016/j.jaci.2013.04.052
- Arima, K., Umeshita-Suyama, R., Sakata, Y., Akaiwa, M., Mao, X. Q., Enomoto, T., et al. (2002). Upregulation of IL-13 concentration in vivo by the IL13 variant associated with bronchial asthma. *J. Allergy Clin. Immunol.* 109, 980–987. doi: 10.1067/mai.2002.124656
- Aron, J. L., and Akbari, O. (2017). Regulatory T cells and type 2 innate lymphoid cell-dependent asthma. *Allergy* 72, 1148–1155. doi: 10.1111/all.13139
- Bachert, C., Mannent, L., Naclerio, R. M., Mullol, J., Ferguson, B. J., Gevaert, P., et al. (2016). Effect of subcutaneous dupilumab on nasal polyp burden in patients with chronic sinusitis and nasal polyposis: a randomized clinical trial. *JAMA* 315, 469–479. doi: 10.1001/jama.2015.19330
- Bagnasco, D., Ferrando, M., Varricchi, G., Passalacqua, G., and Canonica, G. W. (2016). A Critical Evaluation of Anti-IL-13 and Anti-IL-4 Strategies in Severe Asthma. *Int. Arch. Allergy Immunol.* 170, 122–131. doi: 10.1159/000447692
- Bagnasco, D., Ferrando, M., Varricchi, G., Puggioni, F., Passalacqua, G., and Canonica, G. W. (2017). Anti-Interleukin 5 (IL-5) and IL-5Ra biological drugs: efficacy, safety, and future perspectives in severe eosinophilic asthma. *Front. Med. (Lausanne)* 4, 135. doi: 10.3389/fmed.2017.00135
- Barcelo, B., Pons, J., Fuster, A., Sauleda, J., Noguera, A., Ferrer, J. M., et al. (2006). Intracellular cytokine profile of T lymphocytes in patients with chronic obstructive pulmonary disease. *Clin. Exp. Immunol.* 145, 474–479. doi: 10.1111/j.1365-2249.2006.03167.x
- Bel, E. H., Wenzel, S. E., Thompson, P. J., Prazma, C. M., Keene, O. N., Yancey, S. W., et al. (2014). Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N. Engl. J. Med.* 371, 1189–1197. doi: 10.1056/NEJMoa1403291
- Berry, M. A., Parker, D., Neale, N., Woodman, L., Morgan, A., Monk, P., et al. (2004). Sputum and bronchial submucosal IL-13 expression in asthma and eosinophilic bronchitis. *J. Allergy Clin. Immunol.* 114, 1106–1109. doi: 10.1016/j.jaci.2004.08.032
- Bhattacharjee, A., Shukla, M., Yakubenko, V. P., Mulya, A., Kundu, S., and Cathcart, M. K. (2013). IL-4 and IL-13 employ discrete signaling pathways for target gene expression in alternatively activated monocytes/macrophages. *Free Radic. Biol. Med.* 54, 1–16. doi: 10.1016/j.freeradbiomed.2012.10.553
- Bjerner, L., Lemiere, C., Maspero, J., Weiss, S., Zangrilli, J., and Germinar, M. (2016). Reslizumab for inadequately controlled asthma with elevated blood eosinophil levels: a randomized phase 3 study. *Chest* 150, 789–798. doi: 10.1016/j.chest.2016.03.032
- Bochner, B. S., Klunk, D. A., Sterbinsky, S. A., Coffman, R. L., and Schleimer, R. P. (1995). IL-13 selectively induces vascular cell adhesion molecule-1 expression in human endothelial cells. *J. Immunol.* 154, 799–803.
- Bodey, K. J., Semper, A. E., Redington, A. E., Madden, J., Teran, L. M., Holgate, S. T., et al. (1999). Cytokine profiles of BAL T cells and T-cell clones obtained from human asthmatic airways after local allergen challenge. *Allergy* 54, 1083–1093. doi: 10.1034/j.1398-9995.1999.00889.x
- Boita, M., Bucca, C., Riva, G., Heffler, E., and Rolla, G. (2016). Release of Type 2 Cytokines by Epithelial Cells of Nasal Polyps. *J. Immunol. Res.* 2016, 2643297. doi: 10.1155/2016/2643297
- Boniface, S., Koscher, V., Mamessier, E., El Biaze, M., Dupuy, P., Lorec, A. M., et al. (2003). Assessment of T lymphocyte cytokine production in induced sputum from asthmatics: a flow cytometry study. *Clin. Exp. Allergy* 33, 1238–1243. doi: 10.1046/j.1365-2222.2003.01762.x
- Borriello, F., Longo, M., Spinelli, R., Pecoraro, A., Granata, F., Staiano, R. I., et al. (2015). IL-3 synergises with basophil-derived IL-4 and IL-13 to promote the alternative activation of human monocytes. *Eur. J. Immunol.* 45, 2042–2051. doi: 10.1002/eji.201445303
- Borriello, F., Galdiero, M. R., Varricchi, G., Loffredo, S., Spadaro, G., and Marone, G. (2019). Innate Immune Modulation by GM-CSF and IL-3 in Health and Disease. *Int. J. Mol. Sci.* 20, E834. doi: 10.3390/ijms20040834
- Brightling, C. E., Chanez, P., Leigh, R., O'Byrne, P. M., Korn, S., She, D., et al. (2015). Efficacy and safety of tralokinumab in patients with severe uncontrolled asthma: a randomised, double-blind, placebo-controlled, phase 2b trial. *Lancet Respir. Med.* 3, 692–701. doi: 10.1016/S2213-2600(15)00197-6
- Burd, P. R., Thompson, W. C., Max, E. E., and Mills, F. C. (1995). Activated mast cells produce interleukin 13. *J. Exp. Med.* 181, 1373–1380. doi: 10.1084/jem.181.4.1373
- Burmeister Getz, E., Fisher, D. M., and Fuller, R. (2009). Human pharmacokinetics/pharmacodynamics of an interleukin-4 and interleukin-13 dual antagonist in asthma. *J. Clin. Pharmacol.* 49, 1025–1036. doi: 10.1177/0091270009341183
- Busse, W. W., Brusselle, G. G., Korn, S., Kuna, P., Magnan, A., Cohen, D., et al. (2019). Tralokinumab did not demonstrate oral corticosteroid-sparing effects in severe asthma. *Eur. Respir. J.* 53, 1800948. doi: 10.1183/13993003.00948-2018
- Cameron, L., Webster, R. B., Stempel, J. M., Kiesler, P., Kubesch, M., Ramachandran, H., et al. (2006). Th2 cell-selective enhancement of human IL13 transcription by IL13-1112C>T, a polymorphism associated with allergic inflammation. *J. Immunol.* 177, 8633–8642. doi: 10.4049/jimmunol.177.12.8633
- Cao, H., Zhang, J., Liu, H., Wan, L., Zhang, H., Huang, Q., et al. (2016). IL-13/STAT6 signaling plays a critical role in the epithelial-mesenchymal transition of colorectal cancer cells. *Oncotarget* 7, 61183–61198. doi: 10.18632/oncotarget.11282
- Castro, M., Corren, J., Pavord, I. D., Maspero, J., Wenzel, S., Rabe, K. F., et al. (2018). Dupilumab efficacy and safety in moderate-to-severe uncontrolled asthma. *N. Engl. J. Med.* 378, 2486–2496. doi: 10.1056/NEJMoa1804092
- Chachi, L., Shikotra, A., Duffy, S. M., Tliba, O., Brightling, C., Bradding, P., et al. (2013). Functional KCa3.1 channels regulate steroid insensitivity in bronchial smooth muscle cells. *J. Immunol.* 191, 2624–2636. doi: 10.4049/jimmunol.1300104
- Chachi, L., Abbasian, M., Gavrilu, A., Alzahrani, A., Tliba, O., Bradding, P., et al. (2017). Protein phosphatase 5 mediates corticosteroid insensitivity in airway smooth muscle in patients with severe asthma. *Allergy* 72, 126–136. doi: 10.1111/all.13003
- Chatila, T. A. (2004). Interleukin-4 receptor signaling pathways in asthma pathogenesis. *Trends Mol. Med.* 10, 493–499. doi: 10.1016/j.molmed.2004.08.004
- Chen, W., Sivaprasad, U., Tabata, Y., Gibson, A. M., Stier, M. T., Finkelman, F. D., et al. (2009). IL-13R alpha 2 membrane and soluble isoforms differ in humans and mice. *J. Immunol.* 183, 7870–7876. doi: 10.4049/jimmunol.0901028
- Chen, W., Sivaprasad, U., Gibson, A. M., Ericksen, M. B., Cunningham, C. M., Bass, S. A., et al. (2013). IL-13 receptor alpha2 contributes to development of experimental allergic asthma. *J. Allergy Clin. Immunol.* 132, 951–958. doi: 10.1016/j.jaci.2013.04.016
- Choy, D. F., Hart, K. M., Borthwick, L. A., Shikotra, A., Nagarkar, D. R., Siddiqui, S., et al. (2015). TH2 and TH17 inflammatory pathways are reciprocally regulated in asthma. *Sci. Transl. Med.* 7, 301ra129. doi: 10.1126/scitranslmed.aab3142
- Chu, H. W., Halliday, J. L., Martin, R. J., Leung, D. Y., Szefer, S. J., and Wenzel, S. E. (1998). Collagen deposition in large airways may not differentiate severe asthma from milder forms of the disease. *Am. J. Respir. Crit. Care Med.* 158, 1936–1944. doi: 10.1164/ajrccm.158.6.9712073
- Corne, J., Chupp, G., Lee, C. G., Homer, R. J., Zhu, Z., Chen, Q., et al. (2000). IL-13 stimulates vascular endothelial cell growth factor and protects against hyperoxic acute lung injury. *J. Clin. Invest.* 106, 783–791. doi: 10.1172/JCI9674
- Corren, J., Lemanske, R. F., Hanania, N. A., Korenblat, P. E., Parsey, M. V., Arron, J. R., et al. (2011). Lebrikizumab treatment in adults with asthma. *N. Engl. J. Med.* 365, 1088–1098. doi: 10.1056/NEJMoa1106469
- Cosmi, L., Annunziato, F., Galli, M. I. G., Maggi, R. M. E., Nagata, K., and Romagnani, S. (2000). CRTH2 is the most reliable marker for the detection of circulating human type 2 Th and type 2 T cytotoxic cells in health and disease. *Eur. J. Immunol.* 30, 2972–2979. doi: 10.1002/1521-4141(200010)30:10<2972::AID-IMMU2972>3.0.CO;2-
- Dakhama, A., Collins, M. L., Ohnishi, H., Goleva, E., Leung, D. Y., Alam, R., et al. (2013). IL-13-producing BLT1-positive CD8 cells are increased in asthma and are associated with airway obstruction. *Allergy* 68, 666–673. doi: 10.1111/all.12135
- de Vries, J. E. (1998). The role of IL-13 and its receptor in allergy and inflammatory responses. *J. Allergy Clin. Immunol.* 102, 165–169. doi: 10.1016/S0091-6749(98)70080-6
- Detoraki, A., Staiano, R. I., Granata, F., Giannattasio, G., Prevete, N., de Paulis, A., et al. (2009). Vascular endothelial growth factors synthesized by human lung mast cells exert angiogenic effects. *J. Allergy Clin. Immunol.* 123, 1142–1149. doi: 10.1016/j.jaci.2009.01.044

- Detoraki, A., Granata, F., Staibano, S., Rossi, F. W., Marone, G., and Genovese, A. (2010). Angiogenesis and lymphangiogenesis in bronchial asthma. *Allergy* 65, 946–958. doi: 10.1111/j.1398-9995.2010.02372.x
- Doucet, C., Brouty-Boye, D., Pottin-Clemenceau, C., Canonica, G. W., Jasmin, C., and Azzarone, B. (1998). Interleukin (IL) 4 and IL-13 act on human lung fibroblasts. Implication in asthma. *J. Clin. Invest.* 101, 2129–2139. doi: 10.1172/JCI741
- Fahy, J. V. (2015). Type 2 inflammation in asthma—present in most, absent in many. *Nat. Rev. Immunol.* 15, 57–65. doi: 10.1038/nri3786
- Fichtner-Feigl, S., Strober, W., Kawakami, K., Puri, R. K., and Kitani, A. (2006). IL-13 signaling through the IL-13 $\alpha$ 2 receptor is involved in induction of TGF- $\beta$ 1 production and fibrosis. *Nat. Med.* 12, 99–106. doi: 10.1038/nm1332
- Finkelman, F. D., Shea-Donohue, T., Morris, S. C., Gildea, L., Strait, R., Madden, K. B., et al. (2004). Interleukin-4 and interleukin-13-mediated host protection against intestinal nematode parasites. *Immunol. Rev.* 201, 139–155. doi: 10.1111/j.0105-2896.2004.00192.x
- FitzGerald, J. M., Bleeker, E. R., Nair, P., Korn, S., Ohta, K., Lommatzsch, M., et al. (2016). Benralizumab, an anti-interleukin-5 receptor alpha monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet* 388, 2128–2141. doi: 10.1016/S0140-6736(16)31322-8
- FitzGerald, J. M., Bleeker, E. R., Menzies-Gow, A., Zangrilli, J. G., Hirsch, I., Metcalfe, P., et al. (2018). Predictors of enhanced response with benralizumab for patients with severe asthma: pooled analysis of the SIROCCO and CALIMA studies. *Lancet Respir. Med.* 6, 51–64. doi: 10.1016/S2213-2600(17)30344-2
- Fujisawa, T., Joshi, B., Nakajima, A., and Puri, R. K. (2009). A novel role of interleukin-13 receptor  $\alpha$ 2 in pancreatic cancer invasion and metastasis. *Cancer Res.* 69, 8678–8685. doi: 10.1158/0008-5472.CAN-09-2100
- Fushimi, T., Okayama, H., Shimura, S., Saitoh, H., and Shirato, K. (1998). Dexamethasone suppresses gene expression and production of IL-13 by human mast cell line and lung mast cells. *J. Allergy Clin. Immunol.* 102, 134–142. doi: 10.1016/S0091-6749(98)70064-8
- Gandhi, N. A., Pirozzi, G., and Graham, N. M. H. (2017). Commonality of the IL-4/IL-13 pathway in atopic diseases. *Expert Rev. Clin. Immunol.* 13, 425–437. doi: 10.1080/1744666X.2017.1298443
- Gauvreau, G. M., Boulet, L. P., Cockcroft, D. W., Fitzgerald, J. M., Carlsten, C., Davis, B. E., et al. (2011). Effects of interleukin-13 blockade on allergen-induced airway responses in mild atopic asthma. *Am. J. Respir. Crit. Care Med.* 183, 1007–1014. doi: 10.1164/rccm.201008-1210OC
- Genovese, A., Borgia, G., Björck, L., Petraroli, A., de Paulis, A., Piazza, M., et al. (2003). Immunoglobulin superantigen protein L induces IL-4 and IL-13 secretion from human Fc epsilon RI+ cells through interaction with the kappa light chains of IgE. *J. Immunol.* 170, 1854–1861. doi: 10.4049/jimmunol.170.4.1854
- Gibbs, B. F., Haas, H., Falcone, F. H., Albrecht, C., Vollrath, I. B., Noll, T., et al. (1996). Purified human peripheral blood basophils release interleukin-13 and preformed interleukin-4 following immunological activation. *Eur. J. Immunol.* 26, 2493–2498. doi: 10.1002/eji.1830261033
- Gould, H. J., and Sutton, B. J. (2008). IgE in allergy and asthma today. *Nat. Rev. Immunol.* 8, 205–217. doi: 10.1038/nri2273
- Gour, N., and Wills-Karp, M. (2015). IL-4 and IL-13 signaling in allergic airway disease. *Cytokine* 75, 68–78. doi: 10.1016/j.cyto.2015.05.014
- Graves, P. E., Kabesch, M., Halonen, M., Holberg, C. J., Baldini, M., Fritzsche, C., et al. (2000). A cluster of seven tightly linked polymorphisms in the IL-13 gene is associated with total serum IgE levels in three populations of white children. *J. Allergy Clin. Immunol.* 105, 506–513. doi: 10.1067/mai.2000.104940
- Grunig, G., Warnock, M., Wakil, A. E., Venkayya, R., Brombacher, F., Rennick, D. M., et al. (1998). Requirement for IL-13 independently of IL-4 in experimental asthma. *Science* 282, 2261–2263. doi: 10.1126/science.282.5397.2261
- Grunstein, M. M., Hakonarson, H., Leiter, J., Chen, M., Whelan, R., Grunstein, J. S., et al. (2002). IL-13-dependent autocrine signaling mediates altered responsiveness of IgE-sensitized airway smooth muscle. *Am. J. Physiol. Lung Cell Mol. Physiol.* 282, L520–L528. doi: 10.1152/ajplung.00343.2001
- Guida, G., Rolla, G., Badiu, I., Marsico, P., Pizzimenti, S., Bommarito, L., et al. (2010). Determinants of exhaled nitric oxide in chronic rhinosinusitis. *Chest* 137, 658–664. doi: 10.1378/chest.09-0667
- Gungl, A., Biasin, V., Wilhelm, J., Olschewski, A., Kwapiszewska, G., and Marsh, L. M. (2018). Fra2 Overexpression in Mice Leads to Non-allergic Asthma Development in an IL-13 Dependent Manner. *Front. Immunol.* 9, 2018. doi: 10.3389/fimmu.2018.02018
- Hajoui, O., Janani, R., Tulic, M., Joubert, P., Ronis, T., Hamid, Q., et al. (2004). Synthesis of IL-13 by human B lymphocytes: regulation and role in IgE production. *J. Allergy Clin. Immunol.* 114, 657–663. doi: 10.1016/j.jaci.2004.05.034
- Hanania, N. A., Noonan, M., Corren, J., Korenblat, P., Zheng, Y., Fischer, S. K., et al. (2015). Lebrikizumab in moderate-to-severe asthma: pooled data from two randomised placebo-controlled studies. *Thorax* 70, 748–756. doi: 10.1136/thoraxjnl-2014-206719
- Hanania, N. A., Korenblat, P., Chapman, K. R., Bateman, E. D., Kopecky, P., Paggiaro, P., et al. (2016). Efficacy and safety of lebrikizumab in patients with uncontrolled asthma (LAVOLTA I and LAVOLTA II): replicate, phase 3, randomised, double-blind, placebo-controlled trials. *Lancet Respir. Med.* 4, 781–796. doi: 10.1016/S2213-2600(16)30265-X
- Hancock, A., Armstrong, L., Gama, R., and Millar, A. (1998). Production of interleukin 13 by alveolar macrophages from normal and fibrotic lung. *Am. J. Respir. Cell Mol. Biol.* 18, 60–65. doi: 10.1165/ajrcmb.18.1.2627
- Hart, T. K., Blackburn, M. N., Brigham-Burke, M., Dede, K., Al-Mahdi, N., Zia-Amirhosseini, P., et al. (2002). Preclinical efficacy and safety of pascolizumab (SB 240683): a humanized anti-interleukin-4 antibody with therapeutic potential in asthma. *Clin. Exp. Immunol.* 130, 93–100. doi: 10.1046/j.1365-2249.2002.01973.x
- He, C. H., Lee, C. G., Dela Cruz, C. S., Lee, C. M., Zhou, Y., Ahangari, F., et al. (2013). Chitinase 3-like 1 regulates cellular and tissue responses via IL-13 receptor  $\alpha$ 2. *Cell Rep.* 4, 830–841. doi: 10.1016/j.celrep.2013.07.032
- Heffler, E., Pizzimenti, S., Badiu, I., Guida, G., Ricciardolo, F. L., Bucca, C., et al. (2013). Nasal nitric oxide is a marker of poor asthma control. *J. Breath Res.* 7, 026009. doi: 10.1088/1752-7155/7/2/026009
- Heinzmann, A., Mao, X. Q., Akaiwa, M., Kreomer, R. T., Gao, P. S., Ohshima, K., et al. (2000). Genetic variants of IL-13 signalling and human asthma and atopy. *Hum. Mol. Genet.* 9, 549–559. doi: 10.1093/hmg/9.4.549
- Hershey, G. K., Friedrich, M. F., Esswein, L. A., Thomas, M. L., and Chatila, T. A. (1997). The association of atopy with a gain-of-function mutation in the alpha subunit of the interleukin-4 receptor. *N. Engl. J. Med.* 337, 1720–1725. doi: 10.1056/NEJM199712113372403
- Hershey, G. K. (2003). IL-13 receptors and signaling pathways: an evolving web. *J. Allergy Clin. Immunol.* 111, 677–690. doi: 10.1067/mai.2003.1333
- Hew, M., Gillman, A., Sutherland, M., Wark, P., Bowden, J., and Guo, M. (2016). Real-life effectiveness of omalizumab in severe allergic asthma above the recommended dosing range criteria. *Clin. Exp. Allergy* 46, 1407–1415. doi: 10.1111/cea.12774
- Hinks, T. S., Brown, T., Lau, L. C., Rupani, H., Barber, C., Elliott, S., et al. (2016). Multidimensional endotyping in patients with severe asthma reveals inflammatory heterogeneity in matrix metalloproteinases and chitinase 3-like protein 1. *J. Allergy Clin. Immunol.* 138, 61–75. doi: 10.1016/j.jaci.2015.11.020
- Holgate, S. T., Wenzel, S., Postma, D. S., Weiss, S. T., Renz, H., and Sly, P. D. (2015). Asthma. *Nat. Rev. Dis. Primers.* 1, 15025. doi: 10.1038/nrdp.2015.25
- Horie, S., Okubo, Y., Hossain, M., Sato, E., Nomura, H., Koyama, S., et al. (1997). Interleukin-13 but not interleukin-4 prolongs eosinophil survival and induces eosinophil chemotaxis. *Intern. Med.* 36, 179–185. doi: 10.2169/internalmedicine.36.179
- Huang, S. K., Xiao, H. Q., Kleine-Tebbe, J., Paciotti, G., Marsh, D. G., Lichtenstein, L. M., et al. (1995). IL-13 expression at the sites of allergen challenge in patients with asthma. *J. Immunol.* 155, 2688–2694.
- Ingram, J. L., and Kraft, M. (2012). IL-13 in asthma and allergic disease: asthma phenotypes and targeted therapies. *J. Allergy Clin. Immunol.* 130, 829–842. doi: 10.1016/j.jaci.2012.06.034
- Jia, Y., Fang, X., Zhu, X., Bai, C., Zhu, L., Jin, M., et al. (2016). IL-13(+) Type 2 innate lymphoid cells correlate with asthma control status and treatment response. *Am. J. Respir. Cell Mol. Biol.* 55, 675–683. doi: 10.1165/rcmb.2016.00990C
- Kanoh, S., Tanabe, T., and Rubin, B. K. (2011). IL-13-induced MUC5AC production and goblet cell differentiation is steroid resistant in human airway cells. *Clin. Exp. Allergy* 41, 1747–1756. doi: 10.1111/j.1365-2222.2011.03852.x
- Kasaian, M. T., Raible, D., Marquette, K., Cook, T. A., Zhou, S., Tan, X. Y., et al. (2011). IL-13 antibodies influence IL-13 clearance in humans by modulating scavenger activity of IL-13R $\alpha$ 2. *J. Immunol.* 187, 561–569. doi: 10.4049/jimmunol.1100467



- Kaur, R., and Chupp, G. (2019). Phenotypes and endotypes of adult asthma: Moving toward precision medicine. *J. Allergy Clin. Immunol.* 144, 1–12. doi: 10.1016/j.jaci.2019.05.031
- Kaur, D., Hollins, F., Woodman, L., Yang, W., Monk, P., May, R., et al. (2006). Mast cells express IL-13R alpha 1: IL-13 promotes human lung mast cell proliferation and Fc epsilon RI expression. *Allergy* 61, 1047–1053. doi: 10.1111/j.1398-9995.2006.01139.x
- Kawakami, K., Taguchi, J., Murata, T., and Puri, R. K. (2001). The interleukin-13 receptor alpha2 chain: an essential component for binding and internalization but not for interleukin-13-induced signal transduction through the STAT6 pathway. *Blood* 97, 2673–2679. doi: 10.1182/blood.V97.9.2673
- Kerkhof, M., Tran, T. N., Soriano, J. B., Golam, S., Gibson, D., Hillyer, E. V., et al. (2018). Healthcare resource use and costs of severe, uncontrolled eosinophilic asthma in the UK general population. *Thorax* 73, 116–124. doi: 10.1136/thoraxjnl-2017-210531
- Khatri, S., Moore, W., Gibson, P. G., Leigh, R., Bourdin, A., Maspero, J., et al. (2019). Assessment of the long-term safety of mepolizumab and durability of clinical response in patients with severe eosinophilic asthma. *J. Allergy Clin. Immunol.* 143, 1742–1751 e1747. doi: 10.1016/j.jaci.2018.09.033
- Kirstein, F., Nieuwenhuizen, N. E., Jayakumar, J., Horsnell, W. G. C., and Brombacher, F. (2016). Role of IL-4 receptor alpha-positive CD4(+) T cells in chronic airway hyperresponsiveness. *J. Allergy Clin. Immunol.* 137, 1852–1862 e1859. doi: 10.1016/j.jaci.2015.10.036
- Komai, M., Tanaka, H., Masuda, T., Nagao, K., Ishizaki, M., Sawada, M., et al. (2003). Role of Th2 responses in the development of allergen-induced airway remodeling in a murine model of allergic asthma. *Br. J. Pharmacol.* 138, 912–920. doi: 10.1038/sj.bjp.0705105
- Kondo, M., Tamaoki, J., Takeyama, K., Isono, K., Kawatani, K., Izumo, T., et al. (2006). Elimination of IL-13 reverses established goblet cell metaplasia into ciliated epithelia in airway epithelial cell culture. *Allergol. Int.* 55, 329–336. doi: 10.2332/allergolint.55.329
- Kroegel, C., Julius, P., Matthys, H., Virchow, J. C. Jr., and Luttmann, W. (1996). Endobronchial secretion of interleukin-13 following local allergen challenge in atopic asthma: relationship to interleukin-4 and eosinophil counts. *Eur. Respir. J.* 9, 899–904. doi: 10.1183/09031936.96.09050899
- Kruse, S., Japha, T., Tedner, M., Sparholt, S. H., Forster, J., Kuehr, J., et al. (1999). The polymorphisms S503P and Q576R in the interleukin-4 receptor alpha gene are associated with atopy and influence the signal transduction. *Immunology* 96, 365–371. doi: 10.1046/j.1365-2567.1999.00705.x
- Kumar, R. K., Herbert, C., Webb, D. C., Li, L., and Foster, P. S. (2004). Effects of anticytokine therapy in a mouse model of chronic asthma. *Am. J. Respir. Crit. Care Med.* 170, 1043–1048. doi: 10.1164/rccm.200405-681OC
- Kuperman, D. A., Huang, X., Koth, L. L., Chang, G. H., Dolganov, G. M., Zhu, Z., et al. (2002). Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat. Med.* 8, 885–889. doi: 10.1038/nm734
- Lang, D. M., and Polansky, M. (1994). Patterns of asthma mortality in Philadelphia from 1969 to 1991. *N. Engl. J. Med.* 331, 1542–1546. doi: 10.1056/NEJM199412083312302
- Lange, P., Ulrik, C. S., and Vestbo, J. (1996). Mortality in adults with self-reported asthma. Copenhagen City Heart Study Group. *Lancet* 347, 1285–1289. doi: 10.1016/S0140-6736(96)90937-X
- Laporte, J. C., Moore, P. E., Baraldo, S., Jouvin, M. H., Church, T. L., and Schwartzman, I. N. (2001). Direct effects of interleukin-13 on signaling pathways for physiological responses in cultured human airway smooth muscle cells. *Am. J. Respir. Crit. Care Med.* 164, 141–148. doi: 10.1164/ajrccm.164.1.2008060
- LaPorte, S. L., Juo, Z. S., Vaclavikova, J., Colf, L. A., Qi, X., Heller, N. M., et al. (2008). Molecular and structural basis of cytokine receptor pleiotropy in the interleukin-4/13 system. *Cell* 132, 259–272. doi: 10.1016/j.cell.2007.12.030
- Leigh, R., Ellis, R., Wattie, J., Donaldson, D. D., and Inman, M. D. (2004). Is interleukin-13 critical in maintaining airway hyperresponsiveness in allergen-challenged mice? *Am. J. Respir. Crit. Care Med.* 170, 851–856. doi: 10.1164/rccm.200311-1488OC
- Li, L., Xia, Y., Nguyen, A., Lai, Y. H., Feng, L., Mosmann, T. R., et al. (1999). Effects of Th2 cytokines on chemokine expression in the lung: IL-13 potently induces eotaxin expression by airway epithelial cells. *J. Immunol.* 162, 2477–2487.
- Lightwood, D., Tservistas, M., Zehentleitner, M., Sarkar, K., Turner, A., Bracher, M., et al. (2018). Efficacy of an inhaled IL-13 antibody fragment in a model of chronic asthma. *Am. J. Respir. Crit. Care Med.* 198, 610–619. doi: 10.1164/rccm.201712-2382OC
- Lin, D. C. (2011). Association between severity of asthma and degree of chronic rhinosinusitis. *Am. J. Rhinol. Allergy* 205–208. doi: 10.2500/ajra.2011.25.3613
- Liu, Z., Li, P., Wang, J., Fan, Q., Yan, P., Zhang, X., et al. (2014). A meta-analysis of IL-13 polymorphisms and pediatric asthma risk. *Med. Sci. Monit.* 20, 2617–2623. doi: 10.12659/MSM.891017
- Liu, J., Li, Y. Y., Andiappan, A. K., Yan, Y., Tan, K. S., Ong, H. H., et al. (2018). Role of IL-13/Ralpha2 in modulating IL-13-induced MUC5AC and ciliary changes in healthy and CRSwNP mucosa. *Allergy* 73, 1673–1685. doi: 10.1111/all.13424
- Lozano, R., Naghavi, M., Foreman, K., Lim, S., Shibuya, K., Aboyans, V., et al. (2012). Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380, 2095–2128. doi: 10.1016/S0140-6736(12)61728-0
- Lupardus, P. J., Birnbaum, M. E., and Garcia, K. C. (2010). Molecular basis for hared cytokine recognition revealed in the structure of an unusually high affinity complex between IL-13 and IL-13Ralpha2. *Structure* 18, 332–342. doi: 10.1016/j.str.2010.01.003
- Luttmann, W., Knoechel, B., Foerster, M., Matthys, H., Virchow, J. C. Jr., and Kroegel, C. (1996). Activation of human eosinophils by IL-13. Induction of CD69 surface antigen, its relationship to messenger RNA expression, and promotion of cellular viability. *J. Immunol.* 157, 1678–1683.
- Ma, Y., HayGlass, K. T., Becker, A. B., Fan, Y., Yang, X., Basu, S., et al. (2007). Novel recombinant interleukin-13 peptide-based vaccine reduces airway allergic inflammatory responses in mice. *Am. J. Respir. Crit. Care Med.* 176, 439–445. doi: 10.1164/rccm.200610-1405OC
- Ma, Y., Halayko, A. J., Basu, S., Guan, Q., Weiss, C. R., Ma, A. G., et al. (2013). Sustained suppression of IL-13 by a vaccine attenuates airway inflammation and remodeling in mice. *Am. J. Respir. Cell Mol. Biol.* 48, 540–549. doi: 10.1165/rcmb.2012-0060OC
- Macchia, D., Melioli, G., Pravettoni, V., Nucera, E., Piantanida, M., Aminati, M., et al. (2015). Guidelines for the use and interpretation of diagnostic methods in adult food allergy. *Clin. Mol. Allergy* 13, 27. doi: 10.1186/s12948-15-0033-9
- Marone, G., Triggiani, M., and de Paulis, A. (2005). Mast cells and basophils: friends as well as foes in bronchial asthma? *Trends Immunol.* 26, 25–31. doi: 10.1016/j.it.2004.10.010
- Marone, G., Galdiero, M. R., Pecoraro, A., Pucino, V., Criscuolo, G., Triassi, M., et al. (2019). Prostaglandin D2 receptor antagonists in allergic disorders: safety, efficacy, and future perspectives. *Expert Opin. Invest. Drugs* 28, 73–84. doi: 10.1080/13543784.2019.1555237
- Martinez, F. D. (2002). What have we learned from the Tucson Children's respiratory Study? *Paediatr. Respir. Rev.* 3, 193–197. doi: 10.1016/S1526-542(02)00188-4
- Martinez-Nunez, R. T., Louafi, F., and Sanchez-Elsner, T. (2011). The interleukin 13 (IL-13) pathway in human macrophages is modulated by microRNA-155 via direct targeting of interleukin 13 receptor alpha1 IL13Ralpha1. *J. Biol. Chem.* 286, 1786–1794. doi: 10.1074/jbc.M110.169367
- McCormick, S. M., and Heller, N. M. (2015). Commentary: IL-4 and IL-13 receptors and signaling. *Cytokine* 75, 38–50. doi: 10.1016/j.cyt.2015.05.023
- McKenzie, G. J., Fallon, P. G., Emson, C. L., Grecnis, R. K., and McKenzie, A. N. (1999). Simultaneous disruption of interleukin (IL)-4 and IL-13 defines individual roles in T helper cell type 2-mediated responses. *J. Exp. Med.* 189, 1565–1572. doi: 10.1084/jem.189.10.1565
- Mei, Q., and Qu, J. (2017). Interleukin-13 +2044 G/A and +1923C/T polymorphisms are associated with asthma susceptibility in Asians: a meta-analysis. *Med. (Baltimore)* 96, e9203. doi: 10.1097/MD.00000000000009203
- Metha, A. K. (2016). Rhinovirus infection interferes with induction of tolerance to aeroantigens through OX40 ligand, thymic stromal lymphopoietin, and IL-33. *J. Allergy Clin. Immunol.* 137, 278–288. doi: 10.1016/j.jaci.2015.05.007
- Minty, A., Chalou, P., Derocq, J. M., Dumont, X., Guillemot, J. C., Kaghad, M., et al. (1993). Interleukin-13 is a new human lymphokine regulating inflammatory and immune responses. *Nature* 362, 248–250. doi: 10.1038/362248a0
- Munitz, A., Brandt, E. B., Mingler, M., Finkelman, F. D., and Rothenberg, M. E. (2008). Distinct roles for IL-13 and IL-4 via IL-13 receptor alpha1 and the type II IL-4 receptor in asthma pathogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 105, 7240–7245. doi: 10.1073/pnas.0802465105
- Naseer, T., Minshall, E. M., Leung, D. Y., Laberge, S., Ernst, P., Martin, R. J., et al. (1997). Expression of IL-12 and IL-13 mRNA in asthma and their modulation in response to steroid therapy. *Am. J. Respir. Crit. Care Med.* 155, 845–851. doi: 10.1164/ajrccm.155.3.9117015



- Nilsson, G., and Nilsson, K. (1995). Effects of interleukin (IL)-13 on immediate response gene expression, phenotype and differentiation of human mast cells. Comparison with IL-4. *Eur. J. Immunol.* 25, 870–873. doi: 10.1002/eji.1830250337
- Ober, C., Leavitt, S. A., Tsalenko, A., Howard, T. D., Hoki, D. M., Daniel, R., et al. (2000). Variation in the interleukin 4-receptor alpha gene confers susceptibility to asthma and atopy in ethnically diverse populations. *Am. J. Hum. Genet.* 66, 517–526. doi: 10.1086/302781
- Ochensberger, B., Daepf, G. C., Rihs, S., and Dahinden, C. A. (1996). Human lood basophils produce interleukin-13 in response to IgE-receptor-dependent and independent activation. *Blood* 88, 3028–3037. doi: 10.1182/blood.V88.8.3028.bloodjournal8883028
- Oetjen, L. K., Mack, M. R., Feng, J., Whelan, T. M., Niu, H., Guo, C. J., et al. (2017). Sensory neurons co-opt classical immune signaling pathways to mediate chronic itch. *Cell* 171, 217–228 e213. doi: 10.1016/j.cell.2017.08.006
- Oettgen, H. C., and Geha, R. S. (2001). IgE regulation and roles in asthma pathogenesis. *J. Allergy Clin. Immunol.* 107, 429–440. doi: 10.1067/mai.2001.113759
- Ortega, H. G., Liu, M. C., Pavord, I. D., Brusselle, G. G., FitzGerald, J. M., Chetta, A., et al. (2014). Mepolizumab treatment in patients with severe eosinophilic asthma. *N. Engl. J. Med.* 371, 1198–1207. doi: 10.1056/NEJMoa1403290
- Panettieri, R. A. Jr., Sjobring, U., Peterffy, A., Wessman, P., Bowen, K., Piper, E., et al. (2018). Tralokinumab for severe, uncontrolled asthma (STRATOS 1 and STRATOS 2): two randomised, double-blind, placebo-controlled, phase 3 clinical trials. *Lancet Respir. Med.* 6, 511–525. doi: 10.1016/S2213-2600(18)30184-X
- Patella, V., Florio, G., Petraroli, A., and Marone, G. (2000). HIV-1 gp120 induces L-4 and IL-13 release from human Fc epsilon RI+ cells through interaction with the VH3 region of IgE. *J. Immunol.* 164, 589–595. doi: 10.4049/jimmunol.164.2.589
- Pearlman, A. N., Chandra, R. K., Chang, D., Conley, D. B., Tripathi-Peters, A., Grammer, L., et al. (2009). Relationships between severity of chronic rhinosinusitis and nasal polyposis, asthma, and atopy. *Am. J. Rhinol. Allergy* 23, 45–148. doi: 10.2500/ajra.2009.23.3284
- Pepper, A. N., Renz, H., Casale, T. B., and Garn, H. (2017). Biologic Therapy and Novel Molecular Targets of Severe Asthma. *J. Allergy Clin. Immunol. Pract.* 5, 909–916. doi: 10.1016/j.jaip.2017.04.038
- Pham, T. H., Bak, Y., Oh, J. W., Hong, J., Lee, S., Hong, J. T., et al. (2019). Inhibition of IL-13 and IL-13Ralpha2 Expression by IL-32theta in Human Monocytic Cells Requires PKCdelta and STAT3 Association. *Int. J. Mol. Sci.* 20, E1949. doi: 10.3390/ijms20081949
- Piper, E., Brightling, C., Niven, R., Oh, C., Faggioni, R., Poon, K., et al. (2013). A phase II placebo-controlled study of tralokinumab in moderate-to-severe asthma. *Respir. J.* 41, 330–338. doi: 10.1183/09031936.00223411
- Pope, S. M., Brandt, E. B., Mishra, A., Hogan, S. P., Zimmermann, N., Matthaei, K. I., et al. (2001). IL-13 induces eosinophil recruitment into the lung by an L-5- and eotaxin-dependent mechanism. *J. Allergy Clin. Immunol.* 108, 594–601. doi: 10.1067/mai.2001.118600
- Poposki, J. A., Klingler, A. L., Tan, B. K., Soroosh, P., Banie, H., Lewis, G., et al. (2017). Group 2 innate lymphoid cells are elevated and activated in chronic rhinosinusitis with nasal polyps. *Immun. Inflammation Dis.* 5, 233–243. doi: 10.1002/iid3.161
- Prieto, J., Lensmar, C., Roquet, A., van der Ploeg, I., Gigliotti, D., Eklund, A., et al. (2000). Increased interleukin-13 mRNA expression in bronchoalveolar lavage cells of atopic patients with mild asthma after repeated low-dose allergen provocations. *Respir. Med.* 94, 806–814. doi: 10.1053/rmed.2000.0826
- Punnonen, J., Aversa, G., Cocks, B. G., McKenzie, A. N., Menon, S., Zurawski, G., et al. (1993). Interleukin 13 induces interleukin 4-independent IgG4 and IgE synthesis and CD23 expression by human B cells. *Proc. Natl. Acad. Sci. U. S. A.* 90, 730–734. doi: 10.1073/pnas.90.8.3730
- Rabe, K. F., Nair, P., Brusselle, G., Maspero, J. F., Castro, M., Sher, L., et al. (2018). Efficacy and safety of dupilumab in glucocorticoid-dependent severe asthma. *N. Engl. J. Med.* 378, 2475–2485. doi: 10.1056/NEJMoa1804093
- Ragab, A., Clement, P., and Vincken, W. (2004). Objective assessment of lower airway involvement in chronic rhinosinusitis. *Am. J. Rhinol.* 18, 15–21. doi: 10.1177/194589240401800105
- Ramirez-Carrozzi, V., Sambandam, A., Zhou, M., Yan, D., Kang, J., Wu, X., et al. (2017). Combined blockade of the IL-13 and IL-33 pathways leads to a greater inhibition of type 2 inflammation over inhibition of either pathway alone. *J. Allergy Clin. Immunol.* 139, 705–708 e706. doi: 10.1016/j.jaci.2016.08.026
- Redrup, A. C., Howard, B. P., MacGlashan, D. W. Jr., Kagey-Sobotka, A., Lichtenstein, L. M., and Schroeder, J. T. (1998). Differential regulation of IL-4 and IL-13 secretion by human basophils: their relationship to histamine release in mixed eukocyte cultures. *J. Immunol.* 160, 1957–1964.
- Ricciardolo, F. L. M., Sorbello, V., Folino, A., Gallo, F., Massaglia, G. M., avata, G., et al. (2017). Identification of IL-17F/frequent exacerbator endotype in asthma. *J. Allergy Clin. Immunol.* 140, 395–406. doi: 10.1016/j.jaci.2016.10.034
- Robinson, D., Humbert, M., Buhl, R., Cruz, A. A., Inoue, H., Korom, S., et al. (2017). Revisiting Type 2-high and Type 2-low airway inflammation in asthma: current knowledge and therapeutic implications. *Clin. Exp. Allergy* 47, 161–175. doi: 10.1111/cea.12880
- Rosenberg, H. F., Phipps, S., and Foster, P. S. (2007). Eosinophil trafficking in allergy and asthma. *J. Allergy Clin. Immunol.* 119, 1303–1310. doi: 10.1016/j.jaci.2007.03.048
- Russell, R. J., Chachi, L., FitzGerald, J. M., Backer, V., Olivenstein, R., Itlestad, I. L., et al. (2018). Effect of tralokinumab, an interleukin-13 neutralising monoclonal antibody, on eosinophilic airway inflammation in uncontrolled moderate-to-severe asthma (MESOS): a multicentre, double-blind, randomised, placebo-controlled phase 2 trial. *Lancet Respir. Med.* 6, 499–510. doi: 10.1016/S2213-2600(18)30201-7
- Saha, S. K., Berry, M. A., Parker, D., Siddiqui, S., Morgan, A., May, R., et al. (2008). Increased sputum and bronchial biopsy IL-13 expression in severe asthma. *Allergy Clin. Immunol.* 121, 685–691. doi: 10.1016/j.jaci.2008.01.005
- Samitas, K., Delimpoura, V., Zervas, E., and Gaga, M. (2015). Anti-IgE treatment, airway inflammation and remodelling in severe allergic asthma: current knowledge and future perspectives. *Eur. Respir. Rev.* 24, 594–601. doi: 10.1183/16000617.00001715
- Samitas, K., Zervas, E., and Gaga, M. (2017). T2-low asthma: current approach to diagnosis and therapy. *Curr. Opin. Pulm. Med.* 23, 48–55. doi: 10.1097/MCP.0000000000000342
- Schmid-Grendelmeier, P., Altnauer, F., Fischer, B., Bizer, C., Straumann, A., Menz, G., et al. (2002). Eosinophils express functional IL-13 in eosinophilic inflammatory diseases. *J. Immunol.* 169, 1021–1027. doi: 10.4049/jimmunol.169.2.1021
- Sekiya, K., Nakatani, E., Fukutomi, Y., Kaneda, H., Iikura, M., Yoshida, M., et al. (2016). Severe or life-threatening asthma exacerbation: patient heterogeneity identified by cluster analysis. *Clin. Exp. Allergy* 46, 1043–1055. doi: 10.1111/cea.12738
- Settipane, G. A., and Chafee, F. H. (1977). Nasal polyps in asthma and rhinitis. A review of 6,037 patients. *J. Allergy Clin. Immunol.* 59, 17–21. doi: 10.1016/0091-749(77)90171-3
- Shimokawa, C., Kanaya, T., Hachisuka, M., Ishiwata, K., Hisaeda, H., Kurashima, Y., et al. (2017). Mast cells are crucial for induction of group 2 innate lymphoid cells and clearance of helminth infections. *Immunity* 46, 863–874. doi: 10.1016/j.immuni.2017.04.017
- Singh, D., Kane, B., Molfino, N. A., Faggioni, R., Roskos, L., and Woodcock, A. (2010). A phase I study evaluating the pharmacokinetics, safety and tolerability of repeat dosing with a human IL-13 antibody (CAT-354) in subjects with asthma. *MC Pulm. Med.* 10, 3. doi: 10.1186/1471-2466-10-3
- Steinke, J. W. (2004). Anti-interleukin-4 therapy. *Immunol. Allergy Clin. North Am.* 24, 599–614. doi: 10.1016/j.iac.2004.06.008
- Terl, M., Sedlak, V., Cap, P., Dvorakova, R., Kasak, V., Koci, T., et al. (2017). Asthma management: a new phenotype-based approach using presence of eosinophilia and allergy. *Allergy* 72, 1279–1287. doi: 10.1111/all.13165
- Truyen, E., Coteur, L., Dilissen, E., Overbergh, L., Dupont, L. J., Ceuppens, J. L., et al. (2006). Evaluation of airway inflammation by quantitative Th1/Th2 cytokine mRNA measurement in sputum of asthma patients. *Thorax* 61, 202–208. doi: 10.1136/thx.2005.052399
- Varricchi, G., Senna, G., Loffredo, S., Bagnasco, D., Ferrando, M., and Anonica, G. W. (2017). Reslizumab and Eosinophilic Asthma: One Step Closer to Precision Medicine? *Front. Immunol.* 8, 242. doi: 10.3389/fimmu.2017.00242
- Varricchi, G., Galdiero, M. R., Loffredo, S., Lucarini, V., Marone, G., and Mattei, F. (2018a). Eosinophils: the unsung heroes in cancer? *Oncoimmunology* 7, 1393134. doi: 10.1080/2162402X.2017.1393134
- Varricchi, G., Loffredo, S., Galdiero, M. R., Marone, G., Cristinziano, L., and Granata, F. (2018b). Innate effector cells in angiogenesis and lymphangiogenesis. *Curr. Opin. Immunol.* 53, 152–160. doi: 10.1016/j.coi.2018.05.002
- Varricchi, G., Pecoraro, A., Marone, G., Criscuolo, G., Spadaro, G., and enovese, A. (2018c). Thymic stromal lymphopoietin isoforms, inflammatory disorders, and cancer. *Front. Immunol.* 9, 1595. doi: 10.3389/fimmu.2018.01595

- Varricchi, G., Raap, U., Rivellese, F., Marone, G., and Gibbs, B. F. (2018d). Human mast cells and basophils—How are they similar how are they different? *Immunol. Rev.* 282, 8–34. doi: 10.1111/imr.12627
- Varricchi, G., Rossi, F. W., Galdiero, M. R., Granata, F., Criscuolo, G., Spadaro, G., et al. (2019). Physiological roles of mast cells: collegium internationale allergologicum update 2019. *Int. Arch. Allergy Immunol.* 179, 247–261. doi: 10.1159/000500088
- Venkataramani, S., Low, S., Weigle, B., Dutcher, D., Jerath, K., Menzenski, M., et al. (2018). Design and characterization of Zweimab and Doppelmab, high finity dual antagonistic anti-TSLP/IL13 bispecific antibodies. *Biochem. Biophys. Res. Commun.* 504, 19–24. doi: 10.1016/j.bbrc.2018.08.064
- Verma, M., Liu, S., Michalec, L., Sripada, A., Gorska, M. M., and Alam, R. (2018). Experimental asthma persists in IL-33 receptor knockout mice because of the emergence of thymic stromal lymphopoietin-driven IL-9(+) and IL-13(+) type 2 innate lymphoid cell subpopulations. *J. Allergy Clin. Immunol.* 142, 793–803 e798. doi: 10.1016/j.jaci.2017.10.020
- Wallrapp, A., Riesenfeld, S. J., Burkett, P. R., and Kuchroo, V. K. (2018). Type 2 innate lymphoid cells in the induction and resolution of tissue inflammation. *Immunol. Rev.* 286, 53–73. doi: 10.1111/imr.12702
- Walter, D. M., McIntire, J. J., Berry, G., McKenzie, A. N., Donaldson, D. D., DeKruyff, R. H., et al. (2001). Critical role for IL-13 in the development of allergen-induced airway hyperreactivity. *J. Immunol.* 167, 4668–4675. doi: 10.4049/jimmunol.167.8.4668
- Wang, I. M., Lin, H., Goldman, S. J., and Kobayashi, M. (2004). STAT-1 is activated by IL-4 and IL-13 in multiple cell types. *Mol. Immunol.* 41, 873–884. doi: 10.1016/j.molimm.2004.04.027
- Webb, D. C., McKenzie, A. N., Koskinen, A. M., Yang, M., Mattes, J., and Foster, P. S. (2000). Integrated signals between IL-13, IL-4, and IL-5 regulate airways hyperreactivity. *J. Immunol.* 165, 108–113. doi: 10.4049/jimmunol.165.1.108
- Wenzel, S., Wilbraham, D., Fuller, R., Getz, E. B., and Longphre, M. (2007). Effect of an interleukin-4 variant on late phase asthmatic response to allergen challenge in asthmatic patients: results of two phase 2a studies. *Lancet* 370, 1422–1431. doi: 10.1016/S0140-6736(07)61600-6
- Wenzel, S. E. (2012). Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat. Med.* 18, 716–725. doi: 10.1038/nm.2678
- Wills-Karp, M., Luyimbazi, J., Xu, X., Schofield, B., Neben, T. Y., Karp, C. L., et al. (1998). Interleukin-13: central mediator of allergic asthma. *Science* 282, 2258–2261. doi: 10.1126/science.282.5397.2258
- Wills-Karp, M. (2001). IL-12/IL-13 axis in allergic asthma. *J. Allergy Clin. Immunol.* 107, 9–18. doi: 10.1067/mai.2001.112265
- Wood, N., Whitters, M. J., Jacobson, B. A., Witek, J., Sypek, J. P., and Kasaian, M. (2003). Enhanced interleukin (IL)-13 responses in mice lacking IL-13 receptor alpha 2. *J. Exp. Med.* 197, 703–709. doi: 10.1084/jem.20020906
- Wynn, T. A. (2015). Type 2 cytokines: mechanisms and therapeutic strategies. *Nat. Rev. Immunol.* 15, 271–282. doi: 10.1038/nri3831
- Xu, Y., Li, J., Ding, Z., Li, B., Yu, Z., and Tan, W. (2017). Association between IL-13 +1923C/T polymorphism and asthma risk: a meta-analysis based on 26 case-control studies. *Biosci. Rep.* 37, BSR20160505. doi: 10.1042/BSR20160505
- Zhang, F. Q., Han, X. P., Zhang, F., Ma, X., Xiang, D., Yang, X. M., et al. (2017). Therapeutic efficacy of a co-blockade of IL-13 and IL-25 on airway inflammation and remodeling in a mouse model of asthma. *Int. Immunopharmacol.* 46, 133–140. doi: 10.1016/j.intimp.2017.03.005
- Zhang, Y., Cheng, J., Li, Y., He, R., Pan, P., Su, X., et al. (2019). The safety and efficacy of anti-IL-13 treatment with Tralokinumab (CAT-354) in moderate to severe asthma: a systematic review and meta-analysis. *J. Allergy Clin. Immunol. Pract.* 2661–2671. doi: 10.1016/j.jaip.2019.05.030
- Zhu, Z., Homer, R. J., Wang, Z., Chen, Q., Geba, G. P., Wang, J., et al. (1999). Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J. Clin. Invest.* 103, 779–788. doi: 10.1172/JCI5909

**Conflict of Interest:** EH is advisory board member for AstraZeneca, Sanofi-Genzyme, Novartis, GSK, Circassia, Nestlé, and Purina.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Marone, Granata, Pucino, Pecoraro, Heffler, Loffredo, Scadding and Varricchi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.