

Designing a feasible exercise intervention in first-episode psychosis

Fisher, Emily; Wood, Stephen; Upthegrove, Rachel; Aldred, Sarah

DOI:

[10.1016/j.psychres.2020.112840](https://doi.org/10.1016/j.psychres.2020.112840)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Fisher, E, Wood, S, Upthegrove, R & Aldred, S 2020, 'Designing a feasible exercise intervention in first-episode psychosis: exercise quality, engagement and effect', *Psychiatry Research*, vol. 286, 112840. <https://doi.org/10.1016/j.psychres.2020.112840>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Table 1. Baseline characteristics for both exercise and control groups

	<i>Exercise</i>		<i>Control</i>		p	Effect size
	Average	SD	Average	SD		
Age (years)	23.5	3.8	26.1	5.7	0.22	-0.54
Duration of illness (months)	20.6	10.0	18.0	11.7	0.61	0.24
<i>Body composition</i>						
Weight (kg)	82.5	14.7	84.1	16.8	0.82	-0.10
BMI (kg/m ²)	25.5	3.7	25.5	4.7	0.98	-0.01
<i>Smoking status</i>						
Tobacco consumption	73.0%		64.0%			NA
Cannabis consumption	50.0%		91.0%			NA
<i>Habitual activity</i>						
Garmin Steps (steps/day)	6508.4	1712.2	4735.4	2051.2	0.05	0.86
Garmin floors (floors climbed/day)	7.3	4.6	5.5	3.8	0.39	0.42
Garmin intensity minutes (min/day)	29.5	68.6	30.5	32.3	0.94	-0.04
Garmin active calories (kcal/day)	861.4	475.4	1041.8	914.6	0.58	-0.24
IPAQ MET mins (min/day)	884.2	711.3	1231.9	1547.9	0.51	-0.29
IPAQ Sitting mins (min/day)	1939.5	931.4	1491.4	1804.1	0.98	0.01
<i>Symptoms and functional measures</i>						
PANSS Positive	16.7	4.8	17.8	5.6	0.63	-0.21
PANSS Negative	19.6	3.6	19.5	6.1	0.97	0.01
PANSS general	40.2	9.3	39.0	10.0	0.85	0.13
WHOQOL Overall (%)	45.4	17.8	48.1	16.7	0.94	-0.27
WHODAS Overall (%)	40.8	24.3	24.6	16.5	0.08	0.74
	n	%	n	%		
<i>Medication status</i>						
Olanzapine	1	29	2	25		
Amisulpride	3	43	0	0		
Risperidone	1	14	2	25		
Paliperidone	2	29	0	0		
Depixol	1	14	1	13		
Quetiapine	1	14	2	25		
Aripiprazole	1	14	0	0		
Cloxipol	0	0	1	13		
Haloperidol	0	0	1	13		
Anti-depressant prescription	1	14	4	50		
Anti-anxiety prescription	1	14	2	25		

Group mean values, standard deviation, P values (unpaired t-test) and effect sizes (Cohen's D) are expressed. BMI: body mass index, RHR: resting heart rate, PANSS: positive and negative symptom scale, IPAQ: international physical activity questionnaire, WHOQOL: World Health Organisation quality of life scale, WHODAS: World Health Organisation disability assessment scale.