

# Deep eutectic solvents as media for the prebiotic DNA-templated synthesis of peptides

Nuñez-Pertiñez, Samuel; Wilks, Thomas

DOI:

[10.3389/fchem.2020.00041](https://doi.org/10.3389/fchem.2020.00041)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Nuñez-Pertiñez, S & Wilks, T 2020, 'Deep eutectic solvents as media for the prebiotic DNA-templated synthesis of peptides', *Frontiers in Chemistry*, vol. 8, 41. <https://doi.org/10.3389/fchem.2020.00041>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



# Deep Eutectic Solvents as Media for the Prebiotic DNA-Templated Synthesis of Peptides

Samuel Núñez-Pertíñez and Thomas R. Wilks\*

School of Chemistry, University of Birmingham, Birmingham, United Kingdom

## OPEN ACCESS

### Edited by:

Janarthanan Jayawickramarajah,  
Tulane University, United States

### Reviewed by:

Sriram Kanvah,  
Indian Institute of Technology  
Gandhinagar, India  
Harekrushna Sahoo,  
National Institute of Technology  
Rourkela, India

### \*Correspondence:

Thomas R. Wilks  
t.r.wilks@bham.ac.uk

### Specialty section:

This article was submitted to  
Supramolecular Chemistry,  
a section of the journal  
Frontiers in Chemistry

**Received:** 08 November 2019

**Accepted:** 14 January 2020

**Published:** 31 January 2020

### Citation:

Núñez-Pertíñez S and Wilks TR (2020)  
Deep Eutectic Solvents as Media for  
the Prebiotic DNA-Templated  
Synthesis of Peptides.  
Front. Chem. 8:41.  
doi: 10.3389/fchem.2020.00041

Translation of genetic information into peptide products is one of the fundamental processes of biology. How this occurred prebiotically, in the absence of enzyme catalysts, is an intriguing question. Nucleic acid-templated synthesis (NATS) promotes reactions by bringing building blocks tethered to complementary DNA strands into close proximity and has been shown to enable peptide synthesis without enzymes—it could therefore serve as a model for prebiotic translation of information stored in nucleic acid sequences into functional peptides. The decomposition of highly reactive DNA adapters has so far limited the effectiveness of NATS, but these studies have been performed exclusively in aqueous solution. Deep eutectic solvents (DESs) have been proposed as a feasible solvent for prebiotic replication of nucleic acids, and here are studied as media for prebiotic translation using NATS as a model. DESs are shown to enhance the stability of DNA-conjugated activated esters, the precursors of peptides. However, this enhanced stability was coupled with decreased amine reactivity that hampered the formation of peptide bonds in DESs. These properties are exploited to demonstrate the storage of DNA-conjugated activated esters in a DES followed by transfer into aqueous buffer to activate the NATS of peptides “on demand.” These findings, together with the reported functions of DESs in prebiotic processes, shed light on how DESs could have facilitated the non-enzymatic translation of genetic information into functional peptides on the early Earth.

**Keywords:** prebiotic, translation, deep eutectic solvent, templated synthesis, nucleic acids

## INTRODUCTION

Living organisms preserve the information required to synthesize their molecular components in nucleic acid sequences (Crick, 1970). The replication of this genetic material, and its translation into active catalytic proteins is regulated and catalyzed by highly complex molecular machines. Perhaps the most impressive example is the biosynthesis of peptides and proteins by the ribosome (Steitz, 2008; Yonath, 2009). A key question is how these processes were performed prebiotically, in the absence of such complex machinery. Based on the fact that the core of the ribosome is made from RNA, it has been proposed that replication and translation were first performed by simple nucleic acids, giving rise to the “RNA world” (Crick, 1968; Orgel, 1968; Kruger et al., 1982; Guerrier-Takada et al., 1983; Gilbert, 1986) and “RNA/peptide world” hypotheses (Yarus, 2001; Turk et al., 2010). Several nucleic acid systems that display enzyme-free self-replication have been described (Ekland and Bartel, 1996; Johnston et al., 2001; Paul and Joyce, 2002; Li et al., 2017; Hänle and Richert, 2018; Liu et al., 2018; Mariani et al., 2018; Zhang et al., 2019), and there have been a handful of

mechanisms proposed for prebiotic translation (Zhang and Cech, 1997; Tamura and Schimmel, 2001, 2004). However, these studies have highlighted a number of obstacles to achieving efficient replication and translation in the absence of enzymes: (1) the hydrolysis of the highly reactive, activated monomers required for both processes is rapid in aqueous solution (Kanavarioti et al., 1989); (2) hydrolysed monomers can inhibit replication and induce errors in translation (Deck et al., 2011); (3) after replication, long strands will hybridize to form stable duplexes, blocking further replication by template inhibition (Szostak, 2012).

Most studies in this area have assumed that the solvent for these prebiotic processes was aqueous, but deep eutectic solvents (DESSs) have recently emerged as an alternative (He et al., 2017). DESSs are a family of solvents, closely related to ionic liquids, composed of a hydrogen bond donor and a salt that form a eutectic mixture at a specific molar ratio (Abbott et al., 2003). Most DESSs have high viscosity, low vapor pressure and a very high concentration of ionic species (Zhang et al., 2012; Smith et al., 2014), and their basic components, such as glycerol or urea, are likely to have been present on the early Earth (Kaiser et al., 2015; Okamura et al., 2019). DESSs composed of quaternary ammonium salts and hydrogen bond donors (e.g., reline, a mixture of urea and choline chloride) have been shown to be compatible with protein and nucleic acid biomolecules (Wagle et al., 2014; Xu et al., 2017; Pätzold et al., 2019). For example, it was demonstrated that DNA was stable and retained its structural integrity for at least 6 months when stored in a DES (Mondal et al., 2013), that tertiary structures such as A- and B-DNA duplexes, and G-quadruplexes, were maintained (Mamajanov et al., 2010; Lannan et al., 2012; Zhao et al., 2013; Gállego et al., 2015), and that enzymatic peptide synthesis could be performed efficiently (Maugeri et al., 2013). DESSs have also been shown to be favorable media for nucleic acid replication. For example, DESSs promote the formation of activated organophosphates from mineral phosphate, which is a necessary step in the formation of the activated monomers required for replication (Gull et al., 2014, 2017; Burcar et al., 2016). The high viscosities of DESSs also enable more efficient non-enzymatic replication: the faster diffusion of monomers compared to long oligonucleotides in these solvents has been shown to help overcome the template inhibition problem (He et al., 2017). High viscosity also favors intramolecular folding over intermolecular interactions, enabling, for example, the recovery of ribozyme activity following a replication process (He et al., 2019).

Based on this previous work around replication, we asked whether DESSs could also be suitable media for non-enzymatic translation and chose nucleic acid-templated synthesis (NATS) of a dipeptide as a simple model for this process. NATS promotes reactions by using hybridization to bring reactants tethered to complementary oligonucleotides into close proximity, and has been extensively used in the literature for the programmed production of oligopeptides (Li and Liu, 2004; O'Reilly et al., 2017). Importantly, this previous work has been limited by hydrolysis of the activated building blocks required (Meng et al., 2016). Here, we investigate the NATS of peptides in an archetypal DES, glycholine (a

mixture of glycerol and choline chloride). We find that glycholine protects DNA-conjugated activated esters, the necessary precursors of oligopeptide products, from solvolysis. By comparing the NATS of peptides in DESSs and aqueous buffered solution we also uncover the inhibitory effect of glycholine on amide bond formation. We exploit these properties to demonstrate the extended storage of otherwise unstable peptide building blocks in glycholine, followed by shuttling into aqueous solution to activate NATS “on demand.” Combined with recent theories about the role of compartmentalisation in prebiotic self-replicating systems (Mann, 2012), these experiments hint at a possible role for DESSs in enabling non-enzymatic construction of oligopeptides on the early Earth.

## MATERIALS AND METHODS

**Nuclear magnetic resonance (NMR):** One-dimensional  $^1\text{H}$  and  $^{13}\text{C}$  NMR, and two-dimensional NMR spectra were recorded on a Bruker Advance 300 MHz, Bruker Advance III 400 MHz or a Bruker Advance III 500 MHz instrument at 25°C. Deuterated solvents chloroform-*d* (99.8 D atom%), methanol-*d*<sub>4</sub> (99.8 D atom%), dimethyl sulfoxide-*d*<sub>6</sub> (DMSO-*d*<sub>6</sub>, 99.8 D atom%) acetone-*d*<sub>6</sub> (99.9 D atom%) were purchased from Sigma Aldrich. The residual non-deuterated solvent peak was used as a chemical shift ( $\delta$ , ppm) internal standard. The data were processed using Mestrenova (Mestrelab research) v.12.0.2 and ADC/NMR software. **High resolution mass spectrometry (HRMS):** HRMS was performed on a Bruker Q-ToF Maxis Plus spectrometer or on a Waters Xevo GS2-XS qToF system. A relative error under 5 ppm was ensured. **Reversed-phase high performance liquid chromatography (RP-HPLC):** RP-HPLC was performed in a Varian 920LC system with a photodiode array UV detector (PDA), and a fluorescence detector. **Liquid chromatography-Mass spectrometry (LC-MS):** LC-MS was performed on an Agilent 1200 HPLC system coupled to a Bruker AmazonX high resolution ion trap, in negative ion mode. The desalted oligonucleotide samples were eluted through a XBridge oligonucleotide BEH C18 column (130 Å, 2.5  $\mu\text{m}$ , 4.6  $\times$  50 mm) using a 5 vol% MeOH, 10 mM ammonium acetate (buffer A) and a 70 vol% MeOH, 10 mM ammonium acetate (buffer B) solvent system. The data were processed using Compass Data Analysis (Bruker) v.4.1 software, and the MaxEnt integrated deconvolution algorithm. Alternatively, LC-MS was performed on a Waters ACQUITY UPLC system coupled to a Xevo GS2-XS qToF mass spectrometer in negative sensitivity mode with leucine-enkephalin  $[\text{M}-\text{H}]^-$  554.2620 Lockspray. The oligonucleotides were eluted through an ACQUITY UPLC oligonucleotide BEH C18 column (130 Å, 1.7  $\mu\text{m}$ , 2.1  $\times$  50 mm) using a 50 mM triethylammonium acetate (TEAA, pH 7.0) solution in H<sub>2</sub>O (buffer A) and a 50 mM TEAA solution in MeCN (buffer B) at 60°C and a 0.2 mL·min<sup>-1</sup> flow. **Fluorescence spectroscopy:** Fluorescence spectral data were recorded on an Agilent Cary Eclipse fluorescence spectrophotometer equipped with a photomultiplier tube (PMT) detector. Quartz cuvettes from Starna scientific (Type 3/Q/10) with four polished sides were used for fluorescence. The emission

and excitation spectra were recorded using Cary Eclipse v.1.2.0.0 software. **Karl Fischer titration:** The water content in DESSs was determined using an automated CA-200 (Mitsubishi chemicals) Karl Fischer coulometric titrator. Aqualine (Fisher) auxiliary reagent mixture and Aquamicon solution P (Aquamicon, 3.8–4.0 mg H<sub>2</sub>O·mL<sup>-1</sup>) calibration standards were used. **pH measurements:** An Accumet AP110 pH meter kit (Fisher) equipped with a pH sensitive glass electrode was used to determine the pH in aqueous solution. Prior to sample analysis, the instrument was calibrated with pH 4, pH 7, and pH 10 standard solution buffers.

## DNA Sequences and Modifications

Sequences were optimized, and the expected assemblies predicted with NUPACK (Zadeh et al., 2009). Modified DNA strands were purchased from Integrated DNA Technologies and further modified as described below. Sequence data and modification structures are given in **Table S1** and **Scheme S1**.

## Synthesis of DESSs

DESSs were prepared by mixing appropriate amounts of hydrogen bond donor and choline chloride in a Schlenk round bottom flask under positive nitrogen pressure, heating to 100°C and stirring until a homogeneous mixture was obtained. DESSs were dried at 80°C under vacuum prior to use (Abbott et al., 2004). The H<sub>2</sub>O content of glycholine was determined by Karl Fischer titration. H<sub>2</sub>O content was then adjusted to the desired value. For simplicity, we refer to the volume percentage content of H<sub>2</sub>O in the DES using a subscript number e.g., glycholine containing 7 vol% H<sub>2</sub>O is written as Gly<sub>0.07</sub>.

## Measurement of DES Solution pH

DES (1 mL) was diluted in H<sub>2</sub>O (4 mL) and the pH was determined using a pH-sensitive glass electrode. The error was evaluated in two different series of samples, by performing triplicate measurements and determining the standard deviation.

## <sup>1</sup>H-NMR Spectroscopy Study of The Stability of Small Molecule Activated Esters

Solutions of **1** (5 mM) in different solvent mixtures were stirred at 24 °C. After 2, 4, 6 and 24 h the solutions were extracted

with 0.7 mL of either acetone-*d*<sub>6</sub> (DES solutions) or chloroform-*d* (phosphate buffer solutions). The relative proportions of **1** and **3** were determined by <sup>1</sup>H-NMR spectroscopy (**Figure S1**).

## RP-HPLC Stability Study of Activated Esters


Activated ester **1** (4.1 mg) was dissolved in DES (2 mL). Each solution was prepared 40 min after the previous one. The solutions were stirred over a period of 29 h. After stirring for 5, 24, and 29 h, a 100 μL aliquot of each sample was diluted in H<sub>2</sub>O (900 μL) and the sample analyzed by RP-HPLC. Method: column Discovery C18 (Sigma Aldrich, 5 μm, 10 × 4.6 cm), flow 1 mL·min<sup>-1</sup>, temperature 40°C and injection volume 10 μL. Solvent A: H<sub>2</sub>O, 0.05 vol% TFA. Solvent B: 70 vol% MeCN, 0.05 vol% TFA.

## Thermal Stability of dsDNA in DESSs

The thermal stability of dsDNA in DES solutions was determined using fluorophore and quencher labeled DNA strands (**S15** and **S16**). Solutions of 100 nM dsDNA were briefly centrifuged and placed in a qPCR instrument. Samples were heated and cooled at 1°C·min<sup>-1</sup>. The melting temperature (*T*<sub>m</sub>) was determined as the middle-point of the sigmoidal trace defined by the fluorescence emission intensity as a function of temperature.

## Synthesis of NHS-Activated Fluorescein, **S18** (Gao et al., 2002)

Fluorescein free carboxylic acid (**S17**, 3.322 g, 10 mmol) and *N*-hydroxysuccinimide (1.151 g, 10 mmol) were dissolved in dry DMF (19 mL) under positive nitrogen pressure in a dry Schlenk tube (**Scheme 1**). Then, *N,N'*-dicyclohexylcarbodiimide (DCC, 2.063 g, 10 mmol) was added and the resulting solution stirred at 60°C for 2 h under positive nitrogen pressure. *N,N'*-dicyclohexylurea (DCU) was removed by cooling down the reaction mixture to -20°C for 2 h and the white precipitate was filtered off. The solvent was removed under reduced pressure and the crude solid was purified by SiO<sub>2</sub> column chromatography using an EtOAc/acetone gradient. The desired fractions were combined to produce **S18** as a bright orange solid (2.002 g, 47%). The <sup>1</sup>H-NMR spectroscopic analysis was in good agreement with the reported characterization (Gao et al., 2002). **TLC** (EtOAc:MeOH 6:4) *R*<sub>f</sub> = 0.1. **<sup>1</sup>H-NMR** (300 MHz, DMSO-*d*<sub>6</sub>) δ (ppm): 8.36 (1H, *d* *J* = 7.7 Hz, Ar-*H*), 7.96 (2H, *m*, Ar-*H*), 7.65 (1H, *d* *J* = 7.5 Hz, Ar-*H*), 6.79 (2H, *d* *J* = 9.6 Hz, Ar-*H*), 6.53 (4H,



**SCHEME 1** | Synthesis of the *N*-hydroxymaleimide ester **S21**. (i) SuOH, DCC, DMF, 60°C, 2 h; (ii) TsOH·H-Ala-OBn, Et<sub>3</sub>N, DCM, r.t. 3 h; (iii) Pd/C (10 wt%), H<sub>2</sub>, MeOH, r.t., 1.5 h; (iv) *N*-hydroxymaleimide, DCC, EtOAc:1,4-dioxane (2:1), r.t., o.n.

m, Ar-H), 2.73 (4H, s<sup>br</sup>, (CH<sub>2</sub>)<sub>2</sub> succinimide). **HRMS** (ESI<sup>-</sup>) *m/z* [M-H]<sup>-</sup> calcd. 428.0770 found 428.0763.

## Synthesis of S19

**S18** (1.000 g, 2.23 mmol) and TsOH·H-L-Ala-OBn (1.637 g) were dissolved in dry DMF (15 mL) under positive nitrogen pressure in a dry round bottom Schlenk flask. Then, Et<sub>3</sub>N (1.1 mL, 8.16 mmol) was added drop wise over 10 min and the resulting solution was stirred at room temperature for 6 h. The solvent was removed under reduced pressure and the crude product was purified by SiO<sub>2</sub> column chromatography using a DCM/MeOH (100:0 to 75:15) gradient to produce **S19** as a bright orange solid (704 mg, 61%). **TLC** (EtOAc:MeOH 6:4) R<sub>f</sub> = 0.6. **<sup>1</sup>H-NMR** (400 MHz, methanol-*d*<sub>4</sub>) δ (ppm): 7.76 (1H, m, Ar-H), 7.55 (1H, m, Ar-H), 7.41 (1H, m, Ar-H), 7.21 (2H, m, Ar-H), 7.11 (4H, m, Ar-H), 6.88 (1H, m, Ar-H), 6.56 (1H, m, Ar-H), 6.51 (1H, m, Ar-H), 6.45 (1H, m, Ar-H), 6.20 (1H, d *J* = 8.7 Hz, Ar-H), 5.97 (1H, dd *J*<sub>1</sub> = 8.7 Hz *J*<sub>2</sub> = 2.5 Hz, Ar-H), 4.80 (2H, m, OCH<sub>2</sub>), 3.59 (1H, q *J* = 7.2 Hz, CH-CH<sub>3</sub>), 1.02 (3H, d *J* = 7.2 Hz, CH-CH<sub>3</sub>). **<sup>13</sup>C-NMR** (100 MHz, methanol-*d*<sub>4</sub>) δ (ppm): 171.75 (C=O), 168.79 (C=O), 160.29 (C=O), 160.13 (ArC), 154.54 (ArC), 154.36 (ArC), 154.24 (ArC), 136.95 (ArC), 134.40 (ArC-H), 132.13 (ArC-H), 130.71 (ArC-H), 130.60 (ArC-H), 129.89 (ArC-H), 129.53 (ArC), 129.37 (ArC-H), 129.35 (ArC-H), 129.15 (ArC), 129.09 (ArC), 125.13 (ArC-H), 123.55 (ArC-H), 113.26 (ArC-H), 112.95 (ArC-H), 110.26 (ArC), 109.24 (ArC), 103.59 (ArC-H), 103.08 (ArC-H), 68.08 (OCH<sub>2</sub>), 51.55 (NCH), 15.44 (CH<sub>3</sub>). **HRMS** (ESI<sup>-</sup>) *m/z* [M-H]<sup>-</sup> calcd. 492.1447 found 492.1457.

## Synthesis of S20

Pd/charcoal (50 mg, 10 wt%) was placed in a dry round bottom Schlenk flask under positive nitrogen pressure. **S19** (500 mg, 1.01 mmol) was added and dissolved in degassed MeOH (50 mL). The nitrogen atmosphere was replaced with hydrogen by bubbling the solution for 3 min with a hydrogen balloon. The positive hydrogen pressure was maintained for 1.5 h with constant stirring at room temperature. The solid-supported catalyst was removed by filtration through celite. The solvent was removed under reduced pressure to produce **S20** as a bright yellow solid (375 mg, 92%). **TLC** (DCM:MeOH 9:1) R<sub>f</sub> = 0.2. **<sup>1</sup>H-NMR** (500 MHz, methanol-*d*<sub>4</sub>) δ (ppm): 7.05 (1H, m, Ar-H), 6.61 (3H, m, Ar-H), 6.48 (2H, m, Ar-H), 6.41 (1H, dd *J*<sub>1</sub> = 8.7 Hz *J*<sub>2</sub> = 2.4 Hz, Ar-H), 3.67 (1H, q *J* = 7.2 Hz, CHCH<sub>3</sub>), 1.20 (3H, d *J* = 7.2 Hz, CHCH<sub>3</sub>). **<sup>13</sup>C-NMR** (160 MHz, methanol-*d*<sub>4</sub>) δ (ppm): 173.6 (C=O), 169.1 (C=O), 160.6 (C=O), 160.4 (ArC), 154.9 (ArC), 154.6 (ArC), 154.4 (ArC), 134.5 (ArC), 132.5 (ArC), 132.4 (ArC-H), 130.9 (ArC-H), 130.0 (ArC-H), 113.4 (ArC-H), 113.0 (ArC-H), 110.6 (ArC), 109.6 (ArC), 103.7 (ArC-H), 103.2 (ArC-H), 51.7 (CH-CH<sub>3</sub>), 15.7 (CH-CH<sub>3</sub>). **HRMS** (ESI<sup>-</sup>) *m/z* [M-H]<sup>-</sup> calcd. 402.0983 found 402.0982.

## Synthesis of S21

**S20** (50 mg, 0.124 mmol), DCC (25 mg, 0.121 mmol) and *N*-hydroxymaleimide (14 mg, 0.124 mmol) were dissolved in EtOAc/1,4-dioxane (2:1, 2 mL) and stirred at room temperature overnight. Then, the solvent was lyophilised on a Schlenk line, the crude solid was dissolved in EtOAc (2.5 mL) and cooled down

to 4°C for 1 h. The white precipitate of DCU was removed by centrifugation at 4°C and 21 kRCF for 15 min. The supernatant was dried to produce **S21** as a bright yellow powder (58 mg, 95%). The NMR spectra showed extra peaks which were attributed to atropoisomers and the presence of trace amounts of DCU. **<sup>1</sup>H-NMR** (500 MHz, DMSO-*d*<sub>6</sub>) δ (ppm): 9.99 (1H, s, OH or NH), 9.89 (1H, s, OH or NH), 7.83 (1H, m, Ar-H), 7.57 (2H, m, Ar-H), 7.17 (2H, s<sup>br</sup>, maleimide Ar-H), 7.05 (1H, m, Ar-H), 6.61 (2H, m, Ar-H), 6.55 (1H, m, Ar-H), 6.49 (2H, m, Ar-H), 6.38 (1H, m, dd *J*<sub>1</sub> = 8.7 Hz *J*<sub>2</sub> = 2.4 Hz, Ar-H), 4.11 (1H, q *J* = 7.1 Hz, CH-CH<sub>3</sub>), 1.26 (3H, d *J* = 7.1 Hz, CH-CH<sub>3</sub>). **<sup>13</sup>C-NMR** (160 MHz, DMSO-*d*<sub>6</sub>) δ (ppm): 167.28 (C=O), 165.96 (C=O), 164.38 (C=O), 158.71 (C=O), 157.36 (ArC), 153.20 (ArC), 152.13 (ArC), 133.43 and 133.09 (maleimide CH=CH), 131.91 (ArC-H), 130.07 (ArC-H), 129.68 (ArC), 129.45 (ArC-H), 128.78 (ArC-H), 123.85 (ArC-H), 122.58 (ArC-H), 112.37 (ArC-H), 112.11 (ArC-H), 108.47 (ArC), 107.62 (ArC), 102.41 (ArC-H), 102.06 (ArC-H), 48.16 (CH-CH<sub>3</sub>), 15.05 (CH-CH<sub>3</sub>). **HRMS** (ESI<sup>-</sup>) *m/z* [M-H]<sup>-</sup> calcd. 497.0985 found 497.0992.

## Fluorescence Spectroscopy Study of the Stability of DNA-Conjugated Activated Esters

The tetramethylrhodamine (TAMRA)/disulfide-labeled DNA hairpin **S4** (1 μL, 1 mM in H<sub>2</sub>O, 1 nmol) was thermally annealed with the complementary biotinylated strand **S5** (1.1 μL, 1.1 mM in H<sub>2</sub>O, 1.1 nmol) in phosphate buffer solution (PBS, 10 μL, 0.1 M, pH 7.0). The disulfide group was reduced with tris(2-carboxyethyl)phosphine hydrochloride (TCEP, 0.5 μL, 0.5 M in H<sub>2</sub>O) at 18°C for 1.25 h. Then, the resulting double stranded DNA (dsDNA) was conjugated to streptavidin-coated magnetic nanoparticles (Dynabeads, Thermofisher), and transferred into Gly<sub>0.07</sub> (200 μL). The maleimide ester **S21** (1 mg, 2.5 μmol) was added to the solution and allowed to react for 2 h. The nanoparticle-supported DNA-conjugated activated ester was transferred into clean Gly<sub>0.07</sub> and the hairpin was released by toehold-mediated strand exchange by addition of one equivalent of strand **S6** at 18°C for 2 h to produce an approximately 5 μM solution of **4**. Finally, the solution of **4** (5 μL) was diluted in the DES of interest (95 μL), the resulting solution briefly centrifuged, and the fluorescein (FAM) fluorescence recorded over 24 h in a Mx3005P qPCR instrument (Agilent). The spectral overlap between FAM fluorophore (**S21**) and TAMRA-labeled DNA (**S4**) was also determined (**Figure S4**).

## DNA-Templated Synthesis of Peptide Bonds

Disulfide-labeled DNA **S7** (10 μL, 1 mM in H<sub>2</sub>O, 10 nmol) was mixed with TCEP solution (2 μL, pH 4 corrected with 5 M NaOH, 1 μmol) in a microcentrifuge tube. The solution was shaken at 18°C for 1.25 h. Maleimide activated ester **S21** (108 μL, 37 mM in DMF, 4 μmol) was added and the combined solution shaken at 18°C for 1.0 h to produce the DNA-conjugated activated ester, **6**. The volume of the solution was reduced to approximately 50 μL by freeze-drying. The previous solution was diluted with Gly<sub>0.07</sub> (4,950 μL), and

amine-labeled DNA 7 (10  $\mu\text{L}$ , 1 mM in  $\text{H}_2\text{O}$ , 10 nmol) added. The resulting mixture was vigorously shaken until the solution was homogeneous and then shaken at 18°C overnight. The DES solution was diluted with an excess of  $\text{H}_2\text{O}$  and the DNA strands transferred into  $\text{H}_2\text{O}$  for HPLC/LC-MS analysis by repeated centrifugal ultrafiltration using a 3 kDa molecular weight cut-off (MWCO) Amicon spin filter. When the DNA-templated reaction was performed in aqueous solution, the activated ester solution in DMF was diluted with PBS (4,950  $\mu\text{L}$ , 0.1 M, pH 7.5) containing the complementary amine-labeled DNA 7.

### Amine-Labeled DNA Reactivity in DES/Triethylamine ( $\text{Et}_3\text{N}$ )

Amine-labeled DNA 10 (1  $\mu\text{L}$ , 1 mM in  $\text{H}_2\text{O}$ , 1 nmol) was diluted in the appropriate solvent (100  $\mu\text{L}$ ). *N*-hydroxysuccinimide ester 11 (~0.3 mg, 1  $\mu\text{mol}$ ) was added, and the mixture shaken overnight at 20°C. The product was transferred into  $\text{H}_2\text{O}$  by dilution of the samples and successive centrifugal ultrafiltration using a 3 kDa MWCO Amicon spin filter. The product was analyzed by HPLC and LC-MS.

### Storage of Activated Esters in Gly<sub>0.07</sub>


Disulfide-labeled DNA S11 (3  $\mu\text{L}$ , 1 mM in  $\text{H}_2\text{O}$ , 3 nmol) was mixed with TCEP solution (3  $\mu\text{L}$ , pH 4 corrected with 5 M NaOH, 1.5  $\mu\text{mol}$ ) in a microcentrifuge tube. The solution was shaken at 18°C for 1.25 h. Maleimide activated ester S21 (54  $\mu\text{L}$ , 22 mM in DMF, 1.2  $\mu\text{mol}$ ) was added and the combined solution was shaken at 18°C for 1.0 h to produce the DNA-conjugated activated ester 13. This sample was divided into 3 aliquots and used as follows. (1) *Storage in Gly<sub>0.07</sub>*: the activated ester solution (20  $\mu\text{L}$ ) was diluted with Gly<sub>0.07</sub> (180  $\mu\text{L}$ ), and the resulting solution was shaken at 18°C for 2.0 h. The solution was diluted with PBS (1.3 mL, 0.1 M, pH 7.5) containing complementary amine-labeled DNA 15 (1  $\mu\text{L}$ , 1 mM in  $\text{H}_2\text{O}$ , 1 nmol). The solution was shaken at 18°C for 5 h. Finally, the solution was diluted with 1 vol. of  $\text{H}_2\text{O}$ , the precipitate was centrifuged, and the DNA products were transferred into  $\text{H}_2\text{O}$  by successive steps of dilution of the supernatant with  $\text{H}_2\text{O}$  and concentration by centrifugal ultrafiltration on a 3 kDa MWCO Amicon spin filter. The product was analyzed by HPLC and LC-MS. (2) *Storage in PBS*: the activated ester solution (20  $\mu\text{L}$ ) was diluted with PBS (180  $\mu\text{L}$ , 0.1 M, pH 7.0), and the resulting solution was shaken at 18°C for 2.0 h. Then, the sample was treated as in (1). (3) *Fast reaction in PBS*: the activated ester solution (20  $\mu\text{L}$ ) was diluted with PBS (1,480  $\mu\text{L}$ , 0.1 M, pH 7.5) containing complementary amine-labeled DNA 15 (1 nmol). The solution was shaken at 18°C for 5.0 h. The products were transferred into  $\text{H}_2\text{O}$  for LC-MS analysis as described previously. (4) *Off-template control*: maleimide activated ester (S21) (18  $\mu\text{L}$ , 22 mM in DMF, 400 nmol) was diluted in PBS (1,482  $\mu\text{L}$ , 0.1 M, pH 7.5) containing amine-labeled DNA 15 (1  $\mu\text{L}$ , 1 mM in  $\text{H}_2\text{O}$ , 1 nmol). The solution was shaken at 18°C for 5.0 h and the product was transferred into  $\text{H}_2\text{O}$  for LC-MS analysis as described above.

## RESULTS AND DISCUSSION

### The Stability of Activated Esters in DES and Aqueous Buffer

We used the *N*-hydroxysuccinimide ester of *L*-alanine (NHS-Ala) as a model activated peptide building block. This had similar solution stability to more prebiotically feasible activating groups while allowing us to use established techniques to prepare the required DNA-conjugated esters (He and Liu, 2011). A preliminary stability experiment was performed using a small-molecule NHS-Ala, 1. We followed the hydrolysis of 1 in several glycoline/water mixtures and in aqueous PBS using proton nuclear magnetic resonance ( $^1\text{H-NMR}$ ) spectroscopy (Figure 1 and Figure S1). 1 was dissolved in the appropriate solvent mixture and then left for 24 h at 24°C. Samples were extracted with immiscible deuterated solvent at set timepoints, and the relative proportions of activated ester and degradation products (2 and 3) were determined based on their distinctive methyl  $^1\text{H-NMR}$  signals. Both the proportion of water and the presence of basic additives were investigated for their effect on the rate of ester degradation.

As expected, rapid degradation of the ester in aqueous PBS was observed, with only 25% remaining after 24 h. In stark contrast, degradation in Gly<sub>0.07</sub> was very slow, with ~90% of the activated ester intact after 24 h. Increasing the  $\text{H}_2\text{O}$  content to 50 vol% accelerated solvolysis substantially. We attempted to deliberately accelerate the degradation in glycoline by use of a basic additive, triethylamine ( $\text{Et}_3\text{N}$ ). The presence of 2 mM  $\text{Et}_3\text{N}$  in Gly<sub>0.07</sub> had the same destabilizing effect as increasing the water content to 50 vol%, while 10 mM  $\text{Et}_3\text{N}$  led to a similar rate of degradation to that observed in PBS. The effect of basic additives was studied by diluting the DES in  $\text{H}_2\text{O}$



and measuring the pH of the resulting solution. 2 mM Et<sub>3</sub>N produced a pH of 7.5, while 10 mM Et<sub>3</sub>N produced a pH of 9.9 (Table S2), it therefore seems likely that the observed differences in degradation rate arose as a result of the increased basicity of the solution.

The above <sup>1</sup>H-NMR analysis could have been biased by differences in the extraction efficiency of **1** and **3** into the deuterated solvent used, so a complementary study that did not rely on this assumption was performed using reversed-phase high performance liquid chromatography (RP-HPLC, Figure S2). The results were highly consistent with the <sup>1</sup>H-NMR study, so we concluded that glycoline was indeed effective at stabilizing activated esters, and moved on to DNA-based experiments.


The thermal stability of double-stranded DNA (dsDNA) in glycoline/H<sub>2</sub>O was assessed by fluorescence quenching and revealed that the duplex remained stable down to at least 7 vol% H<sub>2</sub>O: in Gly<sub>0.07</sub> the *T<sub>m</sub>* of a 24 base pair dsDNA was 48°C (Figure S3). The addition of moderate volumes of H<sub>2</sub>O to DES has been used to reduce its viscosity and melting point (Ma et al., 2018; Smith et al., 2019), and resulted in increased conversion in enzymatic reactions (Durand et al., 2013; Guajardo et al., 2017). Furthermore, it has been shown that the microstructure of the DES is at least partially preserved upon the addition H<sub>2</sub>O, up to 50 wt% (Hammond et al., 2017; Gabriele et al., 2019). Given this precedent, as well as the good stability of activated esters demonstrated above, Gly<sub>0.07</sub> was used for experiments from this point onwards.

We were interested in assessing the stability of DNA-conjugated activated esters *in situ*, avoiding the possibility

of degradation occurring during analysis. To this end, a fluorescence quenching experiment was designed (Figure 2). A FAM-labeled NHS-Ala (S21) was synthesized from S17 through a multi-step procedure (Scheme 1 and Scheme S2, intermediates S18 to S20) and conjugated to a TAMRA-labeled DNA hairpin through thia-Michael addition. This design meant that in the ester form (**4**), FAM fluorescence would be quenched by the neighboring TAMRA group. Upon solvolysis of the ester to give **5**, the FAM-labeled amino acid would diffuse away, triggering a measurable increase in fluorescence. We measured the fluorescence excitation and emission spectra of FAM and TAMRA in glycoline to ensure good spectral overlap was maintained in this solvent (Figure S4).

The DNA hairpin was supported on streptavidin-coated magnetic nanoparticles and transferred to Gly<sub>0.07</sub> for activated ester synthesis, ensuring minimal hydrolysis of the activated ester. Excess reagents were removed by magnetic purification, the hairpins were released into solution by toehold-mediated strand displacement and the nanoparticles removed with a magnet. The hairpin solution was diluted into an excess of the solvent of interest, and the evolution of the fluorescence over a 24 h period was recorded (Figure S5).


All experiments showed the expected increase in fluorescence due to solvolysis of the activated ester (Figure 2). While a rigorous quantitative kinetic analysis of the data was not possible due to uncertainties in the minimum and maximum fluorescence values, the results showed exactly the same trend as the small molecule experiments, with slow hydrolysis observed in Gly<sub>0.07</sub>. We therefore concluded that glycoline was also effective at stabilizing DNA-conjugated activated esters.



## Nucleic Acid-Templated Synthesis of Peptide Bonds in DES


Having demonstrated the higher stability of the activated esters in glycoline, we moved to a single-step NATS reaction using

NHS-Ala (**Figure 3A**). The DNA-conjugated activated ester **6** was prepared in 90 vol% DMF and then transferred into a DES solution containing the complementary acceptor strand **7**. The DMF content was reduced to 1 vol%, which did not substantially



**FIGURE 3** | DNA-templated formation of peptide bonds in DES. **(A)** Single-step DNA-templated synthesis of a peptide bond in glycoline and in PBS. The conversion was determined by HPLC integration of the signals corresponding to **7**, **9**, and **9'** (isomers of the product). Peaks were identified by LC-MS (Bruker AmazonX): **7** [M-H]<sup>-</sup> *m/z* calcd. 9337.2 found 9336.0; \*DNA-SH [M-H]<sup>-</sup> *m/z* calcd. 10615.2 found 10599.6; **8** [M-H]<sup>-</sup> *m/z* calcd. 10712.0 found 10712.4; **9** and **9'** (isomers) [M-H]<sup>-</sup> *m/z* calcd. 9723.6 **(B)** comparison of the reactivity of an amine-labeled DNA (**10**) in the presence of an excess of activated ester (**11**) in Gly<sub>0.07</sub> / Et<sub>3</sub>N and 1 M NaCl (aq)/Et<sub>3</sub>N. The conversion was determined by HPLC integration of the signals corresponding to **10** and **12**. Peaks were identified by LC-MS (Bruker AmazonX): **10** [M-H]<sup>-</sup> *m/z* calcd. 3897.7 found 3898.6; **11** [M-H]<sup>-</sup> *m/z* calcd. 4096.8 found 4097.7. Error bars correspond to the standard deviation over multiple integration events.





**FIGURE 4** | Activating NATS “on demand.” **(A)** Scheme of the storage of a DNA conjugated activated ester in PBS (0.1 M pH 7.0) or  $\text{Gly}_{0.07}$ , followed by DNA-templated peptide bond formation in PBS. **(B)** LC-MS chromatograms of the products (Waters Xevo-G2-XS). 15  $[\text{M}]^0$   $m/z$  calcd. 10891.855 found 10891.522; 14  $[\text{M}]^0$   $m/z$  calcd. 9558.632 found 9558.581. 16  $[\text{M}]^0$   $m/z$  calcd. 11201.935 found 11201.489.

affect the duplex  $T_m$ . Alternatively, **6** was synthesized in Gly<sub>0.07</sub>; this required longer reaction times (2 h) to reach the same yield attained in DMF, which we speculate was due to the higher viscosity of the DES. Controls mixing the acceptor strand with the small molecule reagents showed that no off-template reactions took place.

A control NATS reaction in PBS was also performed. While hydrolysis of the activated esters in aqueous solution was very fast, if the DNA conjugate **6** was isolated rapidly and transferred to a solution that contained the complementary amine-labeled DNA **7**, the amide product **9** was formed in 48% conversion. By contrast, the formation of **9** in DES was only detected in very low yield (**Figure 3A**). A series of attempts to improve the conversion through the use of additives were not successful in raising the yield of **9** above ~5% (**Table S3**). We hypothesized that the low conversion in DES was due to a lack of reactivity of the amine moiety. To test this, amino-modified DNA **10** was reacted with a large excess of a small-molecule activated ester **11** in aqueous 1 M NaCl and glycholine solution that contained varying concentrations of Et<sub>3</sub>N (**Figure 3B**). In aqueous solution, the conversion in the absence of base was very low, however the addition of a minimal amount of Et<sub>3</sub>N resulted in nearly quantitative conversion. By contrast, in DES, very large amounts of Et<sub>3</sub>N were required to achieve only moderate conversion (60%). It is interesting to note that NATS of peptide bonds has previously been reported in pH 7.5 buffered solution, similar to the pH of the glycholine/Et<sub>3</sub>N solutions measured above, so pH cannot by itself explain the lack of amine reactivity in these solvent mixtures (He and Liu, 2010).

These results highlighted that glycholine by itself could not have promoted prebiotic peptide synthesis. Stabilization of the activated ester comes at the cost of reduced amine reactivity, and increasing the reactivity of the amine can only be achieved by use of additives that destabilize the ester. However, recent hypotheses have proposed an important role for compartmentalisation in prebiotic processes (Mann, 2012). One could imagine a prebiotic system in which activated species are generated and stored in pockets of DES, and then transferred into aqueous solution to facilitate the formation of peptide products. We designed a transfer experiment to investigate this possibility (**Figure 4A**).

The DNA-conjugated activated ester **13** was stored in PBS or glycholine for 2 h. The solution was then diluted with an excess of PBS containing the acceptor amino-DNA strand **15**. The templated synthesis of the peptide product **16** was then quantified via HPLC by comparing peak areas (**Figure 4B**). The identity of the product was confirmed by LC-MS. A positive control was performed by removing the storage step and adding the acceptor strand (**15**) immediately after the synthesis of the activated ester **13**. This resulted in 68% conversion to the peptide product. To check the importance of DNA templating, an off-template negative control was performed by adding the acceptor strand **15** to a solution of the small molecule activated ester **S21** at the same concentration as the DNA strand (670 nM). This showed no evidence of the

formation of the peptide product **16**, as expected. When the DNA-templated peptide synthesis was attempted following storage of the activated ester in PBS for 2 h, no product was observed (**Figure 4B**). By contrast, DNA-templated synthesis following storage in glycholine resulted in 30% conversion to the peptide product.

## DISCUSSION

Here, we demonstrate that glycholine-based DESs have a positive impact on the stability of DNA-conjugated activated esters when compared with buffered aqueous solutions. However, there is a marked reduction in the reactivity of amines, which prevents the NATS of peptides in these solvents. Our results highlight that it is very challenging to produce activated esters in aqueous media for subsequent templated reactions, as they quickly decompose; however, the synthesis of the activated ester in Gly<sub>0.07</sub> provides a solution to this problem. By applying a two-stage process in which the activated ester was stored in DES and then transferred into PBS solution to trigger templated synthesis we achieved a moderate yield of 30%. By comparison, no product was observed if the activated ester was stored in PBS.

It may be possible to circumvent the issue of amine deactivation we observe by the use of alternative DESs or other viscous solvents. For example, glycerol containing an appropriate concentration of NaCl has been shown to stabilize DNA duplexes (Bonner and Klibanov, 2000). Future work could investigate whether such solvent mixtures are more effective at promoting nucleic acid-templated peptide synthesis, and thus shine further light on whether increased activated ester stability is inevitably linked to decreased amine reactivity in these environments.

Finally, it is interesting to speculate how our results might be relevant to prebiotic translation. The high viscosity of DESs means that they mix only slowly with aqueous solutions, so this may have allowed a primitive form of compartmentalization in which the production and stabilization of activated species took place in a DES phase, followed by diffusion into the aqueous phase where templated chemistry could occur. Future experiments using simple DES/aqueous interfaces of this kind could provide fascinating insights into such prebiotic processes.

## DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

## AUTHOR CONTRIBUTIONS

SN-P and TW designed the experiments, analyzed the data, and wrote and formatted the manuscript. SN-P carried out the experimental work.

## FUNDING

This work was funded entirely by the European Research Council (grant ID 615142), which provided a studentship for SN-P and

a postdoctoral fellowship for TW. Open access publication fees were funded by the University of Birmingham.

## ACKNOWLEDGMENTS

The authors thank Prof. Rachel O'Reilly for her invaluable support during the performance of this work and Prof. Andrew Turberfield, Dr. Jonathan Bath, and Dr. Robert

Oppenheimer for advice on the design of the DNA constructs and helpful discussions.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fchem.2020.00041/full#supplementary-material>

## REFERENCES

- Abbott, A. P., Boothby, D., Capper, G., Davies, D. L., and Rasheed, R. K. (2004). Deep eutectic solvents formed between choline chloride and carboxylic acids: versatile alternatives to ionic liquids. *J. Am. Chem. Soc.* 126, 9142–9147. doi: 10.1021/ja048266j
- Abbott, A. P., Capper, G., Davies, D. L., Rasheed, R. K., and Tambyrajah, V. (2003). Novel solvent properties of choline chloride/urea mixtures. *Chem. Commun.* 2003, 70–71. doi: 10.1039/b210714g
- Bonner, G., and Klibanov, A. M. (2000). Structural stability of DNA in nonaqueous solvents. *Biotechnol. Bioeng.* 68, 339–344. doi: 10.1002/(SICI)1097-0290(20000505)68:33.O.CO;2-O
- Burcar, B., Pasek, M., Gull, M., Cafferty, B. J., Velasco, F., Hud, N. V., et al. (2016). Darwin's warm little pond: a one-pot reaction for prebiotic phosphorylation and the mobilization of phosphate from minerals in a urea-based solvent. *Angew. Chem. Int. Ed.* 55, 13249–13253. doi: 10.1002/anie.201606239
- Crick, F. (1968). The origin of the genetic code. *J. Mol. Biol.* 38, 367–379. doi: 10.1016/0022-2836(68)90392-6
- Crick, F. (1970). Central dogma of molecular biology. *Nature* 227, 561–563. doi: 10.1038/227561a0
- Deck, C., Jauker, M., and Richert, C. (2011). Efficient enzyme-free copying of all four nucleobases templated by immobilized RNA. *Nat. Chem.* 3, 603–608. doi: 10.1038/nchem.1086
- Durand, E., Lecomte, J., Baréa, B., Dubreucq, E., Lortie, R., and Villeneuve, P. (2013). Evaluation of deep eutectic solvent-water binary mixtures for lipase-catalyzed lipophilization of phenolic acids. *Green Chem.* 15, 2275–2282. doi: 10.1039/c3gc40899j
- Ekland, E. H., and Bartel, D. P. (1996). RNA-catalyzed RNA polymerization using nucleoside triphosphates. *Nature* 382, 373–376. doi: 10.1038/382373a0
- Gabriele, F., Chiarini, M., Germani, R., Tiecco, M., and Spreti, N. (2019). Effect of water addition on choline chloride/glycol deep eutectic solvents: characterization of their structural and physicochemical properties. *J. Mol. Liq.* 291:111301. doi: 10.1016/j.molliq.2019.111301
- Gállego, I., Grover, M. A., and Hud, N. V. (2015). Folding and imaging of DNA nanostructures in anhydrous and hydrated deep-eutectic solvents. *Angew. Chem. Int. Ed.* 54, 6765–6769. doi: 10.1002/anie.201412354
- Gao, J., Wang, P., and Giese, R. W. (2002). Xanthamide fluorescent dyes. *Anal. Chem.* 74, 6397–6401. doi: 10.1021/ac020368+
- Gilbert, W. (1986). Origin of life: the RNA world. *Nature* 319:618. doi: 10.1038/319618a0
- Guajardo, N., Domínguez de María, H. P., Ahumada, K., Schreiber, R. A., Ramírez-Tagle, R., Crespo, F. A., et al. (2017). Water as cosolvent: nonviscous deep eutectic solvents for efficient lipase-catalyzed esterifications. *Chem. Cat. Chem.* 9, 1393–1396. doi: 10.1002/cctc.201601575
- Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N., and Altman, S. (1983). The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* 35, 849–857. doi: 10.1016/0092-8674(83)90117-4
- Gull, M., Cafferty, B. J., Hud, N. V., and Pasek, M. A. (2017). Silicate-promoted phosphorylation of glycerol in non-aqueous solvents: a prebiotically plausible route to organophosphates. *Life* 7:29. doi: 10.3390/life7030029
- Gull, M., Zhou, M., Fernández, F. M., and Pasek, M. A. (2014). Prebiotic phosphate ester syntheses in a deep eutectic solvent. *J. Mol. Evol.* 78, 109–117. doi: 10.1007/s00239-013-9605-9
- Hammond, O. S., Bowron, D. T., and Edler, K. J. (2017). Effect of water upon deep eutectic solvent nanostructure: an unusual transition from ionic mixture to aqueous solution. *Angew. Chem. Int. Ed.* 56, 9782–9785. doi: 10.1002/anie.201702486
- Hänle, E., and Richert, C. (2018). Enzyme-free replication with two or four bases. *Angew. Chem. Int. Ed.* 57, 8911–8915. doi: 10.1002/anie.201803074
- He, C., Gállego, I., Laughlin, B., Grover, M. A., and Hud, N. V. (2017). A viscous solvent enables information transfer from gene-length nucleic acids in a model prebiotic replication cycle. *Nat. Chem.* 9, 318–324. doi: 10.1038/nchem.2628
- He, C., Lozoya-Colinas, A., Gállego, I., Grover, M. A., and Hud, N. V. (2019). Solvent viscosity facilitates replication and ribozyme catalysis from an RNA duplex in a model prebiotic process. *Nucleic Acids Res.* 47, 6569–6577. doi: 10.1093/nar/gkz496
- He, Y., and Liu, D. R. (2010). Autonomous multistep organic synthesis in a single isothermal solution mediated by a DNA walker. *Nat. Nanotechnol.* 5, 778–782. doi: 10.1038/nnano.2010.190
- He, Y., and Liu, D. R. (2011). A sequential strand-displacement strategy enables efficient six-step DNA-templated synthesis. *J. Am. Chem. Soc.* 133, 9972–9975. doi: 10.1021/ja201361t
- Johnston, W. K., Unrau, P. J., Lawrence, M. S., Glasner, M. E., and Bartel, D. P. (2001). RNA-catalyzed RNA polymerization: accurate and general RNA-templated primer extension. *Science* 292, 1319–1325. doi: 10.1126/science.1060786
- Kaiser, R. I., Maity, S., and Jones, B. M. (2015). Synthesis of prebiotic glycerol in interstellar ices. *Angew. Chem. Int. Ed.* 54, 195–200. doi: 10.1002/anie.201408729
- Kanavarioti, A., Bernasconi, C. F., Doodokyan, D. L., and Alberas, D. J. (1989). Magnesium ion catalyzed phosphorus-nitrogen bond hydrolysis in imidazole-activated nucleotides. relevance to template-directed synthesis of polynucleotides. *J. Am. Chem. Soc.* 111, 7247–7257. doi: 10.1021/ja00200a053
- Kruger, K., Grabowski, P. J., Zaug, A. J., Sands, J., Gottschling, D. E., and Cech, T. R. (1982). Self-splicing RNA: autoexcision and autocyclization of the ribosomal RNA intervening sequence of tetrahymena. *Cell* 31, 147–157. doi: 10.1016/0092-8674(82)90414-7
- Lannan, F. M., Mamajanov, I., and Hud, N. V. (2012). Human telomere sequence DNA in water-free and high-viscosity solvents: G-quadruplex folding governed by kramers rate theory. *J. Am. Chem. Soc.* 134, 15324–15330. doi: 10.1021/ja303499m
- Li, L., Prywes, N., Tam, C. P., Oflaherty, D. K., Lelyveld, V. S., Izgu, E. C., et al. (2017). Enhanced nonenzymatic RNA copying with 2-aminoimidazole activated nucleotides. *J. Am. Chem. Soc.* 139, 1810–1813. doi: 10.1021/jacs.6b13148
- Li, X., and Liu, D. R. (2004). DNA-templated organic synthesis: nature's strategy for controlling chemical reactivity applied to synthetic molecules. *Angew. Chem. Int. Ed.* 43, 4848–4870. doi: 10.1002/anie.200400656
- Liu, Z., Mariani, A., Wu, L., Ritson, D., Folli, A., Murphy, D., et al. (2018). Tuning the reactivity of nitriles using Cu(ii) catalysis-potentially prebiotic activation of nucleotides. *Chem. Sci.* 9, 7053–7057. doi: 10.1039/C8SC02513D
- Ma, C., Laaksonen, A., Liu, C., Lu, X., and Ji, X. (2018). The peculiar effect of water on ionic liquids and deep eutectic solvents. *Chem. Soc. Rev.* 47, 8685–8720. doi: 10.1039/C8CS00325D
- Mamajanov, I., Engelhart, A. E., Bean, H. D., and Hud, N. V. (2010). DNA and RNA in anhydrous media: duplex, triplex, and G-quadruplex secondary structures in a deep eutectic solvent. *Angew. Chem. Int. Ed.* 49, 6310–6314. doi: 10.1002/anie.201001561

- Mann, S. (2012). Systems of creation: the emergence of life from nonliving matter. *Acc. Chem. Res.* 45, 2131–2141. doi: 10.1021/ar200281t
- Mariani, A., Russell, D. A., Javelle, T., and Sutherland, J. D. (2018). A light-releasable potentially prebiotic nucleotide activating agent. *J. Am. Chem. Soc.* 140, 8657–8661. doi: 10.1021/jacs.8b05189
- Maugeri, Z., Leitner, W., and Domínguez De María, P. (2013). Chymotrypsin-catalyzed peptide synthesis in deep eutectic solvents. *Eur. J. Org. Chem.* 2013, 4223–4228. doi: 10.1002/ejoc.201300448
- Meng, W., Muscat, R. A., McKee, M. L., Milnes, P. J., El-Sagheer, A. H., Bath, J., et al. (2016). An autonomous molecular assembler for programmable chemical synthesis. *Nat. Chem.* 8, 542–548. doi: 10.1038/nchem.2495
- Mondal, D., Sharma, M., Mukesh, C., Gupta, V., and Prasad, K. (2013). Improved solubility of DNA in recyclable and reusable bio-based deep eutectic solvents with long-term structural and chemical stability. *Chem. Commun.* 49, 9606–9608. doi: 10.1039/c3cc45849k
- Okamura, H., Crisp, A., Hübner, S., Becker, S., Rovo, P., and Carell, T. (2019). Proto-urea-RNA (Wöhler RNA) containing unusually stable urea nucleosides. *Angew. Chem. Int. Ed.* 58, 18691–18696. doi: 10.1002/anie.201911746
- O'Reilly, R. K., Turberfield, A. J., and Wilks, T. R. (2017). The evolution of DNA-templated synthesis as a tool for materials discovery. *Acc. Chem. Res.* 50, 2496–2509. doi: 10.1021/acs.accounts.7b00280
- Orgel, L. E. (1968). Evolution of the genetic apparatus. *J. Mol. Biol.* 38, 381–393. doi: 10.1016/0022-2836(68)90393-8
- Pätzold, M., Siebenhaller, S., Kara, S., Liese, A., Syltack, C., and Holtmann, D. (2019). Deep eutectic solvents as efficient solvents in biocatalysis. *Trends Biotechnol.* 37, 943–959. doi: 10.1016/j.tibtech.2019.03.007
- Paul, N., and Joyce, G. F. (2002). A self-replicating ligase ribozyme. *Proc. Natl. Acad. Sci. U.S.A.* 99, 12733–12740. doi: 10.1073/pnas.202471099
- Smith, E. L., Abbott, A. P., and Ryder, K. S. (2014). Deep eutectic solvents (DESSs) and their applications. *Chem. Rev.* 114, 11060–11082. doi: 10.1021/cr30162p
- Smith, P. J., Arroyo, C. B., Lopez Hernandez, F., and Goeltz, J. C. (2019). Ternary deep eutectic solvent behavior of water and urea choline chloride mixtures. *J. Phys. Chem. B* 123, 5302–5306. doi: 10.1021/acs.jpcc.8b12322
- Steitz, T. A. (2008). A structural understanding of the dynamic ribosome machine. *Nat. Rev. Mol. Cell Biol.* 9, 242–253. doi: 10.1038/nrm2352
- Szostak, J. W. (2012). The eightfold path to non-enzymatic RNA replication. *J. Syst. Chem.* 3:2. doi: 10.1186/1759-2208-3-2
- Tamura, K., and Schimmel, P. (2001). Oligonucleotide-directed peptide synthesis in a ribosome- and ribozyme-free system. *Proc. Natl. Acad. Sci. U.S.A.* 98, 1393–1397. doi: 10.1073/pnas.98.4.1393
- Tamura, K., and Schimmel, P. (2004). Chiral-selective aminoacylation of an RNA minihelix. *Science* 305:1253. doi: 10.1126/science.1099141
- Turk, R. M., Chumachenko, N. V., and Yarus, M. (2010). Multiple translational products from a five-nucleotide ribozyme. *Proc. Natl. Acad. Sci. U.S.A.* 107, 4585–4589. doi: 10.1073/pnas.0912895107
- Wagle, D. V., Zhao, H., and Baker, G. A. (2014). Deep eutectic solvents: sustainable media for nanoscale and functional materials. *Acc. Chem. Res.* 47, 2299–2308. doi: 10.1021/ar5000488
- Xu, P., Zheng, G.-W., Zong, M.-H., Li, N., and Lou, W.-Y. (2017). Recent progress on deep eutectic solvents in biocatalysis. *Bioresour. Bioprocess.* 4, 34–52. doi: 10.1186/s40643-017-0165-5
- Yarus, M. (2001). On translation by RNAs alone. *Cold Spring Harb. Symp. Quant. Biol.* 66, 207–216. doi: 10.1101/sqb.2001.66.207
- Yonath, A. (2009). Large facilities and the evolving ribosome, the cellular machine for genetic-code translation. *J. R. Soc. Interface* 6, S575–S585. doi: 10.1098/rsif.2009.0167.focus
- Zadeh, J. N., Steenberg, C. D., Bois, J. S., Wolfe, B. R., Pierce, M. B., Khan, A. R., et al. (2009). NUPACK: analysis and design of nucleic acid systems. *J. Comput. Chem.* 32, 170–173. doi: 10.1002/jcc.21596
- Zhang, B., and Cech, T. R. (1997). Peptide bond formation by *in vitro* selected ribozymes. *Nature* 390, 96–100. doi: 10.1038/36375
- Zhang, Q., De Oliveira Vigier, K., Royer, S., and Jérôme, F. (2012). Deep eutectic solvents: syntheses, properties and applications. *Chem. Soc. Rev.* 41, 7108–7146. doi: 10.1039/c2cs35178a
- Zhang, W., Pal, A., Ricardo, A., and Szostak, J. W. (2019). Template-directed nonenzymatic primer extension using 2-methylimidazole-activated morpholino derivatives of guanosine and cytidine. *J. Am. Chem. Soc.* 141, 12159–12166. doi: 10.1021/jacs.9b06453
- Zhao, C., Ren, J., and Qu, X. (2013). G-quadruplexes form ultrastable parallel structures in deep eutectic solvent. *Langmuir* 29, 1183–1191. doi: 10.1021/la3043186

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Núñez-Pertíñez and Wilks. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.