

The emotional and psychiatric problems of adolescents on parole whose parents are substance users

Perruci, Ludmila Gonçalves; Diehl, Alessandra; Da Silveira, Belisa Vieira; Teixeira, Jessica Adrielle; Souza, Jacqueline; Miasso, Adriana Inocenti; Dos Santos, Patricia Leila; Dos Santos, Manoel Antônio; De Souza, Roberto Molina; Pillon, Sandra Cristina; Wagstaff, Christopher

DOI:

[10.1177/1367493520925661](https://doi.org/10.1177/1367493520925661)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Perruci, LG, Diehl, A, Da Silveira, BV, Teixeira, JA, Souza, J, Miasso, AI, Dos Santos, PL, Dos Santos, MA, De Souza, RM, Pillon, SC & Wagstaff, C 2021, 'The emotional and psychiatric problems of adolescents on parole whose parents are substance users: a Brazilian cross-sectional study', *Journal of Child Health Care*, vol. 25, no. 2, pp. 253-267. <https://doi.org/10.1177/1367493520925661>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the Accepted Manuscript version of an article accepted for publication in *Journal of Child Health Care*.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Emotional and Psychiatric Problems of Adolescents on Parole Whose Parents are Substance Users: A Brazilian Cross-Sectional Study

Journal:	<i>Journal of Child Health Care</i>
Manuscript ID	JCHC-2019-0319.R1
Manuscript Type:	Original Article
Keyword:	Adolescent, Child Abuse, Substance-Related Disorders, Parents, Juvenile Delinquency
Abstract:	<p>The aim of this study is to evaluate the associations between having parents with substance use problems, and having suffered neglect within the family, and behavioural problems (psychological and drug use) amongst adolescents. All the participants were enrolled on the socio-educational parole scheme, 'Assisted Freedom' (AF). A cross-sectional study of 150 adolescents were interviewed using the Drug Abuse Screening Test, Teen Addiction Severity Index and Childhood Trauma Questionnaire. Ninety-five percent of the participants were male [n =143], aged 13-17. Thirty percent of adolescents had a parent who used substances and had experienced neglect from their families. Those adolescents who were living with both parents (Odds Ratio Adjusted [ORA] 2.7 95% CI 1.13;6.37), from a low income family (6.7 ORA 95%CI 1.85;24.22), experienced hallucinations (ORA 2.8 95%CI 1.25;6.14), had problems controlling violent behaviour (ORA 2.6 95%CI 1.12;5.87) and were physically neglected (3.0 ORA 95%CI 1.24;7.49) were more likely to have parents who used substances and to have experienced parental neglect. This paper concludes that adolescents, who are on parole, come from families with high level of psychosocial vulnerabilities, including substance use, experience neglect by their families leading to adverse emotional and psychological states.</p>

SCHOLARONE™
Manuscripts

1
2
3 **The Emotional and Psychiatric Problems of Adolescents on Parole Whose Parents**
4 **are Substance Users: A Brazilian Cross-Sectional Study**
5
6
7
8
9

10 **ABSTRACT**
11

12 The aim of this study is to evaluate the associations between having parents with
13 substance use problems, and having suffered neglect within the family, and behavioural
14 problems (psychological and drug use) amongst adolescents. All the participants were
15 enrolled on the socio-educational parole scheme, 'Assisted Freedom' (AF). A cross-
16 sectional study of 150 adolescents were interviewed using the Drug Abuse Screening
17 Test, Teen Addiction Severity Index and Childhood Trauma Questionnaire. Ninety-five
18 percent of the participants were male [n =143], aged 13-17. Thirty percent of adolescents
19 had a parent who used substances and had experienced neglect from their families. Those
20 adolescents who were living with both parents (Odds Ratio Adjusted [ORA] 2.7 95% CI
21 1.13;6.37), from a low income family (6.7 ORA 95%CI 1.85;24.22), experienced
22 hallucinations (ORA 2.8 95%CI 1.25;6.14), had problems controlling violent behaviour
23 (ORA 2.6 95%CI 1.12;5.87) and were physically neglected (3.0 ORA 95%CI 1.24;7.49)
24 were more likely to have parents who used substances and to have experienced parental
25 neglect. This paper concludes that adolescents, who are on parole, come from families
26 with high level of psychosocial vulnerabilities, including substance use, experience
27 neglect by their families leading to adverse emotional and psychological states.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 **Key words:** Adolescent, Child Abuse, Child Maltreatment, Substance-Related Disorders,
53 Parenting, Juvenile Delinquency
54
55
56
57
58
59
60

Introduction

Substance use is a social and public health problem in many countries; with significantly contribution to the increase in the global burden of diseases and diverse consequences for the individuals, for families and society (Kepple, 2017; Global Burden Disease, 2018). The impact of alcohol use and other substances by parents on the physical and psychological health of their children has been well documented in the literature, including Brazilian studies (Brazilian National Alcohol and Drugs Survey [BNADS], 2013; Institute of Medicine [IOM] and National Research Council [NRC], 2014).

It is estimated that about 2.1 million children in the United States of America (USA) may be suffering as a consequence of substance use within their families, and 2.9% of children under 18 live in households with at least one parent who is a substance user (IOM and NRC, 2014). Brazilian data confirm that social vulnerabilities in households impact on both individuals and the whole family nucleus; for every substance user there are four other people living with the problem in their homes (BNADS, 2013). Half of the Brazilian population consumes alcohol and the problems related to abuse of alcohol are very common. Substance use prevalence in the last 12 months and in life by the Brazilian population (age ≥ 18 years old) is considerably: tranquilizers (6.0 and 9.6%), marijuana (2.5% and 6.8%), cocaine (1.7% and 3.8%), crack (0.7% and 1.3%), respectively (BNADS, 2013).

Substance use by parents has been associated with abuse, mistreatment and family neglect (Solis et al., 2012;) resulting in physical injury and the development of psychological and psychiatric distress; including, sleep disorders, depression, anxiety, post-traumatic stress disorder, and suicidal behavior amongst their children (Leeb et al., 2011; WHO, 2014). Given the environment that the children of parents who use substances grow up in, there is an increased risk of experiencing various forms of violence

1
2
3 (mainly domestic violence) and family dysfunctional, as well, as precarious access to
4
5 basic necessities such as food and clothing (Horgan, 2011; Winters, 2014).
6

7
8 The destitution of family cohesion, the reversal of roles models in the family with
9
10 children taking responsibility and care of parents and other siblings (Solis et al., 2012)
11
12 are also consequences of parental substance use. Additionally, parental substance use
13
14 (PSU) may threaten the realization of children's potential by exposing them to stressful,
15
16 chaotic and often frightening pressures in their homes. Child maltreatment (CM)
17
18 contributes both directly and indirectly to various health and welfare damages, affecting
19
20 the development of children and adolescents, resulting in low performance, repetition and
21
22 school dropout (Berg et al., 2016). PSU use is an important risk factor for the development
23
24 of psychological problems (shyness, impulsivity, low self-esteem, low problem solving
25
26 ability, high levels of pessimism, fear, guilt, relationship difficulties, loss of self-
27
28 confidence and insecurity) (Berg et al., 2016). Additionally, there is an increased chance
29
30 of mental disorders (e.g. depression, anxiety, conduct disorder, social phobia and suicide),
31
32 risk behaviors (teenage pregnancy and sexually transmitted infections) and violence,
33
34 continuing into adulthood (Elgán et al., 2016).
35
36
37
38
39

40 Furthermore, there is an increased risk of substance use, with children who grow
41
42 up in these environments showing higher rates of alcohol and tobacco use when they
43
44 become adults (WHO, 2014; Meyers et al., 2018). International studies have reported
45
46 associations between risky alcohol consumption and childhood abuse (WHO, 2014). Only
47
48 four out of every ten countries carry out research into CM and 60% of these studies come
49
50 from high-income European countries (WHO, 2014).
51
52

53 From an economic perspective, whilst the costs of child abuse are substantial (with
54
55 an increase burden on the judiciary, health and social services) the costs of substance use-
56
57 related mistreatment remain largely under investigated. Due to the different
58
59
60

1
2
3 methodologies used in studies and variety of countries, it is difficult to make comparisons
4
5 between results. Additionally, there is frequently bias in studies due to underreporting of
6
7 CM to the health sector and /or to the judiciary (WHO, 2014).
8
9

10 In the USA, for example, 35% of individuals who committed offenses, and who
11
12 suffered parental abuse, had consumed alcohol or drugs at the time of the incident (WHO,
13
14 2014). Adolescents who have suffered maltreatment are more likely to engage in
15
16 vandalism, crimes against property, physical fights, bearing or threat with weapons. The
17
18 triad of adolescence, delinquency, and substance use are issues of crucial importance as
19
20 they may be predictive factors for a chronic pattern of violent offending into adulthood
21
22 (Leeb et al., 2011; Cicchetti and Handley, 2019).
23
24
25

26 Adolescent offending can be viewed as a dramatic result of the difficulties that
27
28 families face. The association between CM, criminal activity, family neglect and PSU are
29
30 important social and public health problems and have implications for health and social
31
32 services, as well as, the judiciary. Nevertheless, these associations are underexplored in
33
34 the literature, especially in low-income countries such as Brazil (WHO, 2014).
35
36
37

38 Understanding the contexts of parental substance use is a requirement in the public
39
40 policies of some countries (Velleman and Templeton, 2016) as opposed to how children
41
42 and adolescents affected by parental consumption who have become involved in
43
44 criminality and subsequently with social services or judiciary services. Currently there is
45
46 health (Torvik et al., 2011; IOM and NRC, 2014; Diehl et al., 2018), community service
47
48 (Figlie et al., 2004), schools (Filov et al., 2014) and general population (WHO, 2014)
49
50 related research on the subject. There has been limited published literature investigating
51
52 the outcomes for adolescents, who come from families with a substance using parents
53
54 (Winters, 2014; Nadel and Thornberry, 2017), who have been involved the judiciary,
55
56 hence this present study.
57
58
59
60

Aim

The aim of this study is to evaluate the associations between having a parent who used alcohol/ drugs problematically, having experienced some neglect or abuse from their parents and the emotional/ behavioral problems (for example, psychological and drug use) experienced by adolescents enrolled on a Brazilian parole scheme.

Methods

A cross-sectional study was carried out with a sample of 150 (62.7%) adolescents from a total population of 239 (100%) who, in 2018, were enrolled in a Municipality in the Interior 'Assisted Freedom'(AF), socio educational parole scheme in the state of São Paulo, Brazil. AF is a measure provided for by Law 8069/1990 in Brazil for offenders who have committed an offence but are yet to reach the age of majority, allowing them to remain in the community to serve his or her sentence under the supervision of a social worker and an authority designated by the judiciary (Ortegal, 2011). Therefore, it is a non-custodial socio-educational sentence imposed by the courts aimed at assisting young people back into some form of stable living environment. AF aims to address the adolescent's attitudes and develop important family community values. The work is done through educational interventions focused on personalized service, guaranteeing the social promotion of the adolescents through orientation, maintaining family and community ties, schooling, introduction into the labor market and/ or vocational and training courses (Ortegal, 2011).

Participants for this study were adolescent (aged between 12 and 18 years old incomplete) and enrolled in AF, regardless of offence. Participants who were unable to give consent and/ or cognitive impairment (evaluated through a clinical interview) or

1
2
3 were under the influence of alcohol and/ or drugs at the time of the interview were
4
5 excluded from the study. Formal authorization was requested from Judges from the
6
7 relevant municipality and the coordinators of the Non-Governmental Organizations
8
9 (NGOs), in order to carry out data collection. Data collection was from October 2016 to
10
11 April 2017. The interviews were conducted individually and in private rooms. Each
12
13 interview lasted up to one and a half hours. The interviews were conducted by an
14
15 experienced researcher, trained in the application of the instruments. A pilot test had been
16
17 previously conducted with 10 adolescents who met the socio-educational measures,
18
19 which allowed standardizing the instruments for the researched population.
20
21
22
23
24
25

26 **Instruments**

27
28 - **Socio-demographic data:** age, sex, school attendance, religion, race,
29
30 composition of family, how the family income is generated were all noted.
31

32
33 - **Childhood Trauma Questionnaire (CTQ):** a retrospective self-report
34
35 instrument that measures the various forms of emotional, physical, and sexual abuse,
36
37 emotional and physical neglect in adolescence (age 12 years-old) and adults (Grassi-
38
39 Oliveira et al., 2014). The Brazilian version of CTQ consists of 28 statements related to
40
41 situations that may have occurred, and frequency, in childhood using a five-point Likert
42
43 scale. The score ranges from 28 to 140 points. A study that evaluated the internal structure
44
45 of the questionnaire, composed of five factors (abuse: sexual / physical and negligence:
46
47 physical / emotional), demonstrated that the instrument had good internal consistency in
48
49 Brazilian Portuguese (Grassi-Oliveira et al., 2014). In addition, the reliability of CTQ was
50
51 analyzed by test-retest and found to be stable. In the present study, the dependent variable
52
53 was used, *"I have alcoholic parents or drug users and experienced neglect in family*
54
55 *care"*.
56
57
58
59
60

1
2
3 We provide further evidence for the validity and reliability of the CTQ within the
4
5 Brazilian samples 1,925 participants from eight different clinical and non-clinical
6
7 samples including adolescents, adults and elders. In this study the alternative five-factor
8
9 solution Cronbach's alpha revealed acceptable levels of internal consistency (Emotional
10
11 Abuse 0.80, Physical Abuse = 0.80, Sexual Abuse = 0.90, Emotional Negligence 0.91)
12
13 (Grassi-Oliveira et al., 2014).
14
15

16
17 - **Drug Abuse Screening Test (DAST-20):** is composed of 20 items that assess
18
19 abuse, dependence, abstinence (signs and symptoms), social harm, family relationships,
20
21 legal implications, medical problems and previous treatment. The DAST-20 can be
22
23 applied as self-reporting or in interview format (Yudko et al., 2007). In the present study,
24
25 a Brazilian version were used, with good reliability (DAST-20) have a Cronbach's alpha
26
27 (α) range from 0.82–0.96 (Diehl et al., 2014; Diehl et al., 2016) in the versions, either for
28
29 clinical use (Cronbach's $\alpha = 0.74$) and / or as a research tool (Cronbach's $\alpha = 0.96$)
30
31 (Yudko et al., 2007). This classification used was recoded at zero - five points = no
32
33 problems and six or more points (medium to severe problems) (Diehl et al., 2014; Diehl
34
35 et al., 2016).
36
37
38
39

40 - **Teen Addiction Severity Index (T-ASI):** aims to assess the severity of drug
41
42 use, abuse or dependence in adolescents as well as other problems experienced in their
43
44 lives. The Brazilian version of T-ASI is a semi-structured interview consists of 153 items
45
46 with dichotomous responses, divided into seven domains: 1) substances use; 2) school
47
48 situation; 3) employment / livelihood; 4) family relationships; 5) friends / social
49
50 relationships 6) legal status: with a direct question about “whether you have ever been
51
52 caught by the police in possession of drugs (cannabis, cocaine and/or crack) to traffic
53
54 dealing” and, 7) psychiatric status (Sartes et al., 2009). In the present study, two domains
55
56 were used: substance use of alcohol, tobacco, cannabis, inhalants in the last 30 days and
57
58
59
60

1
2
3 psychiatric status (Depression, Severe anxiety or tension, Hallucinations, Problems
4 controlling violent behaviour and problems with cognition, concentration or memory).
5
6 This scale has been validated for use in Brazil and presented good index of Cronbach's
7
8 alpha in areas: substance use ($\alpha = 0.89$), legal ($\alpha = 0.81$), and psychiatric ($\alpha = 0.80$) (Sartes
9
10 et al., 2009).
11
12
13
14
15
16

17 **Ethics**

18
19 Ethical approval was obtained from the University of São Paulo (USP) (protocol
20 number 1672/04), followed the ethical presuppositions recommended by Brazilian
21 Resolution 466/2012 for research with human beings. All the participants signed an assent
22 forms at the start of the interview. As the participants were adolescents, their legal
23 guardians signed the assent terms in accordance with The Free, Prior and Informed
24 Consent (FPIC) process. The participants did not receive any funds or compensation for
25 participating in this study.
26
27
28
29
30
31
32
33
34

35 **Statistical analysis**

36
37 The data were analyzed using Stata version 12 software. Statistical analyses
38 included the Chi-square test (χ^2) and Degrees of freedom (df) for all samples in the
39 bivariate analysis. Socio-demographic characteristics, substance use (T-ASI and DAST-
40 20), maltreatment (CTQ) and, mental health problems (ASI-Teen) were described by PSU
41 and neglect of family care and the values with $p < 0.05$ were considered statistically
42 significant. Logistic regression analysis was performed. Adjusted odds ratios (ORA) with
43 95% confidence intervals (CI) were calculated for the associations between co-variables:
44 information socio-demographic, substance use (T-ASI), type of abuse associated with
45 maltreatment (CTQ) experienced by adolescents and, symptoms and mental health
46 problems (T-ASI) and the dependent variable (PSU and neglect of family care). However,
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 variables such as tobacco and emotional abuse were not included in the multivariate
4
5 analysis, but were considered when significant compared to the non-adjusted OR because
6
7 we seek a parsimonious model, a model using the fewest possible variables. The
8
9 significance level of 5% was used for all statistical tests.
10
11
12
13

14 **Results**

15 **Sociodemographic Profile**

16
17 The participants were male 143 (95.3%), 17 – 18 years old, black 112 (74.6%),
18
19 half of the sample 83 (55.3%) did not profess any religion and a third came from low-
20
21 income families 50 (33.3%). The majority of participants 116 (77.3%) did not live with
22
23 their parents; those living with their parents had families consisting of an average three
24
25 or four people 63 (42.0%) and five or more people 73 (48.7%), and 73 (48.7%) of the
26
27 families were financially supported by robberies, theft or drug trafficking. Living with
28
29 parents ($\chi^2(1) = 5.13$; $p = 0.023$), and financially supported by robberies, theft or drug
30
31 trafficking ($\chi^2(1) = 4.81$; $p = 0.028$) having low family income ($\chi^2(3) = 10.35$; $p = 0.015$)
32
33 were variables associated with belonging to a family with parents used substances and
34
35 neglected family care with values statistically significant in the bivariate and multiple
36
37 analysis (See Table 1).
38
39
40
41
42
43
44

45 As for the parents' educational level, half of the adolescents had mothers with an
46
47 average level of schooling, 51 (34%) did not know how to respond, 13 (8.7%) had primary
48
49 education and only two (1.3%) had a higher education level. Regarding the father,
50
51 although 89 (59.3%) did not know how to respond, only one third had high school, 11
52
53 (7.3%) had primary education and two (1.3%) had higher education. Parents' schooling
54
55 did not present a statistically significant value when compared to the dependent variable
56
57 (Data were not presented in the table).
58
59
60

1
2
3 Amongst participants, 100 (65.3%) were detained for being in possession of drugs
4 (cannabis, cocaine and/or crack) to traffic dealing and were complying with the AF
5 scheme. Only 27 (27.0%) of the adolescents involved in trafficking had a substance using
6 parent and experienced neglect in family care, with no statistically significant differences
7 in the sample ($\chi^2(1) = 0.408$; $p = 0.523$) (data were not presented in the table). In the
8 multiple analysis, the odds ratio shows the contribution of each variable to the increase
9 in probability. Therefore, the adolescents living with both parents in families with 1-2
10 the minimum wage (MW) were at higher risk (Odds Ratio Adjusted = 6.7 CI 95% = 1.85;
11 24.22) of having alcohol-dependent parents and/ or drug users who neglected their family
12 members.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 [Insert Table 1]
27
28
29
30

31 **Family Neglect and Substance Use**

32
33 A quarter of participants reported that they had family problems, of which almost
34 half came from families whose parents had problems related to substance use and
35 neglecting family care (Yes 42.1% versus No 24.1%; $\chi^2 4.495 p = 0.034$), data not shown
36 in the table.
37
38
39
40
41

42 Of the total sample, more than half of the adolescents reported having experienced
43 physical 139 (92.7%) and emotional 135 (90%) neglect; physical 88 (58.7%), emotional
44 90 (60%) and sexual six (4%) abuse (CTQ). Emotional abuse was associated with having
45 parents who used substances and had neglected family care [36 (40%) yes versus 07
46 (11.7%) no, $\chi^2(1) = 14,133$; p value <0.001 , non-adjusted OR = 5.0; CI 95% 2.06-12.34].
47
48
49
50
51
52
53
54 Data presented in a complementary table CI.
55

56 Sexual abuse was a factor in six cases, mostly amongst participants whose parents
57 who used substances and had neglected family care [4 (66.7%) Yes versus 2 (33.3%) No,
58
59
60

1
2
3 *p* value =0.036]. The results showed associations only in the bivariate analysis, between
4 the abuse (sexual and emotional aggressions) and parents with substance use and neglect
5 in family care. Noted that the risk of physical neglect in relation to personal care and
6 hygiene (wearing dirty clothing) risks are tripled (ORA = 3.0 95% CI 1.24-7.49, *p* = 0.02)
7 amongst the participants (Table CII complementary).
8
9
10
11
12
13
14
15
16

17 **Substance use by participants**

18
19 Regarding substance use, participants had a higher prevalence of marijuana use
20 86 (57.3%), alcohol 59 (39.3%) and tobacco 47 (33.5%) in the last 30 days.
21
22

23
24 Additionally, 19 (40.4%) of the participants who had used substances and had
25 been neglected by their family were smokers versus 24 (23.3%) non-smokers ($\chi^2(1) =$
26 4,628; *p* = 0.031) and non-adjusted odds ratio OR 2.2 95% CI 1.07-4.68 *p* = 0.031, (See
27 Table CIII complementary). Of the total sample, 63 (42.0%) were classified with
28 moderate substance use problems (DAST). In the bivariate analysis, no differences were
29 observed between having alcoholic parents who neglected the care and problem levels
30 of DAST-20 (22 (34.9%) moderate versus 22 (24.1%) in the problem; $\chi^2(1) = 2,078$ *p*
31 = 0.149) (See Table 3 complementary table III).
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Symptoms and mental health problems amongst participants**

46
47 All symptoms and mental health problems were significantly associated with have
48 substance using parents who had shown neglect in family care. A predominance of those
49 participants whose parents used substances and had neglected family care, had depressive
50 symptoms [14 (42.4%) yes versus 29 (24.8%) no $\chi^2(1) = 3,916$; *p* = 0.048], experienced
51 hallucinations [17 (46.0%) yes versus 26 (23.0%) no; $\chi^2(1) = 7,171$; *p* = 0.007] and had
52 problems controlling their violent behavior [15 (45.5%) yes versus 28 (23.9%) no; $\chi^2(1)$
53
54
55
56
57
58
59
60

1
2
3 = 5,831 $p= 0.016$]. Moreover, in the multivariate analysis, those participants were almost
4
5 three times more likely to have depressive symptoms (ORA 2.8 CI95% 1.25-6.14) and
6
7 had problems controlling their violent behavior (ORA 2.6 CI 95% 1.12-5.87).
8
9

10
11
12 [*Insert Table 2*]
13
14
15

16 **Discussion**

17
18
19 Of the total sample, 30% of the participants had a parent who both used substances
20
21 and had neglected family care; the majority of participants were non-black male
22
23 adolescents with a low level of education attainment; from single-parent families with
24
25 low levels of schooling and family income (Table 1). These characteristics could be
26
27 explained, or justified, as the participants having experienced neglect by parents and
28
29 living in low socioeconomic circumstances leading to searching for sustenance through
30
31 the money retrieved from illicit trafficking, robbery and thefts. The profile highlights a
32
33 group of adolescents with strong environmental and psychosocial risk factors, making
34
35 them more susceptible to emotional and psychological problems after being exposed to
36
37 multiple vulnerabilities, including substance use and neglect in their family care.
38
39
40

41
42 Likewise, amongst participants who lived in families with an income between one
43
44 to two times the MW, had a greater potential to experience negligence by parents who
45
46 use substances. There is evidence pointing to strong associations between poverty and
47
48 CM in many countries (WHO, 2014; Walsh et al., 2019; Goldberg and Blaauw, 2019).
49
50 Evidence also suggests that the direct and indirect impacts of poverty interact in complex
51
52 ways with other factors that affect parenting and increase the risk of abuse and neglect
53
54 (WHO, 2014; Bywaters et al., 2016). In addition, significant rates of social inequalities
55
56 related to deprivation were considered as one of the major risk factors for human
57
58 development often identified in families receiving support with parenting (Cambron et
59
60

1
2
3 al., 2019). Studies have shown that socially excluded families are vulnerable to problems
4 related to substance related disorders (WHO, 2014; Aldridge et al., 2018). Children who
5
6 related to substance related disorders (WHO, 2014; Aldridge et al., 2018). Children who
7
8 have parents vulnerable to substance related disorders are at higher risk for developing
9
10 their own health problems and psychosocial difficulties (Dyba et al., 2019).
11

12 Although there are no statistically significant differences were found in the school
13 setting, nonetheless half of the participants had dropped out of education (Table 1).
14
15 Literature shows that both maltreatment and substance use by parents, regardless of
16
17 socioeconomic backgrounds and other psychosocial factors, significantly influences
18
19 school performance (Torvik et al., 2011; IOM and NRC, 2014; Gauffin et al., 2015).
20
21 Academic performance has been strongly evaluated in the most recent studies with
22
23 different results. Whilst poverty and deprivation do negatively influence upon education
24
25 (Torvik et al., 2011; IOM and NRC, 2014; WHO, 2014; Gauffin et al., 2015), children
26
27 who have experienced maltreatment do not necessarily under perform at school relative
28
29 to the general population.
30
31
32
33

34
35 Nonetheless, some literature reviews have shown strong links between CM and
36
37 low academic performance (Torvik et al., 2011; IOM and NRC, 2014; Gauffin, et al.,
38
39 2015, Berg et al., 2016). The main problem identified is the higher risk of being
40
41 impoverished at school and being bullied, more likely to need special education and
42
43 greater risk of absence, repetition and dropping out of school (Torvik et al., 2011; IOM
44
45 and NRC, 2014; WHO, 2014; Gauffin et al., 2015; Berg et al., 2016). The same has been
46
47 observed in children whose parents use alcohol problematically who, when compared to
48
49 children of non-problematic drinking parents, are found to have a poorer academic
50
51 performance (Torvik et al., 2011). Low academic achievement is associated with
52
53 delinquency and school failure (IOM and NRC, 2014; WHO, 2014). Students with low
54
55 grades are more likely to be involved in violence and other behavior problems such as
56
57
58
59
60

1
2
3 non-durable links to school and school dropout which in turn potentiate the risk of
4
5 involvement with violence (IOM and NRC, 2014; WHO, 2014).
6
7

8 All forms of parental abuse and neglect were evidenced in the bivariate analysis
9
10 among the participants of the present study (Table CI complementary). Physical and
11
12 emotional neglect were the most prevalent forms of maltreatment. Adolescents who had
13
14 alcohol-dependent parents and neglected family care had experienced emotional abuse
15
16 when compared to other adolescents. Among the main types of emotional violence, verbal
17
18 abuse that generates feelings of rejection on the part of the adolescent are highlighted.
19
20 Nonetheless, this finding is contradicted by the finding that the participants do not want
21
22 to change anything in their family (see table CII complementary). On the other hand,
23
24 adolescents who experienced physical neglect in the form of "hygiene care" were three
25
26 times more likely to have alcohol dependent parents.
27
28
29

30 According to the World Report on Violence Prevention (WHO, 2014) children
31
32 who experience rejection, neglect, severe corporal punishment and sexual abuse - or who
33
34 witness violence at home or in the community - are at greater risk of being involved in
35
36 aggressive behavior and antisocial behavior in later stages of development, including
37
38 violent behaviors in adulthood (WHO, 2014). This underscores the preponderant role that
39
40 substance use plays in all forms of abuse, mistreatment and family neglect.
41
42
43

44 Physical neglect and abuse (See table CI and CII complementary) are severe and
45
46 widespread forms of maltreatment that occur in childhood and adolescence, with potential
47
48 long-term consequences and adverse repercussions throughout life. Abuse rates are
49
50 higher in communities with high levels of unemployment and concentration of poverty
51
52 (Collins, 2016; Goldberg and Blaauw, 2019). Neglect is closely associated with low
53
54 schooling and low income, but it is important to identify the best way to distinguish
55
56 parental neglect from poverty deprivation (WHO, 2014).
57
58
59
60

1
2
3 As noted in the literature, CM contributes to a wide range of adverse physical and
4 mental health outcomes that will affect both the life of the victimized child and society as
5 a whole. A history of child sexual abuse significantly increases the risk of chronic
6 depression in adults (Nelson et al., 2017) and exposure to sexual abuse doubles the
7 chances of developing anxiety and substance use disorders (SUD) (Teicher and Samson,
8 2013).

9
10
11
12
13
14
15
16
17 Mothers with SUD have also been considered as potentially at risk of perpetrating
18 child abuse and neglect (Goldberg and Blaauw, 2019). A significant number (54.2%)
19 mothers living in precarious social and economic conditions experience mental illness,
20 41.9% had a history of criminal involvement and 64.5% had been both physically and
21 sexually abused as children, with 55.4% reporting childhood sexual abuse (Taplin and
22 Mattick, 2011).

23
24
25
26
27
28
29
30
31 In this study, there was considerable marijuana use (57.3%), alcohol (39.3%),
32 tobacco use (31.3%) and moderate problems evaluated by DAST amongst the participants
33 (See Table CIII complementary). Only tobacco was differentiated between the group and
34 adolescents in the sample, it is still noticeable the use of marijuana was highly probable
35 amongst adolescents whose parents who used substances and had neglected family care
36 (OR unadjusted = 2.2) (Table CIII complementary).

37
38
39
40
41
42
43
44
45 There is evidence that the substance use by adolescents are explained both being
46 in a transition phase (adolescence), and also by social economic conditions, family
47 structure and also by the inextricable link between drugs and involvement in criminality
48 (Diehl et al., 2016; Martins and Pillon, 2008). Evidence for this can be seen in that 65.3%
49 of the participants who had been detained for carrying illicit drugs and trafficking and
50 27% of those had been involved in trafficking had a parent who used substances and had
51 neglected family care (Data not shown in table). A cross-sectional study conducted with
52
53
54
55
56
57
58
59
60

1
2
3 more than 3,000 young adults, children of alcohol dependents, were almost twice as likely
4 to be dependent on alcohol or cannabis compared to individuals without a family history
5 of alcohol dependence (Melchior et al., 2011). Marijuana users are more likely to engage
6 in risky behavior once it is perceived as advantageous to achieve a goal (Rogosch, Oshri
7 and Cicchetti, 2010). It is hypothesized that the onset and frequency of use are associated
8 with the absence of conflict between a behavior and a goal, and adolescents identified the
9 use as a positive reinforcement to achieve a goal or solve a problem (Branje, 2018). As
10 adolescents engage in maladaptive behavior, there is a high probability of engaging in
11 other types of problematic behaviors, such as substance use, which are strongly
12 corroborated the juvenile criminality and delinquency (Martins and Pillon, 2008).
13
14
15
16
17
18
19
20
21
22
23
24
25

26 This study examined the symptoms and mental health problems presented by
27 adolescents on parole. In bivariate analysis, anxiety and comprehension, concentration or
28 memory problems were the most prevalent problems and hallucinations almost triple
29 amongst adolescents whose parents who used substances and had neglected family care
30 (Table 2). These findings are corroborated by Gullbrå et al (2016) and Ferreira et al
31 (2018); children of substance using parents are at increased risk of developing emotional,
32 behavioral, and psychosocial problems. Compared with their peers, children with parents
33 who use substances have increased rates of anxiety, depression, oppositional behavior,
34 conduct and aggressive behavior, delinquency, social incongruity and somatic problems,
35 generalized stress and depressed mood, as well as lower levels of self-esteem and social
36 competence.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 **Limitations**

55
56 The results should be interpreted with caution. First, any generalization of the
57 results is limited to young people who are serving socio-educational sentences and who
58
59
60

1
2
3 live in urban environments similar to our sample. Second, the cross-sectional design of
4 the study does not support causal conclusions. Third, the authenticity of the data is at the
5 mercy of human bias, since they were based on the reports of the subjects investigated
6 (sub or overestimated). Another limitation of this study is that parental alcohol use may
7 probably be underestimated, as has been seen in other studies in which the diagnostic
8 criterion is measured by self-reporting, due to "social desirability bias" (Maloney et al.,
9 2010). The information here may also be underestimated as the professionals
10 accompanying the adolescents were collecting information prior to reporting to the judge
11 regarding improvements in the adolescent's health status, behavior and living conditions.
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 **Implications for public health**

27
28 CM is a global problem, with a serious impact on the physical and mental health
29 of the victim's development and well-being throughout the life, extending throughout
30 society; the inherent cultural and social barriers mean the issue is rarely the subject of
31 global debate (WHO, 2014; Ferrara et al., 2016). Preventing CM has not been a political
32 priority in many countries around the world, despite the scale of the problem and growing
33 awareness of the high social costs (WHO, 2014).
34
35
36
37
38
39
40
41

42 The lack of investment in policy is exacerbated by a lack of understanding among
43 the general population about the serious impacts of maltreatment throughout life on health
44 and behaviors, the burden on society and the cost implication for health and social
45 services. As recent studies have shown, maltreatment and other adversities in childhood
46 and adolescence are associated with a broad spectrum of health and mental health
47 behaviors (Goldberg and Blaauw, 2019; Leeb et al., 2011).
48
49
50
51
52
53
54
55

56 CM can be avoidable. Countries have a responsibility to implement preventive
57 measures to reduce the occurrence of this phenomenon. They can provide protection,
58
59
60

1
2
3 access to justice and care for children and adolescents who may be at risk of maltreatment,
4 including implementing psychosocial programs aimed at children, adolescents, parents
5 and caregivers, as well as creating means to identify, treat and follow known cases of
6 maltreatment (Hughes et al., 2017; Kubik, Docherty and Boxer, 2019;).
7
8
9
10
11

12 The identification of adolescents who during childhood were affected by
13 substance use by parents remains a public health challenge, since problematic drinking
14 often remains hidden in the intimacy of many families (Harwin et al., 2014).
15
16
17
18
19

20 21 **Implications for clinical practice**

22 Screening, traditionally, is the means by which a health problem is identified
23 before signs and symptoms appear. In the case of child and adolescent abuse and neglect,
24 screening presents problems, since it would be necessary to rely on information obtained
25 directly from the offender or from observers. Health professionals have a key role to play
26 in the identification, treatment and referral of cases of abuse and neglect and in reporting
27 suspected cases to relevant authorities. It is vital that cases of child and adolescent abuse
28 are detected early and for the necessary services to intervene as soon as possible in order
29 to minimize the consequences for the child/adolescents.
30
31
32
33
34
35
36
37
38
39
40
41

42 Importantly, despite the notable association between parental abuse and negative
43 outcomes for children, the alcohol abuse of parents does not explain solely why these
44 children/adolescents are at risk. Co-occurrence of risk factors including other forms of
45 parent psychopathology and family adversity, including poverty, conflict, family
46 (dis)functioning, sometimes better explain the potential risks of these adolescents
47 experiencing substance abuse and neglect histories of parents (Solis et al., 2012).
48
49
50
51
52
53
54
55
56
57

58 **Conclusion**

59
60

1
2
3 Links between substance use and family neglect by parents and child abuse has
4 previously been established. This paper demonstrates that adolescents, who are on parole,
5 come from families with high level of psychosocial vulnerabilities, including substance
6 use, experience neglect by their families leading to adverse emotional and psychological
7 states. The findings of the study have implications for the judiciary, health and social
8 services.
9
10
11
12
13
14
15
16
17
18
19

20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

- Aldridge, R. W. et al. (2018) 'Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: a systematic review and meta-analysis', *Lancet*, 391(10117), pp. 241-250. doi: 10.1016/S0140-6736(17)31869-X.
- Berg, L. et al. (2016) 'Parental alcohol-related disorders and school performance in 16-year-olds-a Swedish national cohort study', *Addiction*, (Abingdon, England). doi: 10.1111/add.13454.
- Branje, S. (2018) 'Development of parent-adolescent relationships: conflict interactions as a mechanism of change'. *Child Development Perspectives*, 12(3), pp. 171-176. doi: 10.1111/cdep.12278
- Brazilian National Alcohol and Drugs Survey (BNADS). [Levantamento Nacional de Famílias dos Dependentes Químicos. (LENAD)]. (2013) INPAD – UNIFESP. Retrieved from: <http://inpad.org.br/divulgacao-dos-dados-do-levantamento-nacional-com-familiares-dos-dependentes-quimicos/>
- Bywaters, P. et al. (2016). *The relationship between poverty, child abuse and neglect: An evidence review*. York: Joseph Rowntree Foundation.
- CDC (2014) WISQARS (Web-Based Injury Statistics Query and Reporting System), *Choice Reviews Online*. doi: 10.5860/choice.48-4227.
- Cambron, C. et al. (2019) 'Neighborhood structural factors and proximal risk for youth substance use', *Prevention Science*, doi: 10.1007/s11121-019-01072-8.
- Cicchetti, D., and Handley E. D. (2019) 'Child maltreatment and the development of substance use and disorder', *Neurobiology of Stress*, 10, doi:100144 doi.org/10.1016/j.ynstr.2018.100144
- Collins, S. E. (2016) 'Associations between socioeconomic factors and alcohol outcomes', *Alcohol Research: Current Reviews*, 38(1), pp.83-94.
- Diehl, A. et al. (2018) 'Early childhood maltreatment experience and later sexual behavior in Brazilian adults undergoing treatment for substance dependence', *Revista brasileira de psiquiatria (Sao Paulo, Brazil : 1999)*, 41(3), pp. 199-207. doi: 10.1590/1516-4446-2017-0020.
- Diehl, A. et al. (2014) 'Sexual risk behaviors in non-injecting substance-dependent Brazilian patients', *Adicciones*, 26(3), pp.208-220.
- Diehl, A. et al. (2016) 'Criminality and sexual behaviours in substance dependents seeking treatment', *Journal of Psychoactive Drugs*, pp.124-134. doi: 10.1080/02791072.2016.1168534.

- 1
2
3 Dyba, J. et al. (2019) 'Methamphetamine use in German families: Parental
4 substance use, parent-child interaction and risks for children involved', *Substance*
5 *Use and Misuse*, 54(4), pp.583–591. doi: 10.1080/10826084.2018.1528459.
- 6 Elgán, T. H, et al. (2016) 'A web-based group course intervention for 15-25-year-
7 olds whose parents have substance use problems or mental illness: study protocol
8 for a randomized controlled trial', *BMC Public Health*, 16(1), 1011.
- 9 Ferrara, P. et al. (2016) 'Child abuse and neglect and its psycho-physical and social
10 consequences: A review of the literature,' *Current Pediatric Reviews*, 12(4), pp.
11 301–310. doi: 10.2174/1573396312666160914193357.
- 12 Ferreira A. H. et al. (2018) Repercussion of parents' use of alcohol on children and
13 adolescents. *Revista Eletrônica Acervo Saúde/ Electronic Journal Collection Health*.
14 https://doi.org/10.25248/REAS168_2018
- 15 Figlie, N. et al. (2004) 'Children of addicted parents with bio-psychosocial risk
16 factors: Do they need a special care?', *Revista de Psiquiatria Clinica*, 31(2), pp. 53–
17 62. doi: 10.1590/S0101-60832004000200001.
- 18 Filov, I. et al. (2014) 'Relationship between child maltreatment and alcohol abuse -
19 findings from adverse childhood experience study in Republic of Macedonia', *Open*
20 *Access Macedonian Journal of Medical Sciences*, 2(2), pp. 390–394. doi:
21 10.3889/oamjms.2014.067.
- 22 Gauffin, K., Vinnerljung, B. and Hjern, A. (2015) 'School performance and alcohol-
23 related disorders in early adulthood: A Swedish national cohort study', *International*
24 *Journal of Epidemiology*, 44(3), pp. 919–927. doi: 10.1093/ije/dyv006.
- 25 GBD 2017 Causes of Death Collaborators. (2018) 'Global, regional, and national
26 age-sex-specific mortality for 282 causes of death in 195 countries and territories,
27 1980-2017: A systematic analysis for the Global Burden of Disease Study 2017'.
28 *Lancet*, 392(10159), pp. 1736–1788. doi: 10.1016/S0140-6736(18)32203-7.
- 29 Goldberg, A. E. and Blaauw, E. (2019) 'Parental substance use disorder and child
30 abuse: risk factors for child maltreatment?', *Psychiatry, Psychology and Law*. doi:
31 10.1080/13218719.2019.1664277.
- 32 Grassi-Oliveira, R. et al. (2014) 'Childhood Trauma Questionnaire (CTQ) in
33 Brazilian samples of different age groups: Findings from Confirmatory Factor
34 Analysis.' *PLoS ONE* 9(1), e87118. doi:10.1371/journal.pone.0087118.
- 35 Gullbrå, F. et al. (2016) 'III and substance-abusing parents: how can the general
36 practitioner help their children? A qualitative study', *BMC Family Practice*, 17(1),
37 pp. 1–9. doi: 10.1186/s12875-016-0553-5.
- 38 Harwin, J. et al. (2014) *Changing lifestyles, keeping children safe: An evaluation of*
39 *the first Family Drug and Alcohol Court (FDAC) in Care Proceedings*. Brunel
40 University.
- 41 Horgan, J. (2011) 'Parental substance misuse: Addressing its impact on children. A
42 review of the literature'. National Advisory Committee on Drugs. pp.92.
- 43 Hughes, K. et al. (2017) 'The effect of multiple adverse childhood experiences on
44 health: A systematic review and meta-analysis', *Lancet Public Health*, 2(8), pp.
45 e356–e366. doi: 10.1016/S2468-2667(17)30118-4.
- 46 Institute of Medicine (IOM) and National Research Council (NRC). (2014) 'New
47 directions in child abuse and neglect research', Washington, DC: The National
48 Academies Press.
- 49 Kepple, N. J. (2017) 'The complex nature of parental substance use: examining past
50 year and prior use behaviors as correlates of child maltreatment frequency',
51 *Substance Use and Misuse*, 52(6), pp. 811–821. doi:
52 10.1080/10826084.2016.1253747.
- 53
54
55
56
57
58
59
60

- 1
2
3 Kubik, J., Docherty, M., and Boxer, P. (2019) 'The impact of childhood
4 maltreatment on adolescent gang involvement', *Child Abuse Neglect*, 96, 104096.
5 doi: 10.1016/j.chiabu.2019.104096.
6
7 Leeb, R. T., Lewis, T. and Zolotor, A. J. (2011) 'A review of physical and mental
8 health consequences of child abuse and neglect and implications for practice',
9 *American Journal of Lifestyle Medicine*, 5(5), pp. 454–468. doi:
10 10.1177/1559827611410266.
11 Maloney, E. et al. (2010) 'Prevalence and patterns of problematic alcohol use
12 among Australian parents', *Australian and New Zealand Journal of Public Health*,
13 34(5), pp. 495–501. doi: 10.1111/j.1753-6405.2010.00596.x.
14 Martins, M. C. and Pillon, S. C. (2008) 'A relação entre a iniciação do uso de drogas
15 e o primeiro ato infracional entre os adolescentes em conflito com a lei', *Cadernos
16 de Saude Publica*, 24(5), pp. 1112–1120. doi: 10.1590/S0102-311X2008000500018.
17 Melchior, M. et al. (2011) 'Parental alcohol dependence, socioeconomic
18 disadvantage and alcohol and cannabis dependence among young adults in the
19 community', *European Psychiatry*, 26(1), pp. 13–17. doi:
20 10.1016/j.eurpsy.2009.12.011.
21 Meyers, J. L. et al. (2018) 'Childhood interpersonal violence and adult alcohol,
22 cannabis, and tobacco use disorders: Variation by race/ethnicity?', *Psychology and
23 Medicine*. 48(9), pp. 1540–1550. doi.org/10.1017/S0033291717003208
24 Nadel, E. L., and Thornberry, T. P. (2017). Intergenerational consequences of
25 adolescent substance use: Patterns of homotypic and heterotypic continuity.
26 *Psychology of Addictive Behaviors*, 31(2), 200–211. doi.org/10.1037/adb0000248
27 Nelson, J. et al. (2017) 'Childhood maltreatment and characteristics of adult
28 depression: Meta-analysis', *British Journal of Psychiatry*, pp. 96–104. doi:
29 10.1192/bjp.bp.115.180752.
30 Ortegá, L. R. O. (2011) [A medida socioeducativa de liberdade assistida:
31 Fundamentos e contexto atual]. 145 f. [Master Dissertation] (Master in Social
32 Policy) — Universidade de Brasília, Brasília.
33 Rogosch, F.A., Oshri, A., and Cicchetti, D., (2010) 'From child maltreatment to
34 adolescent cannabis abuse and dependence: A developmental cascade model',
35 *Development and Psychopathology*, 22, pp. 883–897.
36 doi.org/10.1017/s0954579410000520
37 Sartes, L. M. A., De Micheli, D. and Souza-Formigoni, M. L. O. (2009)
38 'Psychometric and discriminative properties of the teen addiction severity index
39 (Brazilian Portuguese version)', *European Child and Adolescent Psychiatry*, 18(11),
40 pp. 653–661. doi: 10.1007/s00787-009-0021-z.
41 Solis, J. M. et al. (2012) 'Understanding the diverse needs of children whose parents
42 abuse substances', *Current Drug Abuse Reviews*, pp. 135–147.
43 Taplin, S. and Mattick, R. (2011) *Child protection and mothers in substance abuse
44 treatment*. Sydney: National Drug and Alcohol Research Centre.
45 Teicher, M. H. and Samson, J. A. (2013) 'Childhood maltreatment and
46 psychopathology: A case for ecophenotypic variants as clinically and
47 neurobiologically distinct subtypes', *American Journal of Psychiatry*, 170(10), pp.
48 1114–1133. doi: 10.1176/appi.ajp.2013.12070957.
49 Torvik, F. A. et al. (2011) 'Parental alcohol use and adolescent school adjustment in
50 the general population: Results from the HUNT study', *BMC Public Health*, 11. doi:
51 10.1186/1471-2458-11-706.
52 Velleman, R. and Templeton, L. J. (2016) 'Impact of parents' substance misuse on
53 children: an update', *BJPsych Advances*, 22(2), pp. 108–117. doi:
54
55
56
57
58
59
60

1
2
3 10.1192/apt.bp.114.014449.

4 Verdolini, N. et al. (2015) 'Traumatic events in childhood and their association with
5 psychiatric illness in the adult', *Psychiatria Danubina*, 27, pp. S60–S70.

6 Walsh, D. et al. (2019) 'Relationship between childhood socioeconomic position
7 and adverse childhood experiences (ACEs): A systematic review', *Journal of
8 Epidemiol Community Health*, 73(12), pp. 1087–1093. doi: 10.1136/jech-2019-
9 212738.

10
11 Winters, A. (2014) 'Influential factors of parental substance and alcohol abuse on
12 children's academic achievement', *Research Journal*, pp.1-13. Portland States
13 University.

14 World Health Organization (2014).) Global status report on violence prevention.

15 World Health Organization. Retrieved from:

16 <http://www.who.int/iris/handle/10665/145086>

17
18 Yudko, E., Lozhkina, O. and Fouts, A. (2007) 'A comprehensive review of the
19 psychometric properties of the Drug Abuse Screening Test', *Journal of Substance
20 Abuse Treatment*, 32(2), pp. 189–198. doi: 10.1016/j.jsat.2006.08.002.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Sociodemographic profile of participants

		Parental substance use and neglect of family care [n (%)]			
		Yes	No	OR (95%CI)	ORA (95%CI)
Gender	Male	104 (72.7)	39 (27.3)	Ref.	
	Female	3 (42.9)	4 (57.1)	3.6 (0.76;16.61)	-
		$\chi^2(1) = 2.912 p=0.088$			
Age group (years)	13-14	8 (57.1)	6 (42.9)	2.4 (0.72;8.02)	-
	15-16	48 (76.2)	15 (23.8)	Ref.	-
	17-18	51 (69.9)	22 (30.1)	1.4 (0.64;2.97)	-
		$\chi^2(2) = 2.183 p=0.336$			
Schooling	No studying	54 (72.0)	21 (28.0)	Ref.	-
	Studying	53 (70.7)	22 (29.3)	1.0 (0.53;2.17)	-
		$\chi^2(1) = 0.033 p=0.857$			
Religion	None	62 (74.7)	21 (25.3)	1.1 (0.32;3.75)	-
	Catholic	18 (72.0)	7 (28.0)	1.2 (0.30;5.23)	-
	Evangelic	13 (76.5)	4 (23.5)	Ref.	-
	Others	14 (56.0)	11 (44.0)	2.5 (0.65;10.06)**	-
		$\chi^2(3) = 3.559 p=0.313$			
Skin color	White	27 (71.0)	11 (28.9)	1.0 (0.45;2.29)	-
	Non-White	80 (71.4)	32 (28.6)	Ref.	-
		$\chi^2(2) = 0.019 p=0.991$			
Living with parents	Yes	19 (55.9)	15 (44.1)*	2.5 (1.12;5.52)**	2.7 (1.13;6.37)**
	No	88 (75.9)	28 (24.1)	Ref.	Ref.
		$\chi^2(1) = 5.13; p = 0.023^*$			
Supported by funds from theft	Yes	46 (63.0)	27 (37.0)*	2.2 (1.08;4.63)**	2.0 (0.96;4.48)
	No	61 (79.2)	16 (20.8)	Ref.	Ref.
		$\chi^2(1) = 4.81; p = 0.028^*$			
Supported by funds from drug trafficking	Yes	84 (68.8)	38 (31.2)	2.0 (0.74;5.89)	-
	No	23 (82.1)	5 (17.9)	Ref.	-
		$\chi^2(1) = 1.967 p=0.161$			
Familiar income	<1 MW	5 (35.7)	9 (64.3)*	1.6 (0.64;4.15)	1.5 (0.58;4.14)
	1 - 2	25 (69.4)	11(30.6)*	6.6 (1.90;23.27)*	6.7 (1.85;24.22)*
	≥ 2	29 (74.4)	10 (25.6)	1.3 (0.50;3.27)	1.0 (0.38;2.87)
	Don't know	48 (78.7)	13 (21.3)	Ref.	Ref.
		$\chi^2(3) = 10.35; p = 0.015^*$			

Note: Minimum Wage [MW]. (Value of MW in Brazilian Real R\$ 937.00).

Chi-square test χ^2 (Degrees of freedom).

**p-value ≤ 0.05 and *p-value ≤ 0.01 .

Odds Ratio Unadjusted (OR) and Odds Ratio Adjusted (ORA).

Confidential Interval (CI)

N = 150

Table 2. Symptoms and mental health problems (ASI-Teen) experienced by participants

		PSU and neglect of family care [n (%)]		OR (CI95%)	ORA (CI95%)
		Yes	No		
Depression	Yes	14 (42.4)	19 (57.6)	2.2 (0.99;5.02)	-
	No	29 (24.8)	88 (75.2)	Ref.	-
		$\chi^2(1) = 3.916 p=0.048^{**}$			
Severe anxiety or tension	Yes	26 (33.3)	52 (66.7)	1.6 (0.79;3.32)	-
	No	17 (23.6)	55 (76.4)	Ref.	-
		$\chi^2(1) = 1.731 p=0.188$			
Halucinations	Yes	17 (46.0)	20 (54.0)	2.8 (1.30;6.21)*	2.8 (1.25;6.14)*
	No	26 (23.0)	87 (77.0)	Ref.	Ref.
		$\chi^2(1) = 7.171 p=0.007^{**}$			
Problems controlling violent behaviour	Yes	15(45.5)	18 (54.5)	2.6 (1.18;5.93)**	2.6 (1.12;5.87)**
	No	28 (23.9)	89 (76.1)	Ref.	Ref.
		$\chi^2(1) = 5.831 p=0.016^{**}$			
Problems with cognition, concentration or memory.	Yes	23 (33.3)	46 (66.7)	1.5 (0.75;3.10)	-
	No	20 (24.7)	61 (75.3)	Ref.	-
		$\chi^2(1) = 1.361 p=0.243$			

Note: Chi-square test χ^2 (Degrees of Freedom).

*p-value ≤ 0.01 and **p-value ≤ 0.05

Odds Ratio Unadjusted (OR) and Odds Ratio Adjusted (ORA).

Confidential Interval (CI)

N = 150

For Peer Review

Table I. PSU and neglect of family care and Childhood Trauma Questionnaire (CTQ).

		PSU and neglect of family care [n (%)]			
		Yes	No	OR Unadjusted (95%CI)	p-value
<i>Physical Neglect</i>					
Yes		39 (28.1)	100 (71.9)	Ref.	
No		4 (36.4)	7 (63.6)	1.5 (0.41;5.29)	0.56
		$\chi^2(1) = 0.344 p=0.558$			
<i>Emocional Neglect</i>					
Yes		36 (26.7)	99 (73.3)	Ref.	
No		7 (46.7)	8 (53.3)	2.4 (0.81;7.11)	0.11
		$\chi^2(1) = 2.641 p=0.104$			
<i>Physical Abuse</i>					
Yes		29 (33.0)	59 (67.0)	1.7 (0.80;3.54)	0.17
No		14 (22.6)	48 (77.4)	Ref.	
		$\chi^2(1) = 1.914 p=0.166$			
<i>Sexual Abuse</i>					
Yes		4 (66.7)	2 (33.3)	5.4 (0.95;30.58)	0.057
No		39 (27.1)	105 (72.9)	Ref.	
		$\chi^2(1) = 4,413 p=0.036^{**}$			
<i>Emocional Abuse</i>					
Yes		36 (40.0)	54 (60.0)	5.0 (2.06;12.34)	<0.01**
No		7 (11.7)	53 (88.3)	Ref.	
		$\chi^2(1) = 14,133 p<0.001^*$			

Note: Chi-square test χ^2 (Degrees of freedom).

*p-value ≤ 0.01 and **p-value ≤ 0.05

Odds Ratio Unadjusted (OR)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential Interval (CI)
N = 150

For Peer Review

Table II. Type of abuse associated with maltreatment (CTQ) experienced by adolescents.

		PSU and neglect of family care [n (%)]		OR (CI95%)	ORA (CI95%)
		Yes	No		
Emotional Abuse					
<i>People in my family called me things like “stupid”, “lazy”, or “too ugly”.</i>	Yes	21 (44.7)*	26 (55.3)	2.9 (1.41;6.25)*	1.8 (0.72;4.63)
	No	22 (21.4)	81 (78.6)	Ref.	Ref.
		$\chi^2(1)=8.584$ $p=0.003$			
<i>There was nothing I wanted to change in my family.</i>	Yes	26 (40.6)*	38 (59.4)	2.8 (1.34;5.75)*	1.9 (0.88;4.48)
	No	17 (19.8)	69 (80.2)	Ref.	Ref.
		$\chi^2(1) = 7.806$ $p=0.005$			
<i>I thought that my parents wished that I had never been born.</i>	Yes	18(42.9)*	24 (57.1)	2.5 (1.17;5.31)**	1.1 (0.45;2.87)
	No	25 (23.2)	83 (76.8)	Ref.	Ref.
		$\chi^2(1) = 5.744$ $p=0.017$			
<i>People in my family said things that hurt or offended me.</i>	Yes	23 (43.4)*	30 (56.6)	2.9 (1.42;6.14)*	1.4 (0.56;3.65)
	No	20 (20.6)	77 (79.4)	Ref.	Ref.
		$\chi^2(1) = 8.696$ $p=0.003$			
Physical Abuse					
<i>I got beaten so much by somebody from my family that therefore I had to go to a hospital or to see a doctor</i>	Yes	6 (66.7)*	3 (33.3)	5.6 (1.34;23.63)**	2.6 (0.36;18.86)
	No	37 (26.2)	104 (73.8)	Ref.	Ref.
		$\chi^2(1) = 6.761$ $p=0.009$			
<i>Somebody in my family hit me so much that it left me with marks or bruises</i>	Yes	15 (51.7)*	14 (48.3)	3.5 (1.53;8.26)*	1.4 (0.45;4.42)
	No	28 (23.1)	93 (76.9)	Ref.	Ref.
		$\chi^2(1) = 9.347$ $p=0.002$			
<i>I got beaten so much that it was seen by a teacher, neighbour or doctor.</i>	Yes	6 (60.0)*	4 (40.0)	4.2 (1.12;15.63)**	0.91 (0.12;6.77)
	No	37 (26.4)	103 (73.6)	Ref.	Ref.
		$\chi^2(1) = 5.144$ $p=0.023$			
Physical Neglect					
<i>I had to wear dirty clothes.</i>	Yes	16 (50.0)*	16 (50.0)	3.4 (1.49;7.62)*	3.0 (1.24;7.49)**
	No	27 (22.9)	91 (77.1)	Ref.	Ref.
		$\chi^2(1) = 9.053$ $p=0.003$			

Note: Chi-square test χ^2 (Degrees of freedom)

*p-value ≤ 0.01 and **p-value ≤ 0.05

Odds Ratio Unadjusted (OR) and Odds Ratio Adjusted (ORA)

Confidential Interval (CI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

N = 150

For Peer Review

Table III. Substance used in the last 30 days by adolescents (ASI-Teen and DAST-20).

		PSU and neglect of family care [n (%)]		
		Yes	No	OR (CI95%)
Tobacco	Yes	19 (40.4)*	28 (59.6)	2.2 (1.07;4.68)*
	No	24 (23.3)	79 (76.7)	Ref.
$\chi^2(1) = 4.628 p = 0.031$				
Inhalants	Yes	13 (31.7)	28 (68.3)	1.2 (0.56;2.67)
	No	30 (27.5)	79 (72.5)	Ref.
$\chi^2(1) = 0.255 p=0.614$				
Cannabis	Yes	25 (29.1)	61 (70.9)	1.0 (0.51;2.14)
	No	18 (28.1)	46 (71.9)	Ref.
$\chi^2(1) = 0.016 p=0.899$				
Alcohol	Yes	20 (33.9)	39 (66.1)	1.5 (0.74;3.11)
	No	23 (25.3)	68 (74.7)	Ref.
$\chi^2(1) = 1.302 p=0.254$				
DAST-20	Moderate	22 (34.9)	41 (65.1)	1.7 (0.83;3.44)
	No problem	21 (24.1)	66 (75.9)	Ref.
$\chi^2(1) = 2.078 p=0.149$				

Note: Chi-square test χ^2 (Degrees of freedom).

*p-value ≤ 0.01 and **p-value ≤ 0.05 .

Odds Ratio Unadjusted (OR) and Odds Ratio Adjusted (ORA).

Confidential Interval (CI)

N = 150