

Newborn pulse oximetry screening in the UK

Brown, Sebastian ; Liyanage, Shehan; Mikrou, Paraskevi; Singh, Anju; Ewer, Andrew

DOI:

[10.1016/S0140-6736\(20\)31959-0](https://doi.org/10.1016/S0140-6736(20)31959-0)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Brown, S, Liyanage, S, Mikrou, P, Singh, A & Ewer, A 2020, 'Newborn pulse oximetry screening in the UK: 2020 survey', *The Lancet*, vol. 396, no. 10255, P881. [https://doi.org/10.1016/S0140-6736\(20\)31959-0](https://doi.org/10.1016/S0140-6736(20)31959-0)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Lancet

Newborn pulse oximetry screening in the UK: a survey of practice in 2020

--Manuscript Draft--

Manuscript Number:	
Article Type:	Correspondence
Keywords:	Newborn; screening; pulse oximetry; congenital heart defect
Corresponding Author:	Andrew Ewer Birmingham Women's Hospital Birmingham, UNITED KINGDOM
First Author:	Andrew Ewer
Order of Authors:	Andrew Ewer Anju Singh Sebastian Brown Paraskemi Mikrou Shehan Liyanage
Manuscript Region of Origin:	UNITED KINGDOM
Additional Information:	
Question	Response
If you are writing in response to published content in The Lancet, please type the full reference to that content here (for example: Cuzick J et al. Use of anastrozole for breast cancer prevention (IBIS-II): long-term results of a randomised controlled trial. Lancet 2020; 394: 117-22).	N/A
Author Comments:	<p>Dear Editor</p> <p>Pulse oximetry screening (POS) has attracted considerable international attention over the past 10 years and the Lancet has been at the forefront, publishing data and opinion which has influenced POS practice worldwide.</p> <p>POS has been recommended by many countries. Recent USA data show that POS reduces death from critical congenital heart defects (CCHD) by over one third.</p> <p>The UK National Screening Committee (NSC) has been considering routine POS since 2013. However, despite overwhelming supporting evidence and two national public consultations, the NSC has yet to recommend this life saving test. As a result newborn babies in the UK are subject to an inequality of service with many denied the opportunity and as a consequence some serious conditions are missed.</p> <p>This survey from all UK neonatal units reflects the further change in practice which has occurred, not least because of the Lancet's support for this important intervention.</p> <p>We hope this will give a vital impetus to the goal of achieving universal POS in the UK and beyond. Many UK units are still waiting for national guidance on this issue.</p>

More babies will have died from undiagnosed CCHD in the last months than from Covid 19.

Prof Andrew Ewer

Dear Editor

Newborn pulse oximetry screening in the UK: a survey of practice in 2020

Routine newborn pulse oximetry screening (POS) identifies babies with critical congenital heart defects (CCHD) that would otherwise have been missed by antenatal ultrasound and postnatal examination.¹ POS reduces mortality from CCHD², and identifies babies with other important conditions such as respiratory disorders and sepsis.

Many countries now recommend POS, but the UK National Screening Committee (NSC) is still undecided.³ Despite the absence of a national recommendation, many UK neonatal units have introduced POS. In 2017 uptake of POS was 40% (an increase from 7% in 2010).⁴

Following a second public consultation by the NSC in 2019, we surveyed all UK neonatal units to see if practice had changed.

Between January and May 2020, lead clinicians from all 189 UK Neonatal Units were invited to complete an online questionnaire with telephone follow-up for non-responders.

189 units responded (100%). Ninety-six units (51%) are currently using routine POS, including the majority of Neonatal Intensive Care Units (55%) and Special Care Units (60%). POS was used in 43% of Local Neonatal Units. Uptake across the UK varied from 100% in Wales and 51% in England, to 29% in Scotland and Northern Ireland. Although there are screening units in all regions of England, uptake in the North is greater than in the South (Figure 1).

Most units screen within 24 hours (57%) or before discharge (29%) and the PulseOx study⁵ algorithm (pre and post ductal saturations less than 95% or > 2% difference being abnormal).

Of the 93 non-screening units, 28% were considering introduction in the near future, but 66% were waiting for a recommendation by the NSC. Perceived barriers to introduction included resources and staffing (47%), lack of local echocardiography (25%), training (21%) and concerns about false positives (9%). Additional perceived barriers were the lack of national screening guidelines (18%), and 'change' itself (12%). Only 6 units (6%) were not considering POS. All six cited concerns over false positives, three were unconvinced by the data, two had insufficient staff and one felt that POS was an unnecessary test.

We asked screening units if they felt that POS had led to an increase in unnecessary investigations. 78% felt that screening did not increase these investigations, and 10% felt that any small increase was justified and offset by the benefits of identifying significant cardiac and non-cardiac pathology.

The majority of UK Neonatal units now use routine POS, although there are still regional differences in uptake and variation in implementation. The need for a national recommendation is more compelling than ever.

References

1. Plana MN, Zamora J, Suresh G, Fernandez-Pineda L, Thangaratinam S, Ewer AK. Pulse oximetry screening for critical congenital heart defects. *Cochrane Database Syst Rev.* 2018;3(3):CD011912. Published 2018 Mar 1. doi:10.1002/14651858.CD011912.pub2
2. Abouk R, Grosse SD, Ailes EC, Oster ME. Association of US State Implementation of Newborn Screening Policies for Critical Congenital Heart Disease With Early Infant Cardiac Deaths [published correction appears in JAMA. 2018 Sep 25;320(12):1288]. *JAMA.* 2017;318(21):2111-2118. doi:10.1001/jama.2017.17627
3. Oddie S, Stenson B, Wyllie J, Ewer AK. UK consultation on pulse oximetry screening for critical congenital heart defects in newborns. *Lancet* 2019;394:103-104.[doi.org/10.1016/S0140-6736\(19\)31515-6](https://doi.org/10.1016/S0140-6736(19)31515-6).
4. Mikrou P, Singh A, Ewer AK. Pulse oximetry screening for critical congenital heart defects: a repeat UK national survey. *Arch Dis Child Fetal Neonatal Ed.* 2017;102(6):F558. doi:10.1136/archdischild-2017-313378
5. Ewer AK, Middleton LJ, Furmston AT, et al. Pulse oximetry screening for congenital heart defects in newborn infants (PulseOx): a test accuracy study. *Lancet.* 2011;378(9793):785-794. doi:10.1016/S0140-6736(11)60753-8

Seb Brown,¹ Shehan Liyanage¹ , P Mikrou², Anju Singh,¹ Andrew K Ewer^{1,3}

- 1 Neonatal Unit, Birmingham Women's Hospital, Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK
- 2 General Paediatric Department, Staffordshire Children's Hospital at Royal Stoke, University Hospital of North Midlands NHS Trust, UK
- 3 Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

Corresponding author

Prof AK Ewer Neonatal Unit

Birmingham Women's Hospital

Edgbaston

Birmingham

UK

B15 2TG

Email: a.k.ewer@bham.ac.uk

Tel: +44 121 472 1377

Fax: +44 121 627 2749




Figure 1 - Percentage of units performing POS in different regions of England