

Cost-effectiveness of bariatric surgery versus community weight management to treat obesity-related idiopathic intracranial hypertension

Elliot, Laura; Frew, Emma; Mollan, Susan; Mitchell, James; Yiangou, Andreas; Alimajstorovic, Zerin; Ottridge, Ryan; Wakerley, Benjamin; Thaller, Mark; Grech, Olivia; Singhal, Rishi; Tahrani, Abd; Harrison, Mark; Sinclair, Alex; Aguiar, Magda

DOI:

[10.1016/j.soard.2021.03.020](https://doi.org/10.1016/j.soard.2021.03.020)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Elliot, L, Frew, E, Mollan, S, Mitchell, J, Yiangou, A, Alimajstorovic, Z, Ottridge, R, Wakerley, B, Thaller, M, Grech, O, Singhal, R, Tahrani, A, Harrison, M, Sinclair, A & Aguiar, M 2021, 'Cost-effectiveness of bariatric surgery versus community weight management to treat obesity-related idiopathic intracranial hypertension: evidence from a single payer health care system', *Surgery for Obesity and Related Diseases*, vol. 17, no. 7, pp. P1310-1316. <https://doi.org/10.1016/j.soard.2021.03.020>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Original article

Cost-effectiveness of bariatric surgery versus community weight management to treat obesity-related idiopathic intracranial hypertension: evidence from a single-payer healthcare system

Laura Elliot, M.Sc.^a, Emma Frew, Ph.D.^{b,*}, Susan P. Mollan, F.R.C.Ophth.^c,
James L. Mitchell, M.C.R.P.^{d,e}, Andreas Yiangou, M.R.C.P.^{d,e,f},
Zerin Alimajstorovic, Ph.D.^{e,f,g}, Ryan S. Ottridge, M.Phil.^h, Ben R. Wakerley, Ph.D.ⁱ,
Mark Thaller, M.R.C.P.^{d,e}, Olivia Grech, M.Res.^{e,f,g}, Rishi Singhal, F.A.C.S.^{j,k},
Abd A. Tahrani, Ph.D.^{f,g,l}, Mark Harrison, Ph.D.^{m,n}, Alexandra J. Sinclair, Ph.D.^{d,e,f},
Magda Aguiar, Ph.D.^m

^aDepartment of Economics and Related Studies, University of York, York, United Kingdom

^bInstitute of Applied Health Research, University of Birmingham, Birmingham, United Kingdom

^cBirmingham Neuro-Ophthalmology, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital, Birmingham, United Kingdom

^dDepartment of Neurology, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital, Birmingham, United Kingdom

^eMetabolic Neurology, Institute of Metabolism and Systems Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom

^fCentre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, United Kingdom

^gInstitute of Metabolism and Systems Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom

^hBirmingham Clinical Trials Unit, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom

ⁱDepartment of Neurology, Gloucestershire Hospitals NHS Foundation Trust, Gloucester, United Kingdom

^jUpper GI Unit and Minimally Invasive Unit, Birmingham Heartlands Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom

^kInstitute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, United Kingdom

^lDepartment of Endocrinology and Diabetes, University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom

^mCollaboration for Outcomes Research and Evaluation, Faculty of Pharmaceutical Sciences, University of British Columbia, Vancouver, Canada

ⁿCentre for Health Evaluation and Outcome Sciences, St. Paul's Hospital, Vancouver, Canada

Received 12 February 2021; accepted 21 March 2021

Abstract

Background: Idiopathic intracranial hypertension (IIH) is associated with significant morbidity, predominantly affecting women of childbearing age living with obesity. Weight loss has demonstrated successful disease-modifying effects; however, the long-term cost-effectiveness of weight loss interventions for the treatment of IIH has not yet been established.

Objectives: To estimate the cost-effectiveness of weight-loss treatments for IIH.

Setting: Single-payer healthcare system (National Health Service, England).

Methods: A Markov model was developed comparing bariatric surgery with a community weight management intervention over 5-, 10-, and 20-year time horizons. Transition probabilities, utilities, and resource use were informed by the IIH Weight Trial (IIH:WT), alongside the published literature. A probabilistic sensitivity analysis was conducted to characterize uncertainty within the model.

Alexandra J. Sinclair, and Magda Aguiar are joint senior authors.

E-mail address: E.Frew@bham.ac.uk (E. Frew).

*Correspondence: Emma Frew, Ph.D., Institute of Applied Health Research, University of Birmingham, Birmingham, B15 2TT, UK.

Results: In the base case analysis, over a 20-year time horizon, bariatric surgery was “dominant,” led to cost savings of £49,500, and generated an additional 1.16 quality-adjusted life years in comparison to the community weight management intervention. The probabilistic sensitivity analysis indicated a probability of 98% that bariatric surgery is the dominant option in terms of cost-effectiveness.

Conclusion: This economic modeling study has shown that when compared to community weight management, bariatric surgery is a highly cost-effective treatment option for IIH in women living with obesity. The model shows that surgery leads to long-term cost savings and health benefits, but that these do not occur until after 5 years post surgery, and then gradually increase over time. (*Surg Obes Relat Dis* 2021;17:1310–1316.) © 2021 American Society for Bariatric Surgery. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Keywords:

Cost-effectiveness; Bariatric surgery; Idiopathic intracranial hypertension; Weight loss

Idiopathic intracranial hypertension (IIH) is a highly incapacitating condition, characterized by raised intracranial pressure (ICP) [1,2], that mainly affects women living with obesity [3]. Elevated ICP can lead to swelling of the optic nerve head, referred to as papilledema, which can potentially result in blindness [4]. IIH is typically associated with significant morbidity due to chronic, disabling headaches [5] and varying degrees of visual disturbances that cause a reduction in health-related quality of life (HRQoL) and productivity loss in the workplace [6].

The pathogenesis of IIH remains unclear [7]; however, it predominately affects females aged between 25 and 36 years, and obesity is a major risk factor [8]. The annual incidence of IIH in female patients increased from 2.5 per 100,000 person-years in 2005 to 9.3 per 100,000 person-years in 2017 [9]. Once considered a rare condition, the burden of IIH is rapidly increasing [3]. Hospital IIH admissions in England have risen by 442% between 2002 and 2014, and associated costs over the same period increased from £9.2 million to £50 million per annum [3]. This rising incidence of IIH is significantly correlated with rising obesity levels [3]. If the IIH incidence continues to increase at the same rate, these annual costs are projected to increase to £462 million by 2030 [3]. These increasing excessive costs have also been shown in the United States, where the total hospital costs per IIH admission in 2007 were 4 times greater than for a population-based per-person admission, with the total economic costs of IIH patients exceeding \$444 million [10].

Weight loss is currently the only established disease-modifying therapy for IIH [8,11]. Lifestyle behavioral weight-loss interventions [12] resulting in 15% weight loss are effective in lowering ICP, improving papilledema and visual functions, and decreasing headache frequency and severity, with concomitant reduction in analgesia usage [11], but long-term weight maintenance is challenging, leading to weight regain in the majority of patients [6]. A recent meta-analysis found reports and series that have shown the beneficial clinical effects of bariatric surgery in resolving IIH [13]. Bariatric surgery is safe and is the

most effective method of achieving weight loss (25%–30%) that is sustainable over the long term [14], and consequently reducing IIH, but with high upfront costs. Hence, more research is needed to understand the long-term cost-effectiveness of surgery when compared to lifestyle weight loss interventions.

In England, the National Institute for Health and Care Excellence (NICE) currently recommends bariatric surgery for people with a body mass index (BMI) over 40 kg/m² and for people with a BMI over 35 kg/m² who fulfill certain criteria. IIH is not currently considered as a significant comorbidity associated with obesity that would qualify for bariatric surgery in patients below a BMI of 40 kg/m².

With the goal of informing new treatment recommendations for IIH and guidelines for bariatric surgery, the IIH weight trial (IIH:WT) was designed to compare the efficacy and cost-effectiveness of bariatric surgery and community weight management interventions [15]. The IIH:WT was conducted within England, and therefore within the context of a single-payer healthcare system (National Health Service [NHS]). This paper reports on the economic evaluation by extrapolating the trial data and modeling the long-term cost-effectiveness from the perspective of the healthcare service in England. The gains are captured using quality-adjusted life years (QALYs), which are commonly used to enable judgements about cost-effectiveness. The IIH:WT findings are modeled over 5-, 10-, and 20-year time horizons to assess the long-term cost-effectiveness of bariatric surgery as a treatment for IIH, compared to a community weight management program.

Methods

Study design

IIH:WT was a randomized, controlled, parallel-group, multi-center trial [16]. Inclusion criteria for the trial were female patients with active IIH and no other significant comorbidity, aged between 18–55 years, with a BMI ≥ 35 kg/m², consistent with the NICE obesity guideline [17]. The National Research Ethics Committee West Midlands

approved the trial (14/WM/0011). The economic evaluation was conducted using a decision-analytic model to facilitate extrapolation of trial findings over an extended time horizon.

Model structure

A Markov model was applied to reflect cyclical fluctuations in weight over time and to allow for the consideration of weight recidivism following bariatric surgery. The model structure is shown in Fig. 1. The model compares bariatric surgery with community weight management, and therefore the surgical arm comprises the suite of surgical procedures that were performed within the IIH:WT. A cycle length of 1 year was used, as this was considered to be the shortest sufficient time period for patients to change between BMI categories. The time horizons for the model were 5, 10, and 20 years.

The model started with a hypothetical cohort of 1000 patients who were considered to have a health state of either obesity or severe obesity, as determined by their BMI and in line with NICE guidance for eligibility for bariatric surgery [17]. Classifying the states according to weight (BMI) category enabled the effects of weight recidivism on IIH status to be captured, as well as allowed for the wider consideration of the health impacts of obesity, such as the incidence of co-morbidities other than IIH.

Patients were distributed between the baseline health states in line with the distribution of BMI from the IIH:WT. Following intervention, the patients then progressed to 1 of 3 states: severe obesity (BMI ≥ 40 kg/m²), obesity (BMI 30–39.9 kg/m²), or no obesity (BMI < 30 kg/m²), or into an absorbing state, death. Maximum weight loss generally occurs between 12–24 months following surgery [18], and from that point onwards, patients typically experience

some degree of weight regain. Weight regain continues gradually until approximately 10 years postsurgery, at which point it plateaus [19]. To reflect this process, the model required distinct transition probabilities between health states over 4 time periods: cycle 1, cycle 2, cycles 3–10, and cycle 11 onwards.

Sources of data

Both primary and secondary data were used to inform the model parameters (Table S1). Data on resource use, costs, and effectiveness were derived from the IIH:WT and supplemented with data from targeted literature review where necessary.

Transition probabilities

The IIH:WT followed patients for 24 months, and these data were used to estimate the transition probabilities for the first 2 model cycles (Table S2). From year 2 onwards, transition probabilities were calculated using weight regain data from the Swedish Obese Subjects (SOS) study [19], in conjunction with data from the IIH:WT. Once the patients reached cycle 10, their weight was assumed to plateau. Hence, the transition probabilities applied in cycles 11–20 assume only the possibility of transitioning to dead or remaining within the same state. The model included a BMI-specific mortality risk [20] and an additional mortality risk associated with bariatric surgery [21].

Costs

The costs included in the model were those relevant to the healthcare service, detailed in Table S1. All costs are reported in 2017/2018 GBP prices [22]. For the surgical arm, the design of the IIH:WT did not predetermine the




Fig. 1. Markov model structure. The structure is identical for both the bariatric surgery arm and the community weight management intervention arm. Arrows indicate possible transitions between states. Patients can transition to “Dead” from any of the states. The dashed line indicates the interventions taking place following baseline measurements.

Table 1
Cost-utility analysis

	Time horizon for model						Additional sensitivity analysis			
	5 yr		10 yr		20 yr		Scenario analysis		Probabilistic sensitivity analysis	
	Surgery	WM	Surgery	WM	Surgery	WM	Surgery	WM	Surgery	WM
Total costs	£15,900	£27,400	£29,900	£58,600	£55,100	£112,400	£81,100	£143,900	£65,500	£118,600
Incremental costs		–£11,500		–£28,700		–£57,300		–£62,900		–£53,000
Total QALYs	3.83	3.53	7.62	7.04	14.28	13.12	14.28	13.12	14.26	13.14
Incremental QALYs		.29		.58		1.16		1.16		1.12
ICER*		–£39,000		–£49,300		–£49,500		£54,300		–£47,200

WM = weight management; QALY = quality-adjusted life years; ICER = incremental cost-effectiveness ratio.

* ICER = difference in total costs between surgery versus WM/Difference in total QALYs between surgery versus WM.

type of surgery to be given, as this was a pragmatic decision to reflect routine clinical practice. The cost of bariatric surgery therefore was determined by the weighted average of the 3 surgery types performed—laparoscopic adjustable gastric band (LAGB), laparoscopic sleeve gastrectomy (LSG), and Roux-en-Y gastric bypass (RYGB)—as the intention of the model was not to recommend a surgery type, but rather to assess the incremental cost-effectiveness of surgery versus community weight management. The total cost of surgery was inclusive of the weighted costs of revisional surgery for gastric banding. Data on resource use derived from the IIH:WT were used in conjunction with the NHS National Tariff [22,23] to calculate the average costs per patient, per cycle for the first 2 cycles. The cost of the community weight management program comprised WeightWatchers vouchers that provided access to online and mobile tools for 12 months.

For each health state, the average cost of IIH management was estimated by multiplying the incidence of IIH by the annual per-patient costs of IIH [3]. Evidence suggests that weight regain results in recurrence of IIH within both individuals living with obesity and those with a healthy weight [6,24,25]. It was therefore assumed that IIH would recur at the same rate regardless of whether the patient had lost weight and then subsequently regained weight or remained in an obesity health state throughout.

Outcomes

Utility values for each health state were derived from the IIH:WT data (Table S1). HRQoL was assessed using the EuroQoL-5 dimension-5 level (EQ-5D-5L) instrument at baseline and at 12 and 24 months, and it was assumed that utilities remained constant for each health state over time.

All costs and benefits were discounted at 3.5%, consistent with UK NICE guidelines.

Cost-utility analysis

The cost-utility analysis estimated the cost of the change in QALYs due to surgery when compared with that of the weight management program. This results in an incremental

cost-effectiveness ratio, which is the difference in costs divided by the difference in QALYs and gives an estimate of the cost per QALY gained.

Scenario and sensitivity analysis

To assess the uncertainty around the model parameters, a probabilistic sensitivity analysis was conducted. The probability that surgery is cost-effective when compared to weight management is then estimated for different threshold values of willingness to pay for a QALY, presented as a cost-effectiveness acceptability curve.

To account for the wider cost impact from weight loss and lowering the incidence of these co-morbidities associated with obesity, a scenario analysis was conducted that included costs associated with type 2 diabetes and coronary heart disease (CHD). The annual incidences and associated healthcare costs of CHD and type 2 diabetes were obtained [26] and applied to the health states using the same method described above for IIH. Any disutility associated with diabetes and CHD was not applied, as it was felt that this was already captured within the self-reported HRQoL values from the trial data.

Results

Across all 3 time horizons of 5, 10, and 20 years, bariatric surgery dominates (less cost and more benefit) the community weight management intervention, generating more QALYs with cost savings (Table 1). The extent of dominance increases over time, with more cost savings accumulating the longer patients are followed-up. At 20 years, surgery led to an incremental cost saving of £49,500 and an additional 1.16 QALYs when compared to community weight management. When the additional costs of type 2 diabetes and CHD were considered, these cost savings increased to £54,300.

The probabilistic sensitivity analysis confirmed the base case results, producing a cost saving of £47,200 and an incremental gain of 1.12 QALYs. The distribution of incremental cost-effectiveness ratios is shown in Fig. S1. At a willingness to pay £20,000 per QALY, the lower threshold

used by decision-making bodies such as NICE, there is a 98% chance that bariatric surgery is cost-effective when compared to the community weight management intervention (Fig. 2).

Discussion

This paper is the first to report on the long-term cost-effectiveness of bariatric surgery compared to a community weight management intervention to treat IHH. This was done within the context of a single-payer healthcare system. The economic model combined the costs associated with bariatric surgery to treat IHH with the HRQOL benefits and found it to increase QALYs and decrease costs, when compared to community weight management. This finding was robust to sensitivity analysis. Varying the time horizon showed that both the incremental costs saved and the incremental QALYs gained from surgery increase over time.

Although there is evidence for the cost-effectiveness of bariatric surgery and despite multiple guidelines for treatment, including from NICE [17], access to bariatric surgery within the English healthcare service remains limited, with less than .002% of the potentially eligible adults having surgery annually [27]. This is due to barriers for referral from primary care and funding constraints from the commissioners [27]. This economic model demonstrates that improving access to bariatric surgery is likely to be cost

saving, reduce the burden of IHH in women with BMIs ≥ 35 kg/m², and improve patient HRQoL.

The economic model only included the co-morbidities of type 2 diabetes and CHD as part of a scenario analysis. Expanding the model to account for additional cost savings from reducing the risks of further obesity-related co-morbidities will only make bariatric surgery even more cost saving. Furthermore, the model was conducted from a health service perspective, meaning that any out-of-pocket payments or indirect costs of IHH were not included. If these indirect costs, such as days off work or time to travel to appointments to manage ongoing IHH symptoms, were included, then this would make surgery even more cost saving. The model classified health states according to BMI category with the aim of capturing the wider health impacts of obesity (type 2 diabetes and CHD), as well as effects of weight recidivism on IHH status. A strength of the model is that it used data from the IHH:WT study, including EQ-5D-5L data that were directly collected from patients to construct QALYs. And as the patients moved between the BMI health states over time, the effects of weight loss upon IHH symptoms and wider obesity effects will have been captured within this EQ-5D-5L data and modeled accordingly. Using QALYs as a commensurate outcome allows comparisons of cost-effectiveness to be made between alternative treatments across different disease areas.




Fig. 2. Cost-effectiveness acceptability curve. QALY = quality-adjusted life years.

There are some limitations to consider. The model uses data from the IIH:WT, and as IIH is a rare condition, the model was unable to differentiate between the different surgery types offered in IIH:WT due to the reduced sample size within each group, so any differential costs and effects were not included. There is evidence that LAGB is associated with lower procedure costs, but has a much higher rate of revisional surgery, as well as a smaller and less well-maintained effect on body weight than RYGB and LSG [28]. However, the range and distribution of bariatric surgeries performed in the IIH:WT broadly reflect current practice in the English healthcare system [15,21], and therefore the results are applicable to assessing surgery versus community weight management for treatment of IIH. At an international level, more research is needed to fully estimate the differential cost-effectiveness between the surgery types. Within the IIH:WT, only half of the community weight management cohort attended 75%–100% of weekly sessions, but evidence suggests this is similar attendance to that seen in other trials [29], and it is unknown how adherence would have varied outside the trial setting. The model used data from the IIH:WT up to the end of cycle 2; beyond that, data from the SOS study were used, as this study contained some of the longest follow-up data available on bariatric surgery. However, this study was conducted 16 years ago, was based exclusively in Sweden, and the surgical techniques differ, as the SOS study also used vertical banded gastroplasty, which is no longer performed; therefore, surgical outcomes may have since changed. Alternative sources of long-term data were available, including a meta-analysis [30]; however, the data were reported as percentages of expected weight loss and the economic model tracked weight trajectories using BMI categories, making it challenging to use these data for the model structure. Within the SOS study, vertical banded gastroplasty weight loss at 10 years was similar to LABG weight loss, and we know that LSG is more effective than LAGB at achieving long-term weight loss; therefore, by using the SOS data to populate the model, we believe this to be a conservative estimate for the likely long-term effectiveness for the surgical arm.

Conclusion

The results suggest that bariatric surgery is a dominant treatment option for IIH patients living with obesity when compared to a community weight management intervention. It provides evidence to inform funding bodies that IIH should qualify as a co-morbidity of obesity that can be improved with weight loss. Hence, IIH patients with a BMI over 35 kg/m² should meet the criteria to be recommended for bariatric surgery under NICE clinical guidance [17].

Acknowledgments

We acknowledge Birmingham Clinical Trials Unit (BCTU) for trial coordination and data management. We acknowledge support from the Medical Research Council (MRC UK) and the National Institute for Health Research (NIHR UK) who funded the IIH:WT trial, the NIHR Clinical Research Network and the Wellcome Trust Clinical Research Facilities, where IIH:WT was performed.

Disclosures

This Trial was funded by grant NIHR-CS-001-028 (Clinician Scientist Fellowship) from the National Institute for Health Research (Dr Sinclair) and grant MR/K015184/1 from the Medical Research Council UK (Dr Sinclair). EF was funded by a National Institute for Health Research (NIHR) career development fellowship award (NIHR-CDF-2015-08-13) for the duration of the study. AJS was funded by an NIHR clinician scientist fellowship (NIHR-CS-011-028) for the duration of the study; is funded by a Sir Jules Thorn Award for Biomedical Science; and reports personal fees from Invex therapeutics, during the conduct of the study but outside the submitted work. AAT was funded by an NIHR Clinician Scientist Award for part of the duration of the study (CS-2013-13-029) and reports grants from Novo Nordisk, personal fees from Novo Nordisk, nonfinancial support from Novo Nordisk, personal fees from Eli Lilly, nonfinancial support from Eli Lilly, personal fees from Janssen, personal fees from AZ, nonfinancial support from AZ, nonfinancial support from Impeto medical, nonfinancial support from Resmed, nonfinancial support from Aptiva, personal fees from BI, nonfinancial support from BI, personal fees from BMS, nonfinancial support from BMS, personal fees from NAPP, nonfinancial support from NAPP, personal fees from MSD, nonfinancial support from MSD, grants from Sanofi, and personal fees from Sanofi. SPM reports other Invex Therapeutics, other Heidelberg engineering during the conduct of the study; other from Chugai-Roche Ltd, other from Janssen, other from Allergan, other from Santen, other from Roche, and other from Neurodiem, outside the submitted work. OG reports consulting work for Invex Therapeutics during the conduct of the study but outside the submitted work. BRW reports consulting work for Invex Therapeutics during the conduct of the study but outside the submitted work. All other authors declare no competing interests. The views expressed are those of the authors and not necessarily those of the UK National Health Service, the NIHR, or the UK department of Health and Social Care.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.soard.2021.03.020>.

References

- [1] Mollan S, Grech O, Alimajstorovic Z, et al. New horizons for idiopathic intracranial hypertension: advances and challenges. *Brit Med Bull* 2020;136:118–26.
- [2] Virdee J, Lacombe S, Vijay V, et al. Reviewing the recent developments in idiopathic intracranial hypertension. *Ophthalmol Ther* 2020;9:767–81.
- [3] Mollan SP, Aguiar M, Evison F, et al. The expanding burden of idiopathic intracranial hypertension. *Eye* 2019;33:478–85.
- [4] Piper RJ, Kalyvas AV, Young AMH, et al. Interventions for idiopathic intracranial hypertension. *Cochrane Database Syst Rev* 2015;2015(8):CD003434.
- [5] Mollan SP, Hoffmann J, Sinclair AJ. Advances in the understanding of headache in idiopathic intracranial hypertension. *Curr Opin Neurol* 2019;32:92–8.
- [6] Daniels A, Liu G, Volpe N, et al. Profiles of obesity, weight gain, and quality of life in idiopathic intracranial hypertension (pseudotumor cerebri). *Am J Ophthalmol* 2007;143:635–41.
- [7] Hornby C, Mollan SP, Botfield H, et al. Metabolic concepts in idiopathic intracranial hypertension and their potential for therapeutic intervention. *J Neuroophthalmol* 2018;38:522–30.
- [8] Mollan SP, Hornby C, Mitchell J, et al. Evaluation and management of adult idiopathic intracranial hypertension. *Pract Neurol* 2018;18:485–8.
- [9] Adderley N, Subramanian A, Nirantharakuman K. Association between idiopathic intracranial hypertension and risk of cardiovascular diseases in women in the United Kingdom. *JAMA Neurol* 2019;76:1088–98.
- [10] Friesner D, Rosenman R, Lobb BM, et al. Idiopathic intracranial hypertension in the USA: the role of obesity in establishing prevalence and healthcare costs. *Obes Rev* 2011;12:e372–380.
- [11] Sinclair AJ, Burdon MA, Nightingale PG, et al. Low energy diet and intracranial pressure in women with idiopathic intracranial hypertension: prospective cohort study. *BMJ* 2010;341:c2701.
- [12] Mollan SP, Davies B, Silver NC, et al. Idiopathic intracranial hypertension: consensus guidelines on management. *J Neurol Neurosurg Psychiatry* 2018;89:1088–100.
- [13] Sun WYL, Switzer NJ, Dang JT, et al. Idiopathic intracranial hypertension and bariatric surgery: a systematic review. *Can J Surg* 2020;63:E123–8.
- [14] Sjöström L, Narbro K, Sjöström CD, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med* 2007;357:741–52.
- [15] Ottridge R, Mollan S, Botfield H, et al. Randomised controlled trial of bariatric surgery versus a community weight loss programme for the sustained treatment of idiopathic intracranial hypertension: the Idiopathic Intracranial Hypertension Weight Trial (IIH:WT) protocol. *BMJ Open* 2017;7:e017426.
- [16] Mollan S, Mitchell J, Ottridge R, et al. Bariatric surgery versus community weight management intervention for the treatment of idiopathic intracranial hypertension (IIH:WT): a randomized controlled trial. *JAMA Neurol*. Epub 2021 April 26. <https://doi.org/10.1001/jamaneruol.2021.0659>.
- [17] National Institute for Health and Care Excellence [homepage on the Internet]. Obesity: identification, assessment and management: clinical guideline [CG189]. c2014 [cited 18 March, 2021]. Available from: <https://www.nice.org.uk/guidance/cg189>.
- [18] Sjöström L. Review of the key results from the Swedish Obese Subjects (SOS) trial—a prospective controlled intervention study of bariatric surgery. *J Intern Med* 2013;273:219–34.
- [19] Sjöström L, Lindroos A-K, Peltonen M, et al. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *N Engl J Med* 2004;351:2683–93.
- [20] Berrington de Gonzalez A, Hartge P, Cerhan JR, et al. Body-mass index and mortality among 1.46 million white adults. *N Engl J Med* 2010;363:2211–9.
- [21] Welbourn R, Small P, Finlay I. National bariatric surgery register, second registry report. 2014. Dendrite Clinical Systems Ltd: Oxfordshire. Available from: <https://e-dendrite.com/Publishing/Reports/Bariatric/NBSR2014.pdf>.
- [22] NHS Improvement. National tariff payment system 2019/20: a consultation notice. Available from: <https://improvement.nhs.uk/resources/national-tariff-1920-consultation/>. Accessed 18 March, 2021.
- [23] Curtis LA, Burns A. Unit Costs of Health and Social Care 2017, Personal Social Services Research Unit, University of Kent: Canterbury. <https://doi.org/10.22024/UniKent/01.02/65559>.
- [24] Thaller M, Tsermoulas G, Sun R, Mollan SP, Sinclair AJ. The negative impact of COVID-19 lockdown on papilloedema and idiopathic intracranial hypertension. *J Neurol Neurosurg Psychiatry*. Epub 2020 Dec 24. <https://doi.org/10.1136/jnnp-2020-325519>.
- [25] Ko MW, Chang SC, Ridha MA, et al. Weight gain and recurrence in idiopathic intracranial hypertension: a case-control study. *Neurology* 2011;76:1564–7.
- [26] McTigue K, Larson JC, Valoski A, et al. Mortality and cardiac and vascular outcomes in extremely obese women. *JAMA* 2006;296:79–86.
- [27] Hazlehurst JM, Logue J, Parretti HM, et al. Developing integrated clinical pathways for the management of clinically severe adult obesity: a critique of NHS England policy. *Curr Obes Rep* 2020;9:530–43.
- [28] Gulliford MC, Charlton J, Prevost T, et al. Costs and outcomes of increasing access to bariatric surgery: cohort study and cost-effectiveness analysis using electronic health records. *Value Health* 2017;20:85–92.
- [29] Dixon K, Shcherba S, Kipping R. Weight loss from three commercial providers of NHS primary care slimming on referral in North Somerset: service evaluation. *J Pub Health* 2012;34:555–61.
- [30] O'Brien PE, Hindle A, Brennan L, et al. Long-term outcomes after bariatric surgery: a systematic review and meta-analysis of weight loss at 10 or more years for all bariatric procedures and a single-centre review of 20-year outcomes after adjustable gastric banding. *Obes Surg* 2019;29:3–14.