

## The other side of the coin: Blunted cardiovascular and cortisol reactivity are associated with negative health outcomes.

Phillips, Anna; Ginty, Annie; Hughes, Brian

DOI:

[10.1016/j.ijpsycho.2013.02.002](https://doi.org/10.1016/j.ijpsycho.2013.02.002)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Phillips, A, Ginty, A & Hughes, B 2013, 'The other side of the coin: Blunted cardiovascular and cortisol reactivity are associated with negative health outcomes.', *International journal of psychophysiology : official journal of the International Organization of Psychophysiology*, vol. 90, no. 1, pp. 1-7.

<https://doi.org/10.1016/j.ijpsycho.2013.02.002>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

NOTICE: this is the author's version of a work that was accepted for publication in International Journal of Psychophysiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in International Journal of Psychophysiology, VOL 90, ISSUE 1, (2013)

DOI: 10.1016/j.ijpsycho.2013.02.002

Eligibility for repository checked February 2015

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## **The other side of the coin: Blunted cardiovascular and cortisol reactivity are associated with negative health outcomes.**

### **Introductory Paper**

Anna C. Phillips<sup>1</sup>, Annie T. Ginty<sup>1</sup>, and Brian M. Hughes<sup>2</sup>.

<sup>1</sup>School of Sport & Exercise Sciences, College of Life and Environmental Sciences, University of Birmingham, UK

<sup>2</sup>School of Psychology, National University of Ireland, Galway, Ireland

#### Highlights

- Blunted reactions to acute laboratory stress relate to poor health outcomes.
- Blunting may reflect reduced effort, stress perceptions, or physiological capacity.
- Blunted responses may be a marker of central motivational dysregulation.
- Blunted reactivity is neurally based and independent of stress task effort.

Address correspondence to: Anna C. Phillips, PhD, School of Sport and Exercise Sciences, University of Birmingham, Birmingham B15 2TT, England. Tel 0044 121 414398; Fax 0044 121 4144121; e-mail [A.C.Phillips@bham.ac.uk](mailto:A.C.Phillips@bham.ac.uk)

## **Abstract**

A cornerstone for research into the link between stress and health has been the reactivity hypothesis; cardiovascular reactivity to psychological stressors, if prolonged or exaggerated, can promote the development of cardiovascular disease. However, it has recently been argued that low or blunted reactivity is also associated with negative health outcomes. As such, in this special issue we present further evidence implicating that cardiovascular and stress hormone responses to acute stress at the other end of the response spectrum can also be considered a pathway to ill health. In this introductory article, we explore and review the origins of and potential mechanisms underlying blunted responses to acute stress. In so doing, we aim to highlight: what is currently known regarding this new conceptualization of the reactivity hypothesis; the potential explanations for blunted reactivity; the pathways underlying associations with health outcomes; and where this field is headed in terms of developing our understanding of the link between reactivity and health.

**Keywords:** blunted reactivity; cardiovascular reactivity; health outcomes; stress responses;

## 1.1 Introduction

A key focus of research into the link between stress and health for several decades has been the reactivity hypothesis, which proposes that cardiovascular reactivity to psychological stressors, if prolonged or exaggerated, can promote the development of cardiovascular disease (Obrist, 1981). Heightened cardiovascular reactivity is posited to result in changes to the structure and functioning of the heart that potentially promote a number of adverse disease outcomes, including sustained hypertension (Obrist, 1981) cardiac and vascular hypertrophy (Lovallo and Gerin, 2003), oxidation of low-density lipoproteins (Raitakari et al., 1997), and increased serum concentrations of both pro-inflammatory cytokines (Georgiades, 2007) and blood insulin (Nazzaro et al., 2002). Theoretical considerations of how negative health results from reactivity-related adaptations of the cardiovascular system have been corroborated by an extensive empirical literature linking heightened reactivity to objective outcomes, including hypertension (Carroll et al., 2003; Carroll et al., 1995; Carroll et al., 2001; Everson et al., 1996; Markovitz et al., 1998; Matthews et al., 1993; Newman et al., 1999; Treiber et al., 1997), atherosclerosis (Barnett et al., 1997; Everson et al., 1997; Lynch et al., 1998; Matthews et al., 1998), increased left ventricular mass and/or hypertrophy of the heart (Georgiades et al., 1997; Kapuku et al., 1999; Murdison et al., 1998), and cardiovascular disease mortality (Carroll et al., 2012b). Both qualitative reviews and meta-analyses of this evidence confirm the contention that exaggerated stress reactions signal poor future cardiovascular health (Chida and Steptoe, 2010; Gerin et al., 2000) (Schwartz et al., 2003; Taylor et al., 2003) (Treiber et al., 2003).

Given the emphasis placed on associations between exaggerated reactivity and disease pathogenesis, low or blunted reactivity to acute stress has, by implication, been assumed to be

benign or even protective. However, recent evidence suggests that low cardiovascular reactivity to stress, as well as low cortisol reactivity, may actually have serious adverse consequences for health and behaviour. For example, comparatively low cardiovascular and cortisol reactions to acute psychological stress have been found to characterize smokers (al'Absi, 2006; al'Absi et al., 2005; Girdler et al., 1997; Phillips et al., 2009b; Roy et al., 1994; Sheffield et al., 1997) and those with alcohol and other substance addictions (Brenner and Beauchaine, 2011; Lovallo, 2005; Lovallo et al., 2000; Panknin et al., 2002), as well as being associated with obesity (Carroll et al., 2008), depression (Brinkmann et al., 2009; Carroll et al., 2007; de Rooij et al., 2010; Phillips et al., 2011; Rottenberg et al., 2007; Salomon et al., 2009a; Schwerdtfeger and Rosenkaimer, 2011; York et al., 2007), poor self-reported health (De Rooij and Roseboom, 2010; Phillips et al., 2009a), exercise addiction (Heaney et al., 2011a), eating disorders (Ginty et al., 2012a), and poorer cognitive function (Ginty et al., 2011a; Ginty et al., 2011b; Ginty et al., 2012b). Comparatively reduced cardiovascular stress responses have also been associated with personality traits also indicative of future disease, such as Type D personality (Howard et al., 2011) and neuroticism (Hughes et al., 2011), and other studies have found low reactivity among people taking medications known to increase heart disease risk, such as oral contraceptives (Schallmayer and Hughes). In light of accumulating evidence supporting an association between blunted stress reactivity and addiction, and the emerging evidence relating blunted reactivity to other unhealthy behaviours and negative health outcomes, it would appear that both extremes, exaggerated and diminished reactivity are maladaptive responses to stress. This suggests that the most optimally healthy response to stress is a moderate reaction (Lovallo, 2011). This paper will examine the concept of blunted responses to stress, their possible origins and underlying mechanisms as an introduction to this special issue on blunted reactivity to acute stress.

## 2.1. What are the bases of 'blunted' reactivity?

We take the term 'blunted cardiovascular reactivity' to refer to an empirically demonstrable cardiovascular response pattern which is comparatively lower than that seen during typical states of homeostatic function during stress. As yet, the precise mechanism that determines the occurrence of blunting is unclear, and an important distinction needs to be made between reactivity that is blunted (which implies sub-normality) and reactivity that is simply low (which may be biologically normal within a given context). There are several possible bases of blunted reactivity, which we outline below. Overall, it can be noted that while there are a number of possibilities, several remain to be tested empirically.

### 2.1.1 Lower effort

One possibility is that lower reactions to acute stress tasks may reflect *lower effort* on the part of a participant; in other words, that blunted responses are primarily the result of behavioral factors rather than of cognitive or biological factors. Perceptions of how stressful a psychological stress task is might impact upon an individual's willingness to engage with the task, with reduced effort or motivation then underlying a negative health impact. It could be argued that symptoms of conditions such as depression, which are associated with blunted responses, might also relate to lower motivation levels. Certainly, depression is characterized by a reduced degree of motivation, and both depression and low motivation appear to be related to the same gene polymorphisms, such as the met variant of the Val158Met COMT gene (Aberg et al., 2011), which results in different levels of extracellular dopamine within key brain reward system areas. Yet, in neuropsychological tests requiring cognitive effort, depressed participants have not consistently performed worse (Hammar et al., 2011). Further, not all of the behaviours or conditions associated with blunted responses to stress relate to lower motivation or effort; indeed

some such as exercise dependence (Heaney et al., 2011b) require considerable motivation and effort. Finally, in studies produced by the Birmingham group, we have always found blunted responses to be independent of ratings of task stressfulness and engagement, as well as objective performance scores which are a proxy for task engagement (see e.g. (Ginty et al., 2012a; Heaney et al., 2011a)). Taken together, this evidence suggests that individuals characterised by blunted responses are not necessarily demonstrating lower task performance or perceptions of motivation. However, as performance is merely a proxy for effort, and not the same construct, it remains unclear whether or not the tendency to invest less effort accounts for trait-like patterns of blunted responses.

#### 2.1.2 Reduced awareness or perception of stress

An alternative theory of blunted reactivity is that it might reflect a *reduced awareness or perception of stress*; in other words, that blunted responses is primarily the result of cognitive factors rather than of behavioural or biological factors. An inability to detect stressors in the environment, or a tendency to view dangerous stimuli as innocuous, would logically serve to dampen physiological stress responses. It has long been established (a) that perceptual factors are important in determining whether an individual exhibits a physiological stress response (Speisman et al., 1964) and (b) that individuals differ in the extent to which they habitually regard otherwise innocuous stimuli as negatively valenced (Bishop, 2008). It has also been established that training participants to bias their attention away from negative stimuli serves to dampen their cardiovascular responses to subsequent stressors (Higgins and Hughes, 2012), implying that such perceptions are causally responsible for cardiovascular response profiles rather than the other way around. Moreover, such effects are most pronounced among persons with high levels of trait neuroticism (Connor-Smith and Flachsbart, 2007), a group also

characterised by blunted responses to stress (Phillips et al., 2005a). As such, blunted cardiovascular responses may reflect diminutions in stress vigilance and/or detection, although this remains to be tested directly.

### 2.1.3 The impact of task difficulty

One possibility underlying blunted reactivity is that of the impact of task difficulty. In stress tasks which employ active coping, it has been shown that the extent of reactivity is related to the difficulty or challenge levels of the task, such that both easy and over-challenging tasks can result in weaker cardiovascular responses (Richter and Gendolla, 2006). When task difficulty is manipulated, as it increases, so too do the magnitude of cardiovascular responses until difficulty levels where success is impossible (Carroll et al., 1986; Richter et al., 2008), although this has not been shown in all studies (Willemsen et al., 2000). This is considered to be due to effects on motivational intensity in active coping situations (Wright, 1996). Task difficulty is also thought to interact with mood resulting in different levels of perceived subjective demand, which in turn determine the amplitude of cardiovascular reactivity (Gendolla and Krusken, 2001). Indeed these situational manipulations of perceived task demand have been replicated multiple times to show the impact of effort on cardiovascular responses to stress. However, other studies have revealed that in situations of matched task difficulty (Phillips, 2011), and across different types of tasks (de Rooij, 2013 in press) in this special issue), there appear to be a sub-group of individuals who consistently show trait-like blunted responses to acute stress, independent of perceptions of tasks difficulty, see e.g., (Ginty et al., 2012a; Heaney et al., 2011a). This brings us to the next potential explanation for blunting.

### 2.1.4 Reduced physiological capacity to respond

A fourth explanation for blunted reactivity is that the blunted responses may reflect a reduced physiological capacity to respond to stress or a disengagement of the biological systems, rather than of behavioural or cognitive factors. For example, there is reasonably consistent evidence that the sympathetic nervous systems of individuals who have become obese, although characterized by high basal sympathetic nervous system activity (Carroll et al., 2008; Tentolouris et al., 2006), and higher basal cortisol levels (Bjorntorp, 1993; Bjorntorp and Rosmond, 2000), are less responsive to stimulation (Carroll et al., 2008; Tentolouris et al., 2006). There is evidence of a postprandial sympathetic nervous system response, as indicated by higher plasma norepinephrine concentrations and an increased low to high frequency ratio in the HR variability spectrum after ingestion of a meal (Tentolouris et al., 2003; Welle et al., 1981). Similarly, this is smaller among obese individuals than the non-obese (Tentolouris et al., 2003). In addition, the changes in HR and muscle sympathetic nerve stimulation following infusion of anti-hypertensive and anti-hypotensive drugs are significantly smaller in the obese than the non-obese (Grassi et al., 1995), consistent with the view that such blunted responses result from biologically-based attenuation. A further example is that of depression; where basal sympathetic nervous system activity, as indexed by a shift enhanced cardiac sympathetic activity relative to vagal tone (Carney et al., 1988), increased plasma noradrenaline concentrations (Rudorfer et al., 1985), and increased 24-hour urinary noradrenaline excretion (Hughes et al., 2004) has been shown to be higher in individuals with depression or depressive symptomatology compared to controls. This perhaps also makes depressed individuals less responsive to stimulation as shown in reactivity studies (Brinkmann and Gendolla, 2008; Carroll et al., 2007; de Rooij et al., 2010; Phillips et al., 2011; Salomon et al., 2009b; York et al., 2007), potentially via decreased  $\beta$ -adrenergic receptor responsiveness (Mazzola-Pomietto et al., 1994; Yu et al., 2008).

### 2.1.5 Motivational dysregulation within the brain

In line with this argument of a reduced physiological capacity to respond to stress, it has previously been proposed that blunted physiological stress reactivity may be a marker of central motivational dysregulation (Carroll et al., 2009, 2011), a suboptimal functioning of the physiological systems in the brain that support motivation and motivated behavior (e.g. a disengagement in the motivational areas of the brain when faced with an acute challenge). Support comes from the evidence that blunted physiological responses to acute stress are associated with addictions (al'Absi, 2006) (Girdler et al., 1997; Lovallo, 2005) and with a non-substance abuse addiction and disorder, bulimia (Ginty et al., 2012a; Koo-Loeb et al., 1998), related to hypo-functioning of the brain in response to reward (Bohon and Stice). Additionally, individuals with obesity have been observed to show reduced responses to food within the striatum (Stice et al., 2008). Persons with depression are characterised by reduced activation of the striatum to positive stimuli (Epstein et al., 2006) and reduced responsiveness to reward behaviourally through weaker emotional responses, neurologically in the frontostriatal systems, and neurochemically through dysregulated function in brain areas under the control of monoamines (Eshel and Roiser, 2010). Additional support comes from evidence that blunted reactions are associated with poorer cognitive ability in earlier life (Ginty et al., 2011a) and both cross-sectionally (Ginty et al., 2012b) and prospectively (Ginty et al., 2011b); cognitive performance requires the integrity of neural motivational systems (Busato et al., 2000; Dweck, 1986; McClelland et al., 1953; Pintrich and Schunk, 1986). Perhaps the most compelling evidence of motivational dysregulation comes from a recent study showing a hypo-activation during stress in blunted reactors compared to exaggerated reactors. This hypo-activation was concentrated in the anterior mid-cingulate cortex: an area of the brain associated with motivation,

high cognitive functioning, engagement, and emotional control. This study also demonstrated a deactivation during stress compared to rest in the amygdala and frontal cortex: areas of the brain associated with motivation and emotion and blunted reactors reported similar levels of stress task engagement, difficulty, and stressfulness (Ginty et al., 2013) . Thus it is plausible that blunted responses to acute stress situations might reflect unconscious reduced effort (i.e., where a person believes they are trying as hard as everyone else), as is suggested by our previously observed post-task ratings of effort and engagement as well as individuals' stress task performance scores which indicate effort (Ginty et al., 2012a). Indeed, we have recently shown that individuals with inadvertent reduced motivation, measured through forced expiratory volume, are also characterised by blunted stress reactions (Carroll et al., 2012a).

#### 2.1.6 Stifling or titration of the blood pressure response

A final scenario to account for blunting is where the individual detects and recognises stressors, is adequately motivated to respond, and duly initiates a orthodox physiological stress response that is then *stifled or dampened by (or titrated into) the activity of some other physiological response system*. For example, a cardiovascular stress response may be immediately interrupted and attenuated by a competing demand that serves to reduce blood pressure and heart rate. In relation to psychological stress, it is possible that cardiovascular and cortisol responses to stress are sometimes counterbalanced by competing physiological demands arising from emotional (e.g., fear, threat) or bodily (fatigue, hunger) determinants. For example, chronic sleep loss is associated with disruption to blood pressure (e.g., (Kotani et al., 2008)), and so will interfere with measures of cardiovascular stress responses. This in turn creates a likely complication in studies of cardiovascular function in populations whose characteristics are themselves associated with sleep disruption, such as depression (Benca and Peterson, 2008) and obesity (Vgontzas et

al., 2009), both of which have been implicated in cardiovascular blunting. Even within the cardiovascular response system alone, such competing demands play out in the central dynamic homeostatic relationship that exists between cardiac and vascular variables (James et al.). Changes in cardiac output (CO) can often be compensated for by corresponding inverse changes in total peripheral resistance (TPR), without elevating systolic or diastolic blood pressure. Accordingly, even when it appears as though a stress stimulus is eliciting a 'blunted' blood pressure response, the cardiovascular system overall may well be experiencing significant disruption in terms of shifts in CO and TPR. Transitioning from a myocardial (i.e., CO-driven) to a vascular (i.e., TPR-driven) response profile, while maintaining homeostasis in terms of blood pressure stability, is nonetheless indicative of disease risk (Palatini and Julius, 2009). A number of studies have suggested that persons who exhibit low (or blunted) systolic blood pressure reactivity tend to exhibit maladaptive haemodynamic profiles, such as shifting towards vascular-driven responses when experiencing archetypically myocardial stressors (e.g., (Hughes et al., 2011)). Such possibilities highlight the importance of multivariate and multi-system analyses of physiological stress responses, both in general and with reference to blunting in particular.

Given these different potential reasons for blunted reactivity, and the uncertainty as to which might be the most likely explanation or set of explanations, the present special issue attempts to bring about an awareness of this issue by focusing on blunted responses to acute stress and their correlates in order to facilitate progress in developing an integrative explanation for these findings across studies from a variety of sources and populations.

### **3.1 Where does blunted reactivity come from?**

### 3.1.1 Prior events

The question of how people come to exhibit blunted reactivity can be considered with regard to both proximal and distal explanations. In the case of individual lifespan development, it may be that blunted cardiovascular reactivity emerges as the result of *prior psychological or physiological events*. It was originally thought that high biological reactivity to stressors might develop from highly stressful childhood social environments (Boyce and Ellis, 2005). For example, children raised in homes with low levels of organization and consistency show an elevated cortisol response to social stress (Ellenbogen and Hodgins, 2009). Further, in a study of children exposed to highly stressful environments in early childhood, sympathetic and adrenocortical reactivity was higher in comparison to children from moderately stressful environments (Ellis et al., 2005) although children in very supportive low stress environments also displayed high autonomic reactivity also (Ellis et al., 2005). Relatedly, babies whose mothers smoked also displayed heightened blood pressure and heart rate responses to tilt stress (Cohen et al., 2008). The animal literature on early life stress (maternal separation) on reactivity also supports these findings (Sanders and Anticevic, 2007). However, more recently, it has been shown that adults who had experienced significant adverse life events in childhood display smaller heart rate and cortisol responses to mental arithmetic and public speaking stress (Lovallo et al., 2012). Similarly, children who had experienced the stress of maltreatment or bullying showed lower cortisol responses to public speaking stress (Ouellet-Morin et al., 2011). Additionally, young adults with a history of negative family relationships who display hostile verbal behaviours during a role play task are characterised by attenuated HR responses to the task (Luecken and Roubinov, 2012). Finally, in older adults, those who had experienced more past stressful life events in childhood showed lower cortisol responses to the Trier Social Stress

Test (Armbruster et al., 2011). This is also in line with studies which have shown that prior stressful life events and increased perceptions of stressfulness of those life events is related to attenuated cardiovascular reactivity to acute psychological stress (Carroll et al., 2005; Phillips et al., 2005b). It is possible that the differences between studies in terms of early life stress predicting high or low reactivity depends on the age of the individuals undergoing reactivity testing, as many of the studies showing blunting are in adults or elders, which may suggest some type of adaptation of an initially large response in childhood to a blunted response in later life. This brings us on to our next possibility for where blunting comes from.

### 3.1.2 Adaptation

Consistent with some of the longest standing theories in biological psychology (e.g., the flight-or-fight response theory (Cannon, 1932), the cardiovascular stress response can be presumed to be a mechanism that has been naturally selected on the basis of adaptive pressures. As with other developed psychological mechanisms (such as pain detection, anxiety, or depression), one implication of this is that function of mid-range intensity is likely to be the most adaptive, with function at the extremes likely to be less so. As such, within a standard evolutionary framework, it is unsurprising that blunted cardiovascular stress responding (as well as exaggerated responding) is found to be associated with adverse outcomes for the individual organism. Of course, the observed range of cardiovascular reactivity to stress will be influenced by the balancing of adaptive and maladaptive features. Although blunted cardiovascular responses might well have adverse health consequences, they may also confer some ephemeral benefits that result in their being retained in the gene pool. Some initial studies have suggested that limited emotional response states, such as alexithymia, may in fact assist individuals in short-term coping, stress avoidance, and concentration, as well as having some beneficial socially

communicative effects (Nesse and Ellsworth, 2009). In this way, blunted cardiovascular and cortisol reactivity may comprise highly nuanced and developed elements of the physiological response to stress.

#### 4.1 Towards an understanding of blunting: Looking at the other side of the coin

Many psychological, social, and behavioural factors, such as depression, social support, early life adversity, personality, and addictions, have been examined in relation to stress reactivity, and as indicated above, many are emerging as correlates of blunted reactivity. Indeed, it may be that physiological blunted responses are manifest in particular behaviours or personality traits. This Special Issue of the *International Journal of Psychophysiology* provides an opportunity to examine in further depth the range of behavioural and personality manifestations associated with blunted responding, and their underlying commonalities. Some of these personality and behavioural characteristics are also evident among some of the health-related correlates of blunted reactivity, such as bulimia (Ginty et al., 2012a), and exercise dependence (Heaney et al., 2011a).

Knowing what is already established about the negative health outcomes of exaggerated reactivity to stress, it would appear that depending on the outcomes in question, departures from the norm of physiological responding in either direction may pose problems. This suggests that in both instances the system is operating in a biased state, whether at the level of the higher CNS, at the level of the hypothalamus and brainstem, or at the level of the periphery (Carroll et al., 2009). This can be conceptualised as an inverted-U model where high and low reactivity are bad for health depending on the health outcome in question (Carroll et al., 2009) as displayed in Figure 1.

Given their reliance on biological, psychological, and social factors, reactivity paradigms are inherently complex and many issues remain unclear. In order to fully understand the associations between reactivity and health, many important pathways need to be considered in more detail, including the associations between personality effects on motivation and stress responses, cognition and reactivity, and neural correlates of both lowered reactivity and reduced motivation. Understanding these multifaceted links in more detail will enable us to develop greater insight into the routes to risky health behaviours and resultant poor health, and indeed provide the potential for early detection and intervention. For example, blunted stress reactions predict relapse in smokers who have quit (al'Absi et al., 2005) and are also evident in the adolescent offspring of alcoholic parents (Sorocco et al., 2006), suggesting that blunted stress reactivity may have substantial prognostic value.


### 5.1 This special issue

In this special issue, we explore the latest cutting edge research on the psychosocial and behavioural correlates of blunted reactivity and the potential biological and neural pathways by which these factors might have implications for health and disease. In so doing, we aim to highlight what is currently known regarding this new conceptualization of the reactivity hypothesis and the different theories of blunted reactivity, as outlined above, in order to facilitate understanding of what might be the most likely explanation for this phenomenon and where it emerges from. For example, the extent to which of a variety of negative and positive personality traits and mental health disorders or their symptoms are associated with blunted reactivity will be examined by Bibbey et al. (Bibbey et al., 2013 in press) Brindle et al. (Brindle et al., 2013, under

revision), O'Leary et al., (O'Leary et al., 2013 in press) Kupper et al. (Kupper, 2013 in press) Salomon et al. (Salomon et al., 2013 in press) and Schwerdtfeger and Gerteis (Schwerdtfeger and Gerteis). The role that perceptions of stress, behavioural approach/effort, and mood state play in the magnitude of stress reactivity will be examined by Brindle et al. (Brindle et al., 2013, under revision), Schwerdtfeger and Gerteis (Schwerdtfeger and Gerteis), and Salomon et al. (Salomon et al., 2013 in press) respectively. We will also examine the role of early life stress exposure and trauma on the development of blunted responses in papers by Lovallo et al., (Lovallo, 2013 in press) and D'Andrea (D'Andrea, under review), respectively. Further, psychosocial, cognitive, and health correlates of blunted responses and the mechanisms underlying these associations will be considered by Carroll et al. (Carroll et al., 2013 in press), O'Leary et al. (O'Leary et al., 2013 in press), de Rooij (de Rooij, 2013 in press), and Lovallo et al. (Lovallo, 2013 in press), including whether or not blunting can be experimentally induced (O'Leary et al., 2013 in press). Finally, we will address where this field is headed in terms of developing our understanding of the link between reactivity and health through a paper by Ginty (Ginty, 2013 under review) integrating blunted reactivity findings with theory regarding blunted responses to reward, and finally through the integrative commentary by Allen (Allen, 2013, in review). As such, this special issue seeks to expand upon current and previous reactivity literature, so that the contribution of the reactivity hypothesis to explaining the health consequences of key psychosocial and behavioural factors can reach its full potential.

Figure 1: A proposed model linking low and high reactivity to different health outcomes.

Adapted from (Carroll et al., 2009) with permission.



## References

- Aberg, E., Fandino-Losada, A., Sjöholm, L.K., Forsell, Y., Lavebratt, C., 2011. The functional Val158Met polymorphism in catechol-O-methyltransferase (COMT) is associated with depression and motivation in men from a Swedish population-based study. *J Affect Disord* 129, 158-166.
- al'Absi, M., 2006. Hypothalamic-pituitary-adrenocortical responses to psychological stress and risk for smoking relapse. *Int J Psychophysiol* 59, 218-227.
- al'Absi, M., Hatsukami, D., Davis, G.L., 2005. Attenuated adrenocorticotrophic responses to psychological stress are associated with early smoking relapse. *Psychopharmacology (Berl)* 181, 107-117.
- Allen, M., 2013, in review. Integrative commentary: Implications of blunted reactivity. *International Journal of Psychophysiology*.
- Armbruster, D., Mueller, A., Strobel, A., Lesch, K.P., Brocke, B., Kirschbaum, C., 2011. Predicting cortisol stress responses in older individuals: influence of serotonin receptor 1A gene (HTR1A) and stressful life events. *Horm Behav* 60, 105-111.
- Barnett, P.A., Spence, J.D., Manuck, S.B., Jennings, J.R., 1997. Psychological stress and the progression of carotid artery disease. *J Hypertens* 15, 49-55.
- Benca, R.M., Peterson, M.J., 2008. Insomnia and depression. *Sleep Med* 9 Suppl 1, S3-9.
- Bibbey, A., Carroll, D., Roseboom, T.J., Phillips, A.C., de Rooij, S.R., 2013 in press. Personality and physiological reactions to acute psychological stress. *Int J Psychophysiol*.
- Bishop, S.J., 2008. Neural mechanisms underlying selective attention to threat. *Ann N Y Acad Sci* 1129, 141-152.
- Bjorntorp, P., 1993. Visceral obesity: a "civilization syndrome". *Obes Res* 1, 206-222.
- Bjorntorp, P., Rosmond, R., 2000. Obesity and cortisol. *Nutrition* 16, 924-936.
- Bohon, C., Stice, E., 2011. Reward abnormalities among women with full and subthreshold bulimia nervosa: a functional magnetic resonance imaging study. *Int J Eat Disord* 44, 585-595.
- Boyce, W.T., Ellis, B.J., 2005. Biological sensitivity to context: I. An evolutionary-developmental theory of the origins and functions of stress reactivity. *Dev Psychopathol* 17, 271-301.
- Brenner, S.L., Beauchaine, T.P., 2011. Pre-ejection period reactivity and psychiatric comorbidity prospectively predict substance use initiation among middle-schoolers: A pilot study. *Psychophysiology* 48, 1588-1596.
- Brindle, R., Terrill, J.A., Ginty, A.T., Conklin, S.M., 2013, under revision. Is the association between depression and blunted cardiovascular stress reactions mediated by perceptions of stress? *International Journal of Psychophysiology*.
- Brinkmann, K., Gendolla, G.H., 2008. Does depression interfere with effort mobilization? Effects of dysphoria and task difficulty on cardiovascular response. *J Pers Soc Psychol* 94, 146-157.
- Brinkmann, K., Schupbach, L., Joye, I.A., Gendolla, G.H., 2009. Anhedonia and effort mobilization in dysphoria: reduced cardiovascular response to reward and punishment. *Int J Psychophysiol* 74, 250-258.

- Busato, V.V., Prins, F.J., Elshout, J.J., Hanmaker, C., 2000. Intellectual ability, learning style, personality, achievement motivation, and academic success of psychology students in higher education. *Personality and Individual Differences* 29, 1057-1068.
- Cannon, W.B., 1932. *The wisdom of the body*. Norton, New York.
- Carney, R.M., Rich, M.W., teVelde, A., Saini, J., Clark, K., Freedland, K.E., 1988. The relationship between heart rate, heart rate variability and depression in patients with coronary artery disease. *J Psychosom Res* 32, 159-164.
- Carroll, D., Bibbey, A., Roseboom, T.J., Phillips, A.C., Ginty, A.T., De Rooij, S.R., 2012a. Forced expiratory volume is associated with cardiovascular and cortisol reactions to acute psychological stress. *Psychophysiology* 49, 866-872.
- Carroll, D., Ginty, A.T., Der, G., Hunt, K., Benzeval, M., Phillips, A.C., 2012b. Increased blood pressure reactions to acute mental stress are associated with 16-year cardiovascular disease mortality. *Psychophysiology* 49, 1444-1448.
- Carroll, D., Phillips, A.C., Der, G., 2008. Body mass index, abdominal adiposity, obesity, and cardiovascular reactions to psychological stress in a large community sample. *Psychosom Med* 70, 653-660.
- Carroll, D., Phillips, A.C., Der, G., Hunt, K., Bibbey, A., Benzeval, M., Ginty, A.T., 2013 in press. Low forced expiratory volume is associated with blunted cardiac reactions to acute psychological stress in a community sample of middle-aged men and women. *Int J Psychophysiol*.
- Carroll, D., Phillips, A.C., Hunt, K., Der, G., 2007. Symptoms of depression and cardiovascular reactions to acute psychological stress: evidence from a population study. *Biological Psychology* 75, 68-74.
- Carroll, D., Phillips, A.C., Lovallo, W.R., 2009. Are large physiological reactions to acute psychological stress always bad for health? *Social and Personality Compass (Health Section)* 3, 725-743.
- Carroll, D., Phillips, A.C., Lovallo, W.R., 2011. The behavioural and health corollaries of blunted physiological reactions to acute psychological stress: Revising the reactivity hypothesis., in: Wright, R.A., Gendolla, G.H.E. (Eds.), *How motivation affects cardiovascular response: Mechanisms and applications*. APA Press, Washington DC.
- Carroll, D., Phillips, A.C., Ring, C., Der, G., Hunt, K., 2005. Life events and hemodynamic stress reactivity in the middle-aged and elderly. *Psychophysiology* 42, 269-276.
- Carroll, D., Ring, C., Hunt, K., Ford, G., Macintyre, S., 2003. Blood pressure reactions to stress and the prediction of future blood pressure: effects of sex, age, and socioeconomic position. *Psychosom Med* 65, 1058-1064.
- Carroll, D., Smith, G.D., Sheffield, D., Shipley, M.J., Marmot, M.G., 1995. Pressor reactions to psychological stress and prediction of future blood pressure: data from the Whitehall II Study. *BMJ* 310, 771-776.
- Carroll, D., Smith, G.D., Shipley, M.J., Steptoe, A., Brunner, E.J., Marmot, M.G., 2001. Blood pressure reactions to acute psychological stress and future blood pressure status: a 10-year follow-up of men in the Whitehall II study. *Psychosom Med* 63, 737-743.
- Carroll, D., Turner, J.R., Prasad, R., 1986. The effects of level of difficulty of mental arithmetic challenge on heart rate and oxygen consumption. *Int J Psychophysiol* 4, 167-173.
- Chida, Y., Steptoe, A., 2010. Greater cardiovascular responses to laboratory mental stress are associated with poor subsequent cardiovascular risk status: a meta-analysis of prospective evidence. *Hypertension* 55, 1026-1032.

- Cohen, G., Vella, S., Jeffery, H., Lagercrantz, H., Katz-Salamon, M., 2008. Cardiovascular stress hyperreactivity in babies of smokers and in babies born preterm. *Circulation* 118, 1848-1853.
- Connor-Smith, J.K., Flachsbart, C., 2007. Relations between personality and coping: a meta-analysis. *J Pers Soc Psychol* 93, 1080-1107.
- D'Andrea, W., under review. Heterogeneity of autonomic responses after exposure to trauma: blunted reactivity in response to startling sounds. *International Journal of Psychophysiology*.
- de Rooij, S.R., 2013 in press. Blunted cardiovascular and cortisol reactivity to acute psychological stress: A summary of results from the Dutch Famine Birth Cohort Study. *Int J Psychophysiol*.
- De Rooij, S.R., Roseboom, T.J., 2010. Further evidence for an association between self-reported health and cardiovascular as well as cortisol reactions to acute psychological stress. *Psychophysiology* 47, 1172-1175.
- de Rooij, S.R., Schene, A.H., Phillips, D.I., Roseboom, T.J., 2010. Depression and anxiety: Associations with biological and perceived stress reactivity to a psychological stress protocol in a middle-aged population. *Psychoneuroendocrinology* 35, 866-877.
- Dweck, C.S., 1986. Motivational processes affecting learning. *American Psychologist* 41, 1040-1048.
- Ellenbogen, M.A., Hodgins, S., 2009. Structure provided by parents in middle childhood predicts cortisol reactivity in adolescence among the offspring of parents with bipolar disorder and controls. *Psychoneuroendocrinology* 34, 773-785.
- Ellis, B.J., Essex, M.J., Boyce, W.T., 2005. Biological sensitivity to context: II. Empirical explorations of an evolutionary-developmental theory. *Dev Psychopathol* 17, 303-328.
- Epstein, J., Pan, H., Kocsis, J.H., Yang, Y., Butler, T., Chusid, J., Hochberg, H., Murrugh, J., Strohmayer, E., Stern, E., Silbersweig, D.A., 2006. Lack of ventral striatal response to positive stimuli in depressed versus normal subjects. *Am J Psychiatry* 163, 1784-1790.
- Eshel, N., Roiser, J.P., 2010. Reward and punishment processing in depression. *Biol Psychiatry* 68, 118-124.
- Everson, S.A., Kaplan, G.A., Goldberg, D.E., Salonen, J.T., 1996. Anticipatory blood pressure response to exercise predicts future high blood pressure in middle-aged men. *Hypertension* 27, 1059-1064.
- Everson, S.A., Lynch, J.W., Chesney, M.A., Kaplan, G.A., Goldberg, D.E., Shade, S.B., Cohen, R.D., Salonen, R., Salonen, J.T., 1997. Interaction of workplace demands and cardiovascular reactivity in progression of carotid atherosclerosis: population based study. *Bmj* 314, 553-558.
- Gendolla, G.H., Krusken, J., 2001. The joint impact of mood state and task difficulty on cardiovascular and electrodermal reactivity in active coping. *Psychophysiology* 38, 548-556.
- Georgiades, A., 2007. Hyperreactivity (cardiovascular). , in: Fink, G. (Ed.), *Encyclopedia of stress*, 2nd ed. Academic Press, Burlington, MA, pp. 372-376.
- Georgiades, A., Lemne, C., de Faire, U., Lindvall, K., Fredrikson, M., 1997. Stress-induced blood pressure measurements predict left ventricular mass over three years among borderline hypertensive men. *Eur J Clin Invest* 27, 733-739.
- Gerin, W., Pickering, T.G., Glynn, L., Christenfeld, N., Schwartz, A., Carroll, D., Davidson, K., 2000. An historical context for behavioral models of hypertension. *J Psychosom Res* 48, 369-377.
- Ginty, A.T., 2013 under review. Blunted responses to stress and reward: A model of biological disengagement. *International Journal of Psychophysiology*.

- Ginty, A.T., Gianaros, P.J., Derbyshire, S.W.G., Phillips, A.C., Carroll, D., 2013. Blunted cardiac stress reactivity relates to neural hypoactivation. *Psychophysiology* 50, 219-229.
- Ginty, A.T., Phillips, A.C., Der, G., Deary, I.J., Carroll, D., 2011a. Cognitive ability and simple reaction time predict cardiac reactivity in the West of Scotland Twenty-07 Study. *Psychophysiology* 48, 1022-1027.
- Ginty, A.T., Phillips, A.C., Der, G., Deary, I.J., Carroll, D., 2011b. Heart rate reactivity is associated with future cognitive ability and cognitive change in a large community sample. *Int J Psychophysiol* 82, 167-174.
- Ginty, A.T., Phillips, A.C., Higgs, S., Heaney, J.L., Carroll, D., 2012a. Disordered eating behaviour is associated with blunted cortisol and cardiovascular reactions to acute psychological stress. *Psychoneuroendocrinology* 37, 715-724.
- Ginty, A.T., Phillips, A.C., Roseboom, T.J., Carroll, D., Derooij, S.R., 2012b. Cardiovascular and cortisol reactions to acute psychological stress and cognitive ability in the Dutch Famine Birth Cohort Study. *Psychophysiology* 49, 391-400.
- Girdler, S.S., Jamner, L.D., Jarvik, M., Soles, J.R., Shapiro, D., 1997. Smoking status and nicotine administration differentially modify hemodynamic stress reactivity in men and women. *Psychosom Med* 59, 294-306.
- Grassi, G., Seravalle, G., Cattaneo, B.M., Bolla, G.B., Lanfranchi, A., Colombo, M., Giannattasio, C., Brunani, A., Cavagnini, F., Mancina, G., 1995. Sympathetic activation in obese normotensive subjects. *Hypertension* 25, 560-563.
- Hammar, A., Strand, M., Ardal, G., Schmid, M., Lund, A., Elliott, R., 2011. Testing the cognitive effort hypothesis of cognitive impairment in major depression. *Nord J Psychiatry* 65, 74-80.
- Heaney, J.L., Ginty, A.T., Carroll, D., Phillips, A.C., 2011a. Preliminary evidence that exercise dependence is associated with blunted cardiac and cortisol reactions to acute psychological stress. *Int J Psychophysiol* 79, 323-329.
- Heaney, J.L.J., Ginty, A.T., Carroll, D., Phillips, A.C., 2011b. Preliminary evidence that exercise dependence is associated with blunted cardiac and cortisol reactions to acute psychological stress. *International Journal of Psychophysiology* 79, 323-329.
- Higgins, N.M., Hughes, B.M., 2012. Individual differences in the impact of attentional bias training on cardiovascular responses to stress in women. *Anxiety Stress Coping* 25, 381-395.
- Howard, S., Hughes, B.M., James, J.E., 2011. Type D personality and hemodynamic reactivity to laboratory stress in women. *Int J Psychophysiol* 80, 96-102.
- Hughes, B.M., Howard, S., James, J.E., Higgins, N.M., 2011. Individual differences in adaptation of cardiovascular responses to stress. *Biol Psychol* 86, 129-136.
- Hughes, J.W., Watkins, L., Blumenthal, J.A., Kuhn, C., Sherwood, A., 2004. Depression and anxiety symptoms are related to increased 24-hour urinary norepinephrine excretion among healthy middle-aged women. *J Psychosom Res* 57, 353-358.
- James, J.E., Gregg, M.E., Matyas, T.A., Hughes, B.M., Howard, S., Stress reactivity and the Hemodynamic Profile-Compensation Deficit (HP-CD) Model of blood pressure regulation. *Biol Psychol* 90, 161-170.
- Kapuku, G.K., Treiber, F.A., Davis, H.C., Harshfield, G.A., Cook, B.B., Mensah, G.A., 1999. Hemodynamic function at rest, during acute stress, and in the field: predictors of cardiac structure and function 2 years later in youth. *Hypertension* 34, 1026-1031.
- Koo-Loeb, J.H., Pedersen, C., Girdler, S.S., 1998. Blunted cardiovascular and catecholamine stress reactivity in women with bulimia nervosa. *Psychiatry Res* 80, 13-27.

- Kotani, K., Saiga, K., Sakane, N., Mu, H., Kurozawa, Y., 2008. Sleep status and blood pressure in a healthy normotensive female population. *Int J Cardiol* 125, 425-427.
- Kupper, N., 2013 in press. Type D personality is associated with low cardiovascular reactivity to acute stress in heart failure patients *International Journal of Psychophysiology*.
- Lovallo, W.R., 2005. Cardiovascular reactivity: Mechanisms and pathways to cardiovascular disease. *International Journal of Psychophysiology* 58, 119-132.
- Lovallo, W.R., 2013 in press. Early life adversity reduces stress reactivity and enhances impulsive behavior: Implications for health behaviors. *Int J Psychophysiol*.
- Lovallo, W.R., Dickensheets, S.L., Myers, D.A., Thomas, T.L., Nixon, S.J., 2000. Blunted stress cortisol response in abstinent alcoholic and polysubstance-abusing men. *Alcohol Clin Exp Res* 24, 651-658.
- Lovallo, W.R., Farag, N.H., Sorocco, K.H., Cohoon, A.J., Vincent, A.S., 2012. Lifetime adversity leads to blunted stress axis reactivity: studies from the Oklahoma Family Health Patterns Project. *Biol Psychiatry* 71, 344-349.
- Lovallo, W.R., Gerin, W., 2003. Psychophysiological reactivity: mechanisms and pathways to cardiovascular disease. *Psychosom Med* 65, 36-45.
- Luecken, L.J., Roubinov, D.S., 2012. Hostile behavior links negative childhood family relationships to heart rate reactivity and recovery in young adulthood. *Int J Psychophysiol* 84, 172-179.
- Lynch, J.W., Everson, S.A., Kaplan, G.A., Salonen, R., Salonen, J.T., 1998. Does low socioeconomic status potentiate the effects of heightened cardiovascular responses to stress on the progression of carotid atherosclerosis? *Am J Public Health* 88, 389-394.
- Markovitz, J.H., Raczynski, J.M., Wallace, D., Chettur, V., Chesney, M.A., 1998. Cardiovascular reactivity to video game predicts subsequent blood pressure increases in young men: The CARDIA study. *Psychosom Med* 60, 186-191.
- Matthews, K.A., Owens, J.F., Kuller, L.H., Sutton-Tyrrell, K., Lassila, H.C., Wolfson, S.K., 1998. Stress-induced pulse pressure change predicts women's carotid atherosclerosis. *Stroke* 29, 1525-1530.
- Matthews, K.A., Woodall, K.L., Allen, M.T., 1993. Cardiovascular reactivity to stress predicts future blood pressure status. *Hypertension* 22, 479-485.
- Mazzola-Pomietto, P., Azorin, J.M., Tramoni, V., Jeanningros, R., 1994. Relation between lymphocyte beta-adrenergic responsivity and the severity of depressive disorders. *Biol Psychiatry* 35, 920-925.
- McClelland, D.C., Atkinson, J.W., Clark, R.A., Lowell, F.L., 1953. *The achievement motive*. Appleton-Century-Crofts, New York.
- Murdison, K.A., Treiber, F.A., Mensah, G., Davis, H., Thompson, W., Strong, W.B., 1998. Prediction of left ventricular mass in youth with family histories of essential hypertension. *Am J Med Sci* 315, 118-123.
- Nazzaro, P., Ciancio, L., Vulpis, V., Triggiani, R., Schirosi, G., Pirrelli, A., 2002. Stress-induced hemodynamic responses are associated with insulin resistance in mild hypertensives. *Am J Hypertens* 15, 865-871.
- Nesse, R.M., Ellsworth, P.C., 2009. Evolution, emotions, and emotional disorders. *Am Psychol* 64, 129-139.
- Newman, J.D., McGarvey, S.T., Steele, M.S., 1999. Longitudinal association of cardiovascular reactivity and blood pressure in Samoan adolescents. *Psychosom Med* 61, 243-249.

- O'Leary, E.D., Howard, S., Hughes, B.M., James, J.E., 2013 in press. An experimental test of blunting using sleep-restriction as an acute stressor. *International Journal of Psychophysiology*.
- Obrist, P., 1981. *Cardiovascular Psychophysiology: A perspective*. Plenum Press, New York.
- Ouellet-Morin, I., Odgers, C.L., Danese, A., Bowes, L., Shakoor, S., Papadopoulos, A.S., Caspi, A., Moffitt, T.E., Arseneault, L., 2011. Blunted cortisol responses to stress signal social and behavioral problems among maltreated/bullied 12-year-old children. *Biol Psychiatry* 70, 1016-1023.
- Palatini, P., Julius, S., 2009. The role of cardiac autonomic function in hypertension and cardiovascular disease. *Curr Hypertens Rep* 11, 199-205.
- Panknin, T.L., Dickensheets, S.L., Nixon, S.J., Lovallo, W.R., 2002. Attenuated heart rate responses to public speaking in individuals with alcohol dependence. *Alcohol Clin Exp Res* 26, 841-847.
- Phillips, A.C., 2011. Blunted cardiovascular reactivity relates to depression, obesity, and self-reported health. *Biol Psychol* 86, 106-113.
- Phillips, A.C., Carroll, D., Burns, V.E., Drayson, M., 2005a. Neuroticism, cortisol reactivity, and antibody response to vaccination. *Psychophysiology* 42, 232-238.
- Phillips, A.C., Carroll, D., Ring, C., Sweeting, H., West, P., 2005b. Life Events and Acute Cardiovascular Reactions to Mental Stress: A Cohort Study. *Psychosomatic Medicine* 67, 384-392.
- Phillips, A.C., Der, G., Carroll, D., 2009a. Self-reported health and cardiovascular reactions to psychological stress in a large community sample: cross-sectional and prospective associations. *Psychophysiology* 46, 1020-1027.
- Phillips, A.C., Der, G., Hunt, K., Carroll, D., 2009b. Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology* 73, 273-278.
- Phillips, A.C., Hunt, K., Der, G., Carroll, D., 2011. Blunted cardiac reactions to acute psychological stress predict symptoms of depression five years later: evidence from a large community study. *Psychophysiology* 48, 142-148.
- Pintrich, P.H., Schunk, D.H., 1986. *Motivation in education theory, research, and applications*. Prentice Hall, Englewood Cliffs, NJ.
- Raitakari, O.T., Pitkanen, O.P., Lehtimaki, T., Lahdenpera, S., Iida, H., Yla-Herttuala, S., Luoma, J., Mattila, K., Nikkari, T., Taskinen, M.R., Viikari, J.S., Knuuti, J., 1997. In vivo low density lipoprotein oxidation relates to coronary reactivity in young men. *J Am Coll Cardiol* 30, 97-102.
- Richter, M., Friedrich, A., Gendolla, G.H., 2008. Task difficulty effects on cardiac activity. *Psychophysiology* 45, 869-875.
- Richter, M., Gendolla, G.H., 2006. Incentive effects on cardiovascular reactivity in active coping with unclear task difficulty. *Int J Psychophysiol* 61, 216-225.
- Rottenberg, J., Clift, A., Bolden, S., Salomon, K., 2007. RSA fluctuation in major depressive disorder. *Psychophysiology* 44, 450-458.
- Roy, M.P., Steptoe, A., Kirschbaum, C., 1994. Association between smoking status and cardiovascular and cortisol stress responsivity in healthy young men. *Int J Behav Med* 1, 264-283.
- Rudorfer, M.V., Ross, R.J., Linnoila, M., Sherer, M.A., Potter, W.Z., 1985. Exaggerated orthostatic responsivity of plasma norepinephrine in depression. *Arch Gen Psychiatry* 42, 1186-1192.

- Salomon, E.A., Mimiaga, M.J., Husnik, M.J., Welles, S.L., Manseau, M.W., Montenegro, A.B., Safren, S.A., Koblin, B.A., Chesney, M.A., Mayer, K.H., 2009a. Depressive symptoms, utilization of mental health care, substance use and sexual risk among young men who have sex with men in EXPLORE: implications for age-specific interventions. *AIDS Behav* 13, 811-821.
- Salomon, K., Bylsma, L.M., White, K.E., Panaite, V., Rottenberg, J., 2013 in press. Is Blunted Cardiovascular Reactivity in Depression Mood-State Dependent? A Comparison of Major Depressive Disorder, Remitted Depression and Healthy Controls. *International Journal of Psychophysiology*.
- Salomon, K., Clift, A., Karlsdottir, M., Rottenberg, J., 2009b. Major depressive disorder is associated with attenuated cardiovascular reactivity and impaired recovery among those free of cardiovascular disease. *Health Psychol* 28, 157-165.
- Sanders, B.J., Anticevic, A., 2007. Maternal separation enhances neuronal activation and cardiovascular responses to acute stress in borderline hypertensive rats. *Behav Brain Res* 183, 25-30.
- Schallmayer, S., Hughes, B.M., Impact of oral contraception and neuroticism on cardiovascular stress reactivity across the menstrual cycle. *Psychol Health Med* 15, 105-115.
- Schwartz, A.R., Gerin, W., Davidson, K., Pickering, T.G., Brosschot, J.F., Thayer, J.F., Christenfeld, N., Linden, W., 2003. Toward a causal model of cardiovascular responses to stress and the development of cardiovascular disease. *Psychosomatic Medicine* 65, 22-35.
- Schwerdtfeger, A., Rosenkaimer, A.K., 2011. Depressive symptoms and attenuated physiological reactivity to laboratory stressors. *Biol Psychol* 87, 430-438.
- Schwerdtfeger, A.R., Gerteis, A.K., 2013 in press. Is the blunted blood pressure reactivity in dysphoric individuals related to attenuated behavioral approach? *Int J Psychophysiol*.
- Sheffield, D., Smith, G.D., Carroll, D., Shipley, M.J., Marmot, M.G., 1997. The effects of recent food, alcohol, and tobacco intake and the temporal scheduling of testing on cardiovascular activity at rest and during psychological stress. *Psychophysiology* 34, 204-212.
- Sorocco, K.H., Lovallo, W.R., Vincent, A.S., Collins, F.L., 2006. Blunted hypothalamic-pituitary-adrenocortical axis responsivity to stress in persons with a family history of alcoholism. *Int J Psychophysiol* 59, 210-217.
- Speisman, J.C., Lazarus, R.S., Mordkoff, A., Davison, L., 1964. Experimental Reduction of Stress Based on Ego-Defense Theory. *J Abnorm Psychol* 68, 367-380.
- Stice, E., Spoor, S., Bohon, C., Veldhuizen, M.G., Small, D.M., 2008. Relation of reward from food intake and anticipated food intake to obesity: A functional magnetic resonance imaging study. *J Abnorm Psychol* 117, 924-935.
- Taylor, T.R., Kamarck, T.W., Dianzumba, S., 2003. Cardiovascular reactivity and left ventricular mass: an integrative review. *Ann Behav Med* 26, 182-193.
- Tentolouris, N., Liatis, S., Katsilambros, N., 2006. Sympathetic system activity in obesity and metabolic syndrome. *Ann N Y Acad Sci* 1083, 129-152.
- Tentolouris, N., Tsigos, C., Perea, D., Koukou, E., Kyriaki, D., Kitsou, E., Daskas, S., Daifotis, Z., Makrilakis, K., Raptis, S.A., Katsilambros, N., 2003. Differential effects of high-fat and high-carbohydrate isoenergetic meals on cardiac autonomic nervous system activity in lean and obese women. *Metabolism* 52, 1426-1432.
- Treiber, F.A., Kamarck, T., Schneiderman, N., Sheffield, D., Kapuku, G., Taylor, T., 2003. Cardiovascular reactivity and development of preclinical and clinical disease states. *Psychosom Med* 65, 46-62.

- Treiber, F.A., Turner, J.R., Davis, H., Strong, W.B., 1997. Prediction of resting cardiovascular functioning in youth with family histories of essential hypertension: a 5-year follow-up. *International Journal of Behavioral Medicine* 4, 278-291.
- Vgontzas, A.N., Liao, D., Bixler, E.O., 2009. Insomnia and hypertension. *Sleep* 32, 1547.
- Welle, S., Lilavivat, U., Campbell, R.G., 1981. Thermic effect of feeding in man: increased plasma norepinephrine levels following glucose but not protein or fat consumption. *Metabolism* 30, 953-958.
- Willemsen, G., Ring, C., McKeever, S., Carroll, D., 2000. Secretory immunoglobulin A and cardiovascular activity during mental arithmetic: effects of task difficulty and task order. *Biol Psychol* 52, 127-141.
- Wright, R.A., 1996. Brehm's theory of motivation as a model of effort and cardiovascular response, in: Gollwitzer, P.M., Bargh, J.A. (Eds.), *The psychology of action: Linking cognition and motivation to behavior*. Plenum, New York.
- York, K.M., Hassan, M., Li, Q., Li, H., Fillingim, R.B., Sheps, D.S., 2007. Coronary artery disease and depression: patients with more depressive symptoms have lower cardiovascular reactivity during laboratory-induced mental stress. *Psychosom Med* 69, 521-528.
- Yu, B.H., Kang, E.H., Ziegler, M.G., Mills, P.J., Dimsdale, J.E., 2008. Mood states, sympathetic activity, and in vivo beta-adrenergic receptor function in a normal population. *Depress Anxiety* 25, 559-564.