

## The perils of Deep Brain Stimulation in Psychiatry: Big claims, on the backs of small sample sizes.

Mazaheri, Ali; Bahramisharif, Ali

DOI:

[10.1016/j.biopsych.2015.12.014](https://doi.org/10.1016/j.biopsych.2015.12.014)

License:

Creative Commons: Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

*Document Version*

Early version, also known as pre-print

*Citation for published version (Harvard):*

Mazaheri, A & Bahramisharif, A 2016, 'The perils of Deep Brain Stimulation in Psychiatry: Big claims, on the backs of small sample sizes.', *Biological Psychiatry*, vol. 80, no. 7, pp. e61-e62.  
<https://doi.org/10.1016/j.biopsych.2015.12.014>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Psychiatry

Elsevier Editorial System(tm) for Biological

Manuscript Draft

Manuscript Number: BPS-D-15-02076R1

Title: The perils of Deep Brain Stimulation in Psychiatry: Big claims, on the backs of small sample sizes. Reply to Widge et al. - Inconsistent CFC Changes from VC/VS DBS Ventral Capsule/Ventral Striatum Deep Brain Stimulation Does Not Consistently Diminish Occipital Cross-Frequency Coupling

Article Type: Correspondence

Corresponding Author: Dr. Ali Mazaheri, PhD

Corresponding Author's Institution: University of Birmingham

First Author: Ali Mazaheri, PhD

Order of Authors: Ali Mazaheri, PhD; Ali Bahramisharif

Response to Reviewers: We've done the minor changes.

# The perils of Deep Brain Stimulation in Psychiatry: Big claims, on the backs of small sample sizes

*Reply to Widge et al. – Inconsistent CFC Changes from VC/VS DBS*

*Ventral Capsule/Ventral Striatum Deep Brain Stimulation Does Not Consistently Diminish Occipital Cross-Frequency Coupling*

Ali Mazaheri<sup>1</sup> and Ali Bahramisharif<sup>2</sup>

<sup>1</sup>School of Psychology, University of Birmingham, United Kingdom.

<sup>2</sup>Academic Medical Centre, University of Amsterdam, The Netherlands.

Corresponding Author:

Ali Mazaheri

[ali.mazah@gmail.com](mailto:ali.mazah@gmail.com)

School of Psychology,

3.03 Hills Building

University of Birmingham

+44(0)121 414 2863

United Kingdom

While there is overall consensus on how DBS of the basal ganglia of Parkinson's patients leads to symptom reduction benefits, precise therapeutic mechanism of such stimulation in psychiatric disorders is still a matter of investigation. We recently proposed that the high frequency stimulation of the internal capsule of patients with obsessive compulsive disorder (OCD), could disrupt the degree of rhythmicity (i.e. Phase preservation) of ongoing oscillations generated at the cortex which would subsequently reduce the connectivity of the cortex with sub-cortical areas [1].

Supporting this view was our observation that the DBS of the ventral internal capsule attenuated cross-frequency interactions of the phase of ongoing beta (13–30 Hz), and power of low gamma (30–50 Hz), over the occipital cortex of 7 OCD patients [2]. Our study was published on the heels of another study which observed DBS-induced changes in motor cortex beta-gamma coupling during DBS for Parkinson's disease [3].

However, the current study by Widge et al. [4] did not replicate our finding of DBS modulation of the cross-frequency interaction in their sample of 4 depressed and 1 OCD patients which underwent DBS near the internal capsule. We believe Widge et al. provide an important contribution to the field of DBS in psychiatry, by clearly demonstrating that more work needs to be done to understand how DBS of subcortical areas changes activity across the cortex, and ultimately influences cognition. Currently, the field of DBS in psychiatry contains many studies which make significant claims, based on observations on very low number of subjects. These big claims based on small sample sizes make the therapeutic efficacy of DBS in alleviating psychiatric symptoms still a matter of debate. However, we do disagree with Widge et al.'s account of a factor which could NOT be causing the discrepancy in between our observations.

We are in agreement that the most significant difference between our two studies was the duration that DBS was turned off: 1 week in our study, and 2 hours in theirs. While the authors do acknowledge this difference as a possible factor, they underestimate it (we believe incorrectly) based on evidence from a previous animal study [5]. The Ewing and Grace study found that stopping DBS of the nucleus accumbens (NAcc) shell in freely moving rats resulted in a rebound of oscillatory power in the NAcc, as well as coherence between limbic sites. Widge et al's rationale here is if DBS in human patients attenuated any cross-frequency coupling, based on the Ewing and Grace study, there should be an immediate rebound in cross-frequency coupling after the halting of the stimulation. However, we need to point out here that the focus of Ewing and Grace's study was on change in the power of oscillatory activity in the NAcc, and coherence within limbic sites. These findings have little direct relevance to cross-frequency interactions at the cortical layer. Moreover, we would be quite interested to see if Widge et al. were able to replicate the oscillatory power rebounds observed by Ewing and Grace at the level of the cortex.

Additional differences between our studies were that the patients had different psychiatric symptoms, as well as frequency of stimulation. These things could indeed play a factor. However, Widge et al. propose that our opposing findings would likely be due to “un-modeled characteristics of the two samples“ such as pre-operative clinical or electrophysiologic phenotypes, slight variation in electrode placement as well as the location of the active DBS contact, and subtle differences in the testing environment. Here we fully agree with the authors that each of these factors could contribute to difference in findings between centres. One point though that we feel the need to raise is that although Widge et al. cite a recent study by Samaha et al. [6] as evidence that cognitive factors influence the phase and peak frequency property of oscillations, our work (published a few months earlier than Samaha et al.) suggests quite the contrary. We found that in healthy participants the peak frequency and phase property of oscillatory activity factors were not influenced by top-down cognitive factors [7].

Finally, the current discussion inevitably raises a rather important question. Is the attenuation on cross-frequency coupling we observed in our study subtle and dependent on finely tuned environmental and participant characteristics or something more robust and generalizable across patients? At this stage we are fully willing to concede that we currently do not have the answer to this question. We are also in full agreement with Widge et al., that in order to identify the clinical significance of DBS in psychiatry, far more coordination among centres across the world doing such research is needed. This coordination would entail the sharing of critical information such as stimulation settings, the exact location of electrodes superimposed on individual structural scans, patient characteristics, and finally analysis scripts, all of which would allow for testing for the replicability of findings. Admittedly there are likely hurdles to such open access of data, such as patient privacy issues as well as resistance from some principle investigators. However, this is a crucial period for the field of DBS in psychiatry where recent studies are questioning its efficacy [8]. In order to make big claims using small sample sizes, the books have to open.

All authors report no biomedical financial interests or potential conflicts of interest.

1. Smolders R, Mazaheri A, van Wingen G, Figeo M, de Koning PP, Denys D. Deep brain stimulation targeted at the nucleus accumbens decreases the potential for pathologic network communication. *Biol Psychiatry*. 2013;74(10):e27-8. Epub 2013/04/30. doi: 10.1016/j.biopsych.2013.03.012. PubMed PMID: 23623188.
2. Bahramisharif A, Mazaheri A, Levar N, Schuurman PR, Figeo M, Denys D. Deep brain stimulation diminishes cross frequency coupling in OCD. *Biological psychiatry*. 2015.
3. de Hemptinne C, Swann NC, Ostrem JL, Ryapolova-Webb ES, San Luciano M, Galifianakis NB, et al. Therapeutic deep brain stimulation reduces cortical phase-amplitude coupling in Parkinson's

disease. *Nat Neurosci.* 2015;18(5):779-86. doi: 10.1038/nn.3997. PubMed PMID: 25867121; PubMed Central PMCID: PMC4414895.

4. Widge A, Zorowitz S, Link K, Miller E, Deckersbach T, Eskandar E, et al. Ventral Capsule/Ventral Striatum Deep Brain Stimulation Does Not Consistently Diminish Occipital Cross-Frequency Coupling. *Biol Psychiatry.* in press

5. Ewing SG, Grace AA. Long-term high frequency deep brain stimulation of the nucleus accumbens drives time-dependent changes in functional connectivity in the rodent limbic system. *Brain Stimul.* 2013;6(3):274-85. Epub 2012/09/18. doi: 10.1016/j.brs.2012.07.007. PubMed PMID: 22981894; PubMed Central PMCID: PMC3536888.

6. Samaha J, Bauer P, Cimaroli S, Postle BR. Top-down control of the phase of alpha-band oscillations as a mechanism for temporal prediction. *Proc Natl Acad Sci U S A.* 2015;112(27):8439-44. Epub 2015/06/24. doi: 10.1073/pnas.1503686112. PubMed PMID: 26100913; PubMed Central PMCID: PMC4500260.

7. van Diepen RM, Cohen MX, Denys D, Mazaheri A. Attention and temporal expectations modulate power, not phase, of ongoing alpha oscillations. *J Cogn Neurosci.* 2015;27(8):1573-86. Epub 2015/03/17. doi: 10.1162/jocn\_a\_00803. PubMed PMID: 25774428.

8. Dougherty DD, Rezai AR, Carpenter LL, Howland RH, Bhati MT, O'Reardon JP, et al. A Randomized Sham-Controlled Trial of Deep Brain Stimulation of the Ventral Capsule/Ventral Striatum for Chronic Treatment-Resistant Depression. *Biol Psychiatry.* 2015;78(4):240-8. doi: 10.1016/j.biopsych.2014.11.023. PubMed PMID: 25726497.