

The Contribution of Youth Sport Football to Weekend Physical Activity for Males Aged 9 to 16 Years: Variability Related to Age and Playing Position

Fenton, Sally; Duda, Joan; Barrett, Timothy

DOI:
[10.1123/pes.2014-0053](https://doi.org/10.1123/pes.2014-0053)

Document Version
Peer reviewed version

Citation for published version (Harvard):
Fenton, S, Duda, J & Barrett, T 2015, 'The Contribution of Youth Sport Football to Weekend Physical Activity for Males Aged 9 to 16 Years: Variability Related to Age and Playing Position', *Pediatric Exercise Science*, vol. 27, no. 2, pp. 208-18. <https://doi.org/10.1123/pes.2014-0053>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/268230820>

The Contribution of Youth Sport Football to Weekend Physical Activity for Males Aged 9 to 16 Years: Variability Related to Age and Playing Position

ARTICLE *in* PEDIATRIC EXERCISE SCIENCE · NOVEMBER 2014

Impact Factor: 1.45 · DOI: 10.1123/pes.2014-0053 · Source: PubMed

CITATIONS

2

READS

174

3 AUTHORS, INCLUDING:



Sally A.M. Fenton

University of Birmingham

9 PUBLICATIONS 14 CITATIONS

SEE PROFILE



Joan L. Duda

University of Birmingham

332 PUBLICATIONS 9,082 CITATIONS

SEE PROFILE

1 **Doi:** <http://dx.doi.org/10.1123/pes.2014-0053>

2 **Original article available:** <http://journals.humankinetics.com/pes>

3 **The contribution of youth sport football to weekend physical activity for males aged 9-**
4 **to 16- years: Variability related to age and playing position**

5 Sally A. M. Fenton¹, Joan L. Duda¹, Timothy Barrett²

6 ¹ School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham,
7 Edgbaston, B15 2TT, UK.

8 ² School of Clinical and Experimental Medicine, University of Birmingham, Edgbaston,
9 B152TT, UK.

10 **Corresponding Author:** Sally Fenton, School of Sport, Exercise and Rehabilitation

11 Sciences, University of Birmingham, Edgbaston, UK, B15 2TT. Email:

12 s.a.m.fenton@bham.ac.uk. Tel: 0121 4158724. Fax: 0121 414 4121 (School of Sport and
13 Exercise Sciences main office)

14 **Running head:** Youth sport football and weekend physical activity

15

16

17

18

19

20

21

22

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Abstract

The aims of this study were (1) to determine minutes of moderate-to-vigorous physical activity (MVPA) and vigorous physical activity (VPA) accrued in youth sport football (also internationally referred to as soccer), and the contribution towards daily weekend MVPA and VPA for males aged 9 to 16 years, and (2) to investigate variability in these outcomes related to age and playing position. One hundred and nine male grassroots footballers (Mean age = 11.98 ± 1.75 years) wore a GT3X accelerometer for 7 days. Weekend youth sport football participation and playing position were recorded. Youth sport football MVPA ($M = 51.51 \pm 17.99$) and VPA ($M = 27.78 \pm 14.55$) contributed 60.27% and 70.68% towards daily weekend MVPA and VPA, respectively. Overall, 36.70% of participants accumulated ≥ 60 minutes MVPA and 69.70% accrued ≥ 20 minutes of VPA during youth sport. For participants aged 13 to 16 years, youth sport football MVPA and VPA were significantly higher, and contributed a greater amount towards daily weekend MVPA and VPA than for participants aged 9 to 12 years ($p < .01$). Youth sport football is an important source of MVPA and VPA at the weekend for male youth, and particularly for adolescents. Participation may offer opportunity for weekend engagement in VPA towards health enhancing levels.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Introduction

As a result of the escalating global childhood obesity epidemic, considerable attention has been drawn towards the area of physical activity (PA) promotion among youth. Evidence based guidelines state that children and adolescents should engage in ≥ 60 minutes of moderate to vigorous physical activity (MVPA) every day to accrue benefits to health (3;15;23;49). Due to recent studies indicating a dose-response relationship between PA intensity and health outcomes (14;44), the latest guidelines have incorporated a recommendation for vigorous physical activity (VPA), advising participation in VPA on at least three days a week for school-aged youth (15;49). However, concerning the volume of VPA that should be accumulated, recommendations are lacking. Indeed, little empirical evidence exists to inform guidelines for daily accumulation of VPA (21;28). The Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) study is the largest study to date that has sought to determine the duration of daily VPA engagement likely to lead to health benefits among children and adolescents. Results from this study demonstrated approximately 18 minutes of VPA discriminated between normal weight and overweight/obesity in over 2000 European youth (Females = 10 minutes, males = 18 minutes) (28). Accordingly, daily recommendations for VPA were proposed at 10 and 20 minutes for female and male children and adolescents respectively.

Numerous population-based studies indicate few children and adolescents engage in recommended levels of MVPA, and daily levels of VPA are reported to be low (9;14;45;46). As a result increased efforts have been directed towards determining settings likely to be effective towards increasing engagement in MVPA and VPA among youth. The majority of research in this domain has focused upon opportunities offered within the school setting (26;38). In particular, Physical Education (PE) , active transport and school breaks (i.e.,

1 recess or lunchtime/break time) have received considerable attention, with numerous studies
2 investigating the contribution of MVPA accumulated in these environments towards
3 guidelines for MVPA and total daily MVPA and VPA (18;29;39). Recently, research has also
4 examined the extent to which MVPA and VPA accrued within different school PA settings
5 contributes differential amounts for children of different ages and for males and females
6 (29;39). However, to date, youth PA settings outside the school environment that are likely to
7 offer opportunity for regular engagement in MVPA and VPA (i.e., both weekday and
8 weekend) have been relatively overlooked.

9 *Youth sport as an opportunity for PA engagement*

10 Government organisations across the globe have advocated youth sport as a potential
11 vehicle through which levels of PA might be increased among youth (7;10;35). However, few
12 studies have actually examined the opportunity offered by the youth sport setting for
13 engagement in PA. Further, a paucity of studies have employed objective measures of PA
14 (e.g., accelerometers) to determine levels of participation in health enhancing PA during
15 youth sport (33). The few accelerometer-based studies that have been conducted within this
16 domain have primarily examined the opportunity offered by youth sport for engagement in
17 MVPA towards recommended levels (20;25;41;51). Results have revealed that whilst youth
18 sport offers the opportunity to accrue substantial amounts of MVPA, most participants are
19 unlikely to meet daily MVPA guidelines during their involvement in youth sport (20;25;41).
20 To our knowledge, no studies have examined the opportunity provided by youth sport for
21 participation in levels of VPA that have demonstrated associations with health. Given that
22 past research indicates VPA may be more strongly linked to health outcomes than moderate
23 PA (14;44), research determining the role of youth sport for encouraging engagement in
24 health enhancing levels of VPA is warranted. Existing studies have also neglected to
25 examine the degree to which youth sport may offer a source of both MVPA and VPA at the

1 weekend. The importance of finding opportunities to increase weekend engagement in
2 MVPA, and particularly VPA has been emphasised (45). Certainly, relative to weekday PA
3 participation, weekend PA engagement is reported to be lower among youth (9;31).
4 Determining the settings in which youth are likely to accumulate health enhancing PA at the
5 weekend is therefore important from a public health standpoint. Indeed, the nature of youth
6 sport means weekend participation is likely. By contrast, research centred within the school
7 environment will result in a focus towards weekday participation in PA.

8 Previous investigations examining levels of PA engagement within the youth sport
9 context have indicated that this setting may offer differential opportunity for engagement in
10 health enhancing PA depending on a multitude of potential determinants. For example,
11 participation in MVPA and VPA during youth sport has been shown to vary as a function of
12 sex, weight status (i.e., BMI), context (i.e., structure, training sessions versus match play),
13 coach behaviour, age and sport type (8;20;25;41;51). It has been suggested that the
14 association between sport type and youth sport PA engagement may result from the differing
15 physiological demands of specific sports (51). A similar consideration may also be relevant
16 within sports, whereby the skill position (e.g., striker, midfielder) of some players may
17 require participants to engage in higher levels of MVPA and VPA relative to their teammates
18 (e.g., goal keepers). As such, where a diversity of roles is apparent within a sport,
19 participation may provide disparate opportunities for engagement in health enhancing PA (4).
20 However, existing studies have neglected to examine variability in youth sport PA
21 engagement as related to different playing positions within specific sports.

22 Age is also a particularly pertinent factor to consider when examining the opportunity
23 offered by youth sports for engagement in health enhancing PA. Evidence of an age related
24 decline in PA participation among youth emphasises the importance of identifying settings
25 that may encourage engagement in MVPA and VPA as children transition into adolescence

1 (31;42). Adolescent youth may rely more on structured opportunities for PA engagement,
2 relative to their younger counterpart who engage in more unorganised play, or ‘free play’
3 throughout the day (1;34). Youth sport may offer one such organised PA context through
4 which adolescents can accrue substantial amounts of health enhancing PA. To date, only one
5 study has examined age differences with regards to the accumulation of PA during youth
6 sport. Leek et al., (2011), reported younger participants (aged 7 to 10 years) engaged in
7 higher levels of MVPA and VPA during youth sport football and softball/baseball than their
8 older counterparts (aged 11 to 14 years). However, associations between age and variability
9 in youth sport VPA were not examined. Further, this investigation employed age-dependent
10 cut-points to estimate PA intensity (19), which have been shown to misclassify light PA as
11 moderate PA in children ≤ 10 years old (47).

12 Extending the existent literature, the primary aims of the present study were two-fold.
13 First, we determined levels of MVPA and VPA accrued during youth sport football (also
14 internationally referred to as soccer) and the contribution made towards daily weekend
15 MVPA and VPA for males aged 9 to 16 years. The number of youth football participants
16 meeting recommended guidelines for MVPA (≥ 60 minutes) and engaging in levels of VPA
17 identified as preventing excess adiposity among youth (≥ 20 minutes) (27) were also
18 determined. The second aim was to examine variability in these outcomes in relation to
19 participant age and playing position. Football was the sport focused on in the present study
20 due to high rates of participation across the globe, and potential application of findings to
21 large numbers of youth sport participants. Indeed, over 21 million children and adolescents
22 participate in youth sport football across the world (24), and research has indicated football to
23 be among the most popular youth sports internationally (2;32).

24

25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Method

Recruitment

Participants were a subsample of male youth sport football players primarily recruited within a larger multi-method European trial (the Promoting Adolescent Physical Activity (PAPA) Project; www.projectpapa.org)(16). Objective PA data (i.e., PA assessed via accelerometer) were collected from a subsample of participants recruited to the larger project in France, England and Greece (N = 417). The full protocol for the PAPA Project and the core objective PA measurement protocol are detailed elsewhere (16;50). The following sections outline the protocol followed in England where additional measures were included (i.e., PA diaries).

Recruitment protocol

The PAPA project was supported by the lead football associations in each country. To aid recruitment, the English Football Association (FA) sent out regional emails to registered football clubs. Clubs were also approached to take part in the project directly by members of the research team (via phone call and/or email, N = 58). Interested coaches were provided with information about the study protocol and passed this information on to parents and players. A total of 46 clubs (36 male, 2 female and 8 mixed) were willing to participate and were recruited to the English arm of the PAPA project.

Once recruited to the larger PAPA trial, individual teams (male, N = 111, female, N = 23) were provided with information pertaining to an additional protocol that involved measurement of PA via accelerometer for 7 days (50). The present study utilises accelerometer data collected in a sub-sample of male English PAPA participants willing to undertake these extra measures (teams, N = 38, participants, N = 149). Only male footballers were recruited as a result of low female interest in participating in the accelerometer protocol

1 (teams, N = 1). An additional 11 male teams (participants, N = 35) were recruited external to
2 the PAPA project to undertake the accelerometer protocol in order to increase the sample size
3 for the present analyses (total participants, N = 184). Informed consent and assent was
4 obtained from all parents and players before participating in the study. The present research
5 was approved by the local National Health Service Ethics Committee.

6 **The present study**

7 **Participants and protocol**

8 Participants were 184 males aged 9 to 16 years (teams, N = 49) who regularly
9 participate in youth sport football at the grassroots level. Grassroots football is defined as
10 participation at the recreational level, i.e., players are not playing for professional clubs or in
11 regional, national or international teams. Typically, players participating at the recreational
12 level engage in training sessions and/or match play within an organised setting \geq once per
13 week.

14 A trained researcher visited football training sessions to carry out the procedures.
15 During the session, participants' height and weight were measured and accelerometers and
16 PA diaries distributed. Where possible, four players representing different playing positions
17 (i.e., a striker, defender, midfielder and goalkeeper) were recruited from each team.
18 Participants were asked to wear the accelerometer for 7 days. Researchers returned one week
19 later to collect accelerometers and PA diaries.

20 **Measures**

21 *Anthropometrics*

22 Height and weight were measured with participants bare foot and wearing light
23 clothing (e.g., shorts and t-shirt). Height was measured to the nearest 0.1cm using a portable
24 stadiometer (SECA, Leicester height measure). Weight was measured to the nearest 0.1kg
25 using electric scales (Tanita, SC3310). All measures were conducted in duplicate. Average

1 values were calculated for height and weight and used to determine BMI [weight (kg)/ height
2 (m^2)].

3 *Physical activity*

4 The GT3X accelerometer (Actigraph; Pensacola, FL) was used to measure MVPA
5 and VPA. Actigraph accelerometers are the most commonly used accelerometers in PA
6 research and have been validated against criterion measures of PA youth (12;48).
7 Accelerometers were initialised to measure PA in 15 second epochs. Verbal instructions were
8 given by the researcher on how the accelerometer should be worn and a demonstration given.
9 Participants were asked to wear the accelerometer on the right hip for seven days during all
10 waking hours, removing only for participation in water sports (e.g., swimming), bathing and
11 sleeping. Accelerometer log sheets and PA diaries were also distributed to enable participants
12 to record non-wear time and to aid with data cleaning and interpretation. Participants were
13 asked to report times they had participated in youth sport football on Saturday and/or Sunday
14 in PA diaries and to record their playing positions (i.e., striker, midfielder, defender,
15 goalkeeper). Log sheets, PA diaries and accelerometers were collected one week later.

16 **Data processing**

17 Physical activity data were downloaded from the GT3X to a computer and analysed
18 using the Actilife software (version 6.2, Actigraph). Data were checked for spurious values
19 and periods of non-wear. Non-wear time was determined by identifying strings of
20 consecutive zeros in the movements counts recorded by the accelerometer for > 30 minutes,
21 allowing for 1 minute of counts <100 (6). Cut-points derived by Evenson and colleagues
22 were used to determine time spent in MVPA (≥ 2296 cpm) and VPA (≥ 4012 cpm) (17).

23 *Youth sport and daily weekend physical activity*

24 Time spent in youth sport football at the weekend was identified from PA diaries (i.e.,
25 day and time). For each participant, diary entries were compared against graphed data to

1 check for accuracy in self-reported timings. Following this, individual time filters were
2 applied to the Actilife software for the identified time period to determine minutes in MVPA
3 and VPA during youth sport football. Total daily MVPA and VPA (minutes) for the day on
4 which youth sport was participated in (i.e., a Saturday or Sunday) were calculated to
5 represent daily weekend MVPA and VPA. A weekend day was considered valid where ≥ 8
6 hours of wear time had been recorded.

7 **Protocol compliance**

8 Participants were required to have recorded one valid weekend day of accelerometer
9 data (inclusive of their sport participation) to be retained for subsequent analyses. Of the
10 original 184 participants, 15.8% (N = 29) did not record valid daily weekend PA data on the
11 day on which youth sport was participated in, and 9.2% (N = 17) failed to return completed
12 PA diaries. A further 15.8% (N = 29) did not record participation in youth sport at the
13 weekend (e.g., monitor was removed for youth sport participation, youth sport session was
14 cancelled) and/or information given in diaries was deemed to be inaccurate based on visual
15 comparisons with graphed data. The final sample therefore included 109 males aged 9 to 16
16 years (compliance = 58.6%). This is comparable to compliance levels reported in the Health
17 Survey for England 2011 for boys aged 8 to 15 years (i.e., 45-49%) (22).

18 **Statistical analysis**

19 *The contribution of youth sport to daily weekend MVPA and VPA*

20 Descriptive statistics [(M \pm SD and ranges (minutes))], for youth sport MVPA and
21 VPA were calculated. Percentages of youth sport session time engaged in MVPA and VPA
22 were also computed. The number of participants meeting recommended guidelines for
23 MVPA (i.e., ≥ 60 minutes) and achieving ≥ 20 minutes of VPA were determined. The
24 guideline of ≥ 20 minutes VPA was employed based on recommendations derived from the
25 HELENA study in which data was analysed from over European 2000 youth. Youth sport

1 MVPA and VPA (minutes) were used to calculate the contribution (%) of weekend youth
 2 sport football towards daily weekend MVPA and VPA [(Weekend youth sport MVPA/VPA
 3 (mins)/ daily weekend MVPA/VPA) x100]. Data was checked for normality and non-
 4 normally distributed data was log transformed. Where transformations reduced data
 5 skewness, transformed variables were used in subsequent analysis¹

6 *Variability related to age and playing position*

7 Descriptive statistics were calculated for youth sport MVPA and VPA and daily
 8 weekend MVPA and VPA for different playing positions (strikers (N = 25), midfielders (N =
 9 41), defenders (N = 32) and goal keepers (N = 6) and age groups [aged 9 to 12 years, N = 69
 10 (M = 10.72, SD = .95), and 13 to 16 years, N = 40 (M = 13.98, SD = .83)]. Age groups were
 11 devised based on data demonstrating the age related decline in PA is steepest between the
 12 ages of 13 to 18 years (42). Variability in youth sport MVPA and VPA (minutes and
 13 percentage of session time) and daily weekend MVPA and VPA related to participant age and
 14 playing position were examined via ANCOVAs. Youth sport context (training versus match)
 15 was included as a covariate in all ANCOVAs². Where total minutes in MVPA and VPA were
 16 dependent variables, youth sport session length was also included as a covariate. Chi square
 17 tests were conducted to determine if participants were more likely to meet guidelines for
 18 MVPA and VPA during youth sport football based on age group and playing position.
 19 ANCOVAs controlling for youth sport context and youth sport session length were also

¹ Transformations did not reduce data skewness for 2 variables; the contribution of youth sport to daily weekend MVPA (1), and VPA (2). Non-transformed variables were therefore used in subsequent analysis (Figure 1). Non-parametric statistical tests (i.e., Mann-Whitney tests) were conducted and confirmed results from parametric analysis (i.e., ANCOVAs). Results from non-parametric tests are therefore reported in order to demonstrate effect sizes.

² Due to the nature of youth sport, participants reported engaging in a mixture of training sessions and match play across the weekend (training, N = 49 match play = 60). Age was not significantly different across contexts ($t(107) = -.75, p = .46$, training M = 11.77 ± 1.73 , match play = 12.03 ± 1.89), and there was no significant association between age and context ($\chi^2(1) = 3.14, p = .07$, Cramers V = $<.17$).

1 carried out to determine if the contribution of weekend youth sport to daily weekend MVPA
2 and VPA differed significantly between age groups and playing positions.

3 **Results**

4 **Participant characteristics**

5 Physical characteristics by sample (total, included and excluded) are displayed in
6 Table 1. Independent samples t-tests demonstrated excluded participants did not differ
7 significantly from those included with respect to age, height, weight or BMI-SDS (all $p =$
8 $>.05$). Questionnaire responses for participant ethnicity were incomplete (non-respondents;
9 total sample = 20.65%, [included = 11.41%, excluded = 9.24%]). Among all respondents,
10 72.8 % were white, 14.3% were Asian, 9.5% were black and 3.4% were multi-racial. Chi
11 square tests demonstrated the ethnic distribution of participants included in the final sample
12 differed significantly from those excluded ($\chi^2 (3) = 17, p = <.01$, *included*: white = 83.0 %,
13 Asian = 5.7%, black = 6.8%, multi-racial = 4.5%, *excluded*: white = 56.9 %, Asian = 27.6%,
14 black = 13.8%, multi-racial = 1.7%).

15 **MVPA and VPA engagement during youth sport football**

16 Weekend youth sport football session time was between 60 and 160 minutes ($M =$
17 106.74 ± 18.90). Total minutes and percentage of session time engaged in MVPA and VPA
18 during weekend youth sport football are reported in Table 2. Ranges for youth sport MVPA
19 and VPA were 77.0 and 63.75 minutes and 71.33% and 52.54%, respectively. Results
20 demonstrated 36.70% of participants met recommended guidelines for MVPA during youth
21 sport and 69.70% of participants' accrued ≥ 20 minutes of VPA.

22 **The contribution of youth sport football to daily weekend MVPA and VPA**

23 Daily weekend MVPA and VPA were $M = 91.26 \pm 35.44$ and $M = 40.89 \pm 20.73$
24 respectively. MVPA accrued during weekend youth sport football contributed $60.27\% \pm$

1 19.48% (range = 25.31% – 91.73%) towards daily weekend MVPA. For VPA, minutes
 2 accrued contributed $70.68\% \pm 21.53\%$ (range = 24.55% – 100%) towards daily weekend
 3 VPA (Figure 1).

4 **Variability related to age and playing position**

5 Significant differences were found for time spent in weekend youth sport MVPA and
 6 VPA between age groups. Children in the older age group engaged in significantly more
 7 minutes of MVPA ($F(1, 105) = 15.76, p < .01, \eta^2 = .13$) and VPA ($F(1, 105) = 23.72, p =$
 8 $< .01, \eta^2 = .18$), and spent a higher percentage of youth sport session time in MVPA ($F(1,$
 9 $106) = 14.11, p < .01, \eta^2 = .12$) and VPA ($F(1, 106) = 21.13, p < .01, \eta^2 = .13$) than those
 10 in the younger age group (Table 2). Chi square tests indicated older participants were more
 11 likely to accrue ≥ 60 minutes MVPA ($\chi^2(1) = 25.81, p < .01, \text{Cramers } V = .49$), and ≥ 20
 12 minutes of VPA during youth sport ($\chi^2(1) = 6.99, p = .01, \text{Cramers } V = .25$).

13 Daily weekend MVPA was not significantly different between age groups (9 to 12
 14 years, $M = 90.19 \pm 33.48$ vs. 13 to 16 years, $M = 93.11 \pm 38.98, F(1, 105) = .02, p = .89, \eta^2$
 15 $= .00$). However, daily weekend VPA was significantly higher for older compared to younger
 16 participants (9 to 12 years, $M = 36.43 \pm 17.35$ vs. 13 to 16 years, $M = 48.59 \pm 23.85, F(1,$
 17 $105) = 5.20, p < .05, \eta^2 = .05$). Weekend youth sport PA contributed a significantly greater
 18 amount towards daily weekend MVPA and VPA for older compared to younger participants
 19 (Figure 1, MVPA, $F(1, 105) = 14.18, p < .01, \eta^2 = .12$, VPA, $F(1, 105) = 13.14, p < .01,$
 20 $\eta^2 = .11$).

21 No significant differences were reported between playing positions for minutes in
 22 youth sport MVPA ($F(3, 98) = .41, p = .75, \eta^2 = .01$) or VPA ($F(3, 98) = .58, p = .63, \eta^2 =$
 23 $.02$) and percent session time in MVPA ($F(3, 99) = .58, p = .65, \eta^2 = .02$) or VPA ($F(3, 99)$
 24 $= .62, p = .61, \eta^2 = .02$). Analyses were repeated to adjust for the significant association
 25 between age and youth sport MVPA and VPA and remained non-significant (all $p > .38$).

1 There was no significant association between playing position and meeting guidelines for
2 MVPA ($\chi^2(3) = .20, p = .98$) or accruing ≥ 20 minutes VPA, ($\chi^2(3) = .52, p = .92$) during
3 youth sport.

4 Daily weekend MVPA and VPA were not significantly different between playing
5 positions (MVPA, striker, $M = 95.39 \pm 36.97$, midfielder, $M = 90.32 \pm 32.06$, defender, $M =$
6 91.19 ± 42.41 , goal keeper, $M = 79.29 \pm 16.67, F(3, 98) = .46, p = .71, \eta^2 = .01$; VPA,
7 striker, $M = 40.18 \pm 19.47$, midfielder, $M = 41.16 \pm 18.63$, defender, $M = 42.16 \pm 25.99$, goal
8 keeper, $M = 32.38 \pm 9.77, F(3, 98) = .63, p = .60, \eta^2 = .02$). Results also indicated weekend
9 youth sport did not contribute significantly different amounts towards daily weekend MVPA
10 and VPA for participants in different playing positions (Figure 1, MVPA, $F(3, 98) = .53, p =$
11 $.67, \eta^2 = .02$, VPA, $F(3, 98) = .53, p = .67, \eta^2 = .02$, after adjustment for age, all $p > .52$).

12 Discussion

13 The present study examined levels of engagement in MVPA and VPA during youth
14 sport football sessions and determined the contribution made towards daily weekend MVPA
15 and VPA in males aged 9 to 16 years. Further, variability in both youth sport PA and the
16 contribution towards daily weekend PA was investigated relative to participant age and
17 playing position. Results demonstrate youth sport football is an important source of MVPA
18 and VPA at the weekend for young males. In particular, participation may offer the
19 opportunity for weekend engagement VPA towards health enhancing levels. Findings also
20 revealed differences between children (aged 9 to 12 years) and adolescents (aged 13 to 16
21 years) in terms of their levels of engagement in MVPA and VPA during youth sport, and the
22 contribution made towards daily weekend engagement in these behaviours.

23 *MVPA and VPA engagement during youth sport football*

24 Present results demonstrated that on average, male youth sport football participants
25 engaged in approximately 52 and 28 minutes of MVPA and VPA, respectively. As such,

1 findings indicate that participation in weekend youth sport football may encourage
2 engagement in MVPA and VPA towards levels identified as being beneficial for health
3 (23;28). Leek et al., (2011) reported comparable levels of MVPA and VPA participation
4 during youth sport among boys aged 7 to 14 years. Specifically, results indicated
5 softball/baseball and soccer (football) participants accumulated 56 minutes of MVPA and up
6 to 30 minutes of VPA during youth sport. However, other studies have reported lower
7 estimates of youth sport MVPA and VPA engagement among boys of similar ages (41;51).
8 For example, Wickel et al., (2007) found boys participating in basketball, flag football and
9 soccer (football) aged 6 to 12 years to accumulate 26 minutes of MVPA and up to 18 minutes
10 of VPA minutes during youth sport time. Similarly, Satchek et al., (2011) reported boys,
11 aged 7 to 10 years, accrued only 16 minutes of MVPA and 4 minutes of VPA during youth
12 sport football.

13 Differing results across studies may partly reflect sample differences in sport type.
14 For example, Leek et al., (2011) reported male footballers (soccer participants) accrued
15 approximately 56 of MVPA during youth sport, whereas male softball/baseball participants
16 participated in MVPA for approximately 48 minutes³. However, the variability in levels of
17 MVPA and VPA reported may, in part, also be due to differences between studies in terms of
18 youth sport session length. Indeed, after accounting for youth sport time, Wickel et al., (2007)
19 and Leek et al., (2011) reported boys representing a variety of different sports, spent on
20 average 49% and 54% of youth sport time engaged in MVPA, respectively. In the present
21 study, male youth sport footballers were engaged in MVPA for 50% of their youth sport
22 session. In addition, Wickel et al., (2007) demonstrated VPA was participated in for 22% of
23 youth sport time, and current results revealed youth sport footballers spent 27% of youth
24 sport sessions engaged in VPA.

³ Data taken from graph (Leek et al., 2011)

1 Whilst disparate study findings may indeed reflect sample differences related to sport
2 type and youth sport session length, the findings of Sacheck et al., (2011) also point to the
3 need to consider the data reduction procedures when drawing comparisons across
4 investigations. Their study revealed boys accrued only 17 minutes, and spent 30% of youth
5 sport time engaged in MVPA, estimates considerably lower than reported in both past studies
6 and the present research (25;51). Both Wickel et al., (2007) and Leek et al., (2011) used age-
7 dependent accelerometer cut-points to estimate engagement in MVPA and VPA. Conversely,
8 Sacheck, et al., (2011) employed single value cut-points to measure MVPA and VPA in
9 participants aged 7 to 10 years [$\geq 3180 - 8159$ cpm (MVPA), ≥ 8160 cpm (VPA)]. These cut-
10 points are relatively higher than the equivalent age-dependent cut-points for youth of the
11 same age [$\geq 705 - 1017$ cpm (MVPA), $\geq 3136 - 3696$ cpm (VPA)]. Thus, the more
12 conservative cut-points utilized by Sacheck et al., (2011) may have contributed towards the
13 lower levels of MVPA and VPA reported in their study.

14 The present study employed the Evenson et al., (2008) cut-points to estimate MVPA
15 and VPA engagement during youth sport. These single-value cut-points have been reported to
16 demonstrate superior reliability and validity over other independently validated cut-points for
17 youth and are currently recommended for use in children and adolescents (47). Present
18 findings may therefore more accurately reflect frequency, intensity and duration of PA
19 engagement among male youth sport football participants than those reported in some
20 previous studies (25;41;51). To our knowledge, only one study to date has employed the
21 Evenson et al., (2008) cut-points to examine levels of MVPA engagement during youth sport
22 football (8). Cohen et al., (2014) revealed participants accrued only 23 minutes of MVPA
23 during youth sport football and spent 37% of youth sport time engaged in MVPA. The lower
24 levels of MVPA, relative to those reported presently, may be partly due to the inclusion of

1 female participants in the previous study. Certainly, past research has demonstrated girls to
2 engage in significantly lower levels of MVPA during youth sport than boys (20;25).

3 To our knowledge, ours is the first investigation to examine the opportunity provided
4 by youth sport for engagement in levels of VPA that have demonstrated associations with
5 health. Results revealed 70% of participants accrued ≥ 20 minutes of VPA, a cut-off reported
6 to discriminate between normal weight and overweight/obesity in male European youth (28).
7 Thus, participation in youth sport football may be conducive towards engagement in levels of
8 VPA required to prevent the development of excess adiposity in young male footballers.
9 However, whilst 20 minutes of VPA has been found to confer positive health benefits, this
10 cut-off is based on the results of a single study. Hay et al., (2012) reported independent
11 associations between vigorous PA and cardiometabolic risk factors occurred across a narrow
12 range of PA duration, revealing only 7 minutes of VPA to be associated with reduced risk for
13 overweight and elevated systolic blood pressure among Canadian youth (21). In the current
14 study, 98.17% of participants (N = 107) accrued ≥ 7 minutes of VPA during youth sport (data
15 not shown)⁴. Additional research is therefore necessary to determine levels of VPA
16 participation which are associated with benefits across a range of health indicators (e.g.,
17 vascular, metabolic) to inform evidence-based guidelines for VPA. The development of such
18 guidelines will facilitate researchers in their attempts to determine youth PA settings likely to
19 offer children and adolescents the optimal opportunity for engagement in levels of VPA
20 conducive towards enhanced health.

21 Whilst present findings indicate youth sport football may indeed offer opportunities
22 for young males to accrue substantial amounts of VPA, results suggest participation in this
23 PA context alone is unlikely to promote engagement in MVPA toward levels required to lead
24 to health benefits. Results revealed only 37% of participants met recommended guidelines for

⁴ Data available from the first author upon request

1 MVPA during youth sport sessions, and youth sport MVPA was reported to be as low as 14
2 minutes (~14% of session time). Whilst the use of differing accelerometer cut-points
3 somewhat precludes comparisons across studies, current findings are in line with those of
4 Leek et al., (2011) who found only 24% of participants to meet daily guidelines for MVPA
5 during youth sport, and investigations reporting participants can spend up to 70% of youth
6 sport time engaged in PA below a moderate intensity (20;41). Taken together, past and
7 present findings suggest there is scope to increase the duration of engagement in MVPA to
8 further optimise youth sport participation, and in turn, contribute towards obesity and disease
9 prevention among youth active in this context. However, also important to consider is that
10 whilst a number of participants accrued ≥ 20 minutes of VPA, nearly a third did not achieve
11 this recommendation. Moreover, only 27% of youth sport time was spent engaged in VPA.
12 Maximising youth sport as a context for PA engagement should therefore help to contribute
13 towards more youth achieving recommended and health enhancing levels of both MVPA and
14 VPA on youth sport days.

15 *The contribution of youth sport football to daily weekend MVPA and VPA*

16 To our knowledge, only one previous study has examined the contribution of youth
17 sport towards objectively measured daily MVPA, reporting youth sport contributed 23%
18 towards MVPA on a weekday (51). The current study is the first to determine the
19 contribution of youth sport towards daily weekend levels of MVPA, revealing a contribution
20 of approximately 60%. The higher contribution observed in the present research suggests that
21 youth sport likely becomes a more important source of MVPA at the weekend, in the absence
22 of structured opportunities for PA engagement that are inherent within the school day (e.g.,
23 physical education, school breaks/recess). Extant research has also not considered the
24 contribution of youth sport to daily weekday or weekend VPA participation.

1 Previous work has emphasised the necessity to study the main activities undertaken when
2 engaged in VPA (45). The present study, therefore, contributes to the literature by reporting
3 VPA accumulated during youth sport comprises approximately 71% towards daily weekend
4 VPA. As such, results point to youth sport football as an important source of VPA at the
5 weekend on days in which youth sport is participated in. Still, studies that examine the
6 contribution of youth sport to weekday VPA are necessary to determine the value of PA
7 participation in this context as a source of regular VPA throughout the week.

8 *Variability related to age and playing position*

9 Current findings revealed that participants, aged 13 to 16 years, accrued more minutes
10 of MVPA and VPA, and spent a higher percentage time engaged in MVPA and VPA, during
11 youth sport compared to those aged 9 to 12 years. Results also indicated that relative to their
12 younger counterparts, older participants were more likely to meet guidelines for MVPA and
13 VPA during youth sport football. Moreover, PA accumulated in this setting contributed a
14 greater amount towards daily weekend MVPA and VPA for older participants. These results
15 imply that youth sport participation may offer greater opportunity for engagement in MVPA
16 and VPA, and represent a greater source of weekend PA for youth during adolescence than
17 during childhood. Leek et al., (2011) revealed contrasting findings to those reported
18 presently, demonstrating youth sport participants' aged 7-10 years engaged in 7 minutes more
19 MVPA during youth sport than those aged 11-14 years. Divergent findings may be explained
20 by the use of age-dependent cut-points in the past study and likely overestimation of MVPA
21 in participants ≤ 10 years old (47).

22 Given the dearth of studies investigating age related variability in youth sport PA,
23 discussion as to why older participants engaged in higher level of MVPA and VPA during
24 youth sport than younger participants are somewhat speculative. Perhaps, as children reach
25 adolescence, the focus of the coach is removed away from knowledge delivery and

1 instruction regarding basic skills, towards refining skills via game play and activities that
2 facilitate aerobic conditioning, in order to prepare for more competitive play. Indeed, a recent
3 study reported youth sport MVPA to be lower when coaches were providing general
4 instruction and engaged in skill demonstration, relative to during drills, fitness instruction and
5 small group activities (8). Thus, adjustment in the organisation and focus of youth sport
6 sessions as children transition through the stages of sport specialisation (11), coupled with an
7 inherent age related increase in strength and power as youth advance through puberty (40),
8 may result in increased engagement in high intensity PA as children progress into adolescent
9 teams. In turn, youth sport football may likely offer a greater source of health enhancing PA
10 for adolescents relative to children. The present study provides initial evidence for this
11 contention, demonstrating youth sport contributed 72% and 54% towards daily MVPA for
12 adolescents and children respectively. Perhaps as children advance into adolescence, they
13 rely more on organised PA settings (such as youth sport) as opportunities for engagement in
14 health enhancing PA. For example, Nilsson et al., (2009) demonstrated frequency of outdoor
15 play after school to be a significant correlate for daily MVPA in 9 year old European youth,
16 which was attenuated in favour of participation in sport and exercise in clubs in 15 year olds
17 (34). Encouraging participation in youth sport football may therefore have implications for
18 counteracting the age related decline in PA observed among youth during the teenage years
19 (29;30;45). Indeed, a more marked decline in MVPA has been reported to occur between the
20 ages of 13 and 16 in non-sport participants compared to sport participants (27).

21 The present investigation indicated playing position was not related to variability in
22 youth sport MVPA or VPA among male youth sport footballers. By contrast, research
23 conducted in adults indicates position classification to be related to time spent running,
24 sprinting, and engaged in high intensity activity (4;13;30). However, extant studies have
25 largely examined males participating in professional leagues at the elite level. Thus, current

1 results suggest that where players are participating at the earlier stages of sport specialisation
2 (e.g., grassroots participation), the physical demands of different playing positions (and
3 resulting levels of PA engagement), may be more comparable. Nevertheless, there was a
4 trend towards lower levels of MVPA and VPA among goal keepers. The lack of a significant
5 PA differences between outfield players and goal keepers in the present study may be a
6 consequence of the low number of goal keepers recruited and the high variability in youth
7 sport MVPA and VPA among players.

8 Strengths of the present study include the use of accelerometers to measure PA,
9 allowing estimation of duration and intensity of activity during youth sport football sessions.
10 In addition, the broad age range of participants recruited allows important questions to be
11 answered pertaining to the value of youth sport as a source of health enhancing PA for
12 children, relative to adolescents. Moreover, the sample represented youth sport footballers
13 from 49 different teams. As such, results presented are less likely confounded by between-
14 team differences (e.g., coach behaviour) relative to previous studies in which fewer teams
15 have been recruited (8;20;25;41). Limitations to the current research include possible sample
16 bias; participants were self-selected and largely represent white English males who
17 participate in youth sport football. Thus, results presented may not be generalisable to young
18 people representing the wider ethnic distribution of children who participate in youth sport
19 football, females and children and adolescents who participate in other youth sports. Indeed,
20 studies conducted within the youth sport domain highlight the multitude of factors which can
21 affect PA engagement during youth sport (8;20;25;41;51). Thus, future studies are needed to
22 determine the opportunity offered by specific sports for participation in health enhancing PA
23 (and the contribution towards daily PA) across the broad spectrum of youth who participate
24 in sport at grassroots level. In particular and in view of research demonstrating girls engage in

1 lower levels of PA during youth sport than boys, there is a need to examine context specific
2 PA engagement in boys and girls separately (36).

3 The field based nature of data collection procedures also meant we were unable to
4 collect data pertaining to pubertal stage of participants. Data were collected in small groups
5 and as such, asking questions pertaining to stage of biological maturation (e.g., the Tanner
6 scale) in this group setting may have reduced player's willingness to participate. However,
7 whilst we are unable to account for the role of biological maturation when examining age
8 related differences, a recent review concluded associations between biological maturation and
9 levels of PA engagement among youth were inconsistent and often reported to be low (43).
10 Moreover, this review reported chronological age independently influences the association
11 between biological maturation and PA. Indeed, studies which sample sufficiently across age
12 ranges have demonstrated pubertal status is typically highly correlated with age (5;37), which
13 may account for this mediating effect. Thus, whilst associations between biological
14 maturation and youth sport MVPA and VPA may indeed be plausible, this relationship is
15 unlikely to have a significant influence on the results reported herein.

16 In conclusion, results demonstrate that youth sport football is an important source of
17 MVPA and VPA at the weekend for male youth aged 9 to 16 years. Present findings also
18 indicate that participation in youth sport football may encourage weekend engagement in
19 VPA towards health enhancing levels. However, youth sport football may encourage higher
20 engagement in MVPA and VPA, and represent a more important source of weekend MVPA
21 and VPA, for older compared to younger participants. Observed variability in youth sport
22 MVPA and VPA, among both children and adolescents active in this context, highlights the
23 need to optimise the youth sport experience in regard to young people's participation in
24 health enhancing PA.

25

Reference List

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

- (1) Anderson SE, Economos CD, Must A. Active play and screen time in US children aged 4 to 11 years in relation to sociodemographic and weight status characteristics: a nationally representative cross-sectional analysis. BMC Public Health. 2008;8:366.
- (2) Australian Bureau of Statistics. Children's Participation in Cultural and Leisure Activities [internet]. 2009. Available from: <http://www.abs.gov.au/ausstats/abs@.nsf/Products/4901.0~Apr+2009~Main+Features~Organised+sport>
- (3) Australian Government: Department for Health and Ageing. Physical Activity Guidelines for Children (5-12 yrs) [internet]. 2004. Available from: <http://www.beactive.wa.gov.au/index.php?id=474>
- (4) Bloomfield J, Poleman R, O'Donoghue P. Physical demands of different positions in FA Premiere League soccer. J Sports Sci Med. 2007;6(1):63-70.
- (5) Booth A, Johnson DR, Granger DA, Crouter AC, McHale S. Testosterone and child and adolescent adjustment : the moderating role of parent and child relationships. Developmental Psychology. 2003;39(1):85-98.
- (6) Cain K, Sallis JF, Conway TL, Van Dyck D, Calhoun L. Using Accelerometers in Youth Physical Activity Studies: A Review of Methods. J Phys Act Health. 2013;10(3):437-50.

- 1 (7) Centers for Disease Control and Prevention. Promoting Better Health for Young
2 People through Physical Activity and Sports [report]. 2000. Available from:
3 http://usa.usembassy.de/etexts/sport/Promoting_better_health
- 4 (8) Cohen A, McDonald S, McIver K, Pate R, Trost S. Assessing Physical Activity
5 During Youth Sport: The Observational System for Recording Activity in Children:
6 Youth Sports. *Pediatr Exerc Sci*. 2014;26(2):203-9.
- 7 (9) Collings PJ, Wijndaele K, Corder K, Westgate K, Ridgway CL, Dunn V, et al.
8 Levels and patterns of objectively-measured physical activity volume and intensity
9 distribution in UK adolescents: the ROOTS study. *Int J Behav Nutr Phys Act*.
10 2014;11(23).
- 11 (10) Commission of the European Communities. White Paper on Sport [report]. 2007.
12 Available from: [http://europa.eu/documentation/official-docs/white-](http://europa.eu/documentation/official-docs/white-papers/index_en.htm)
13 [papers/index_en.htm](http://europa.eu/documentation/official-docs/white-papers/index_en.htm)
- 14 (11) Côté J. The influence of the family in the development of talent in sports. *Sports*
15 *Psychologist*. 1999;13:395-417.
- 16 (12) de Vries SI, Van Hirtum H, Bakker I, Hopman-Rock M, Hirasing RA, Van
17 Mechelen W. Validity and Reproducibility of Motion Sensors in Youth: A
18 Systematic Update. *Med Sci Sports Exerc*. 2009;41(4):818-27.
- 19 (13) Dellal A, Chamari K, Wong PL, Ahmaidi S, Keller D, Barros R, et al. Comparison
20 of physical and technical performance in European soccer match-play: FA Premier
21 League and La Liga. *Eur J Sport Sci*. 2011;11(1):51-9.

- 1 (14) Dencker M, Andersen LB. Health-related aspects of objectively measured daily
2 physical activity in children. *Clin Physiol Funct Imaging*. 2008;28(3):133-44.
- 3 (15) Department of Health. UK physical activity guidelines [internet]. 2011. Available
4 from: <http://www.dh.gov.uk>
- 5 (16) Duda JL, Quested E, Haug E, Samdal O, Wold B, Balaguer I, et al. Promoting
6 Adolescent health through an intervention aimed at improving the quality of their
7 participation in Physical Activity (PAPA): Background to the project and main trial
8 protocol. *International Journal of Sport and Exercise Psychology*. 2013;11(4):319-
9 27.
- 10 (17) Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two
11 objective measures of physical activity for children. *J Sports Sci*. 2006;24(14):1557-
12 65.
- 13 (18) Fairclough S, Stratton G. 'Physical education makes you fit and healthy'. *Physical*
14 *education's contribution to young people's physical activity levels*. *Health Educ Res*.
15 2005;20(1):14-23.
- 16 (19) Freedson P, Pober D, Janz KF. Calibration of accelerometer output for children.
17 *Med Sci Sports Exerc*. 2005;37(11 Suppl):S523-S530.
- 18 (20) Guagliano JM, Rosenkranz RR, Kolt GS. Girls' Physical Activity Levels during
19 Organized Sports in Australia. *Med Sci Sports Exerc*. 2013;45(1):116-22.

- 1 (21) Hay J, Maximova K, Durksen A, Carson V, Rinaldi RL, Torrance B, et al. Physical
2 Activity Intensity and Cardiometabolic Risk in Youth. Arch Pediatr Adolesc Med.
3 2012;166(11):1022-9.
- 4 (22) Health and Social Care Information Centre. Health Survey for England, 2011:
5 Health, social care and lifestyles. Summary of key findings [report]. 2012. Available
6 from: www.ic.nhs.uk/pubs/hse11report
- 7 (23) Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity
8 and fitness in school-aged children and youth. Int J Behav Nutr Phys Act.
9 2010;7:40.
- 10 (24) Kunz M. 265 million playing football. FIFA Magazine, July issue [internet]. 2007.
11 Available from:
12 http://www.fifa.com/mm/document/fifafacts/bcoffsurv/emaga_9384_10704.pdf
13
14
- 15 (25) Leek D, Carlson JA, Cain KL, Henrichon S, Rosenberg D, Patrick K, et al. Physical
16 activity during youth sports practices. Arch Pediatr Adolesc Med. 2011;165(4):294-
17 9.
- 18 (26) Lonsdale C, Rosenkranz RR, Peralta LR, Bennie A, Fahey P, Lubans DR. A
19 systematic review and meta-analysis of interventions designed to increase moderate-
20 to-vigorous physical activity in school physical education lessons. Prev Med.
21 2013;56(2):152-61.

- 1 (27) Machado-Rodrigues AmM, Coelho e Silva MJ, Mota J, Santos RM, Cummin SP,
2 Malina RM. Physical Activity and Energy Expenditure in Adolescent Male Sport
3 Participants and Nonparticipants Aged 13 to 16 Years. *J Phys Act Health*.
4 2012;9(5):626-33.
- 5 (28) Martinez-Gomez D, Ruiz JR, Ortega FB, Veiga OL, Moliner-Urdiales D, Mauro B,
6 et al. Recommended Levels of Physical Activity to Avoid an Excess of Body Fat in
7 European Adolescents: The HELENA Study. *Am J Prev Med*. 2010;39(3):203-11.
- 8 (29) Meyer U, Roth R, Zahner L, Gerber M, Puder JJ, Hebestreit H, et al. Contribution
9 of physical education to overall physical activity. *Scand J Med Sci Sports*.
10 2013;23(5):600-6.
- 11 (30) Mohr M, Krstrup P, Bangsbo J. Match performance of high-standard soccer
12 players with special reference to development of fatigue. *J Sports Sci*.
13 2003;1;21(7):519-28.
- 14 (31) Nader PR, Bradley RH, Houts RM, McRitchie SL, O'Brien M. Moderate-to-
15 vigorous physical activity from ages 9 to 15 years. *JAMA*. 2008;300(3):295-305.
- 16 (32) National Council of Youth Sports. Report on Trends and Participation in Organized
17 Youth Sports [report]. 2008. Available from:
18 <http://www.ncys.org/publications/2008-sports-participation-study.php>
- 19 (33) Nelson TF, Stovitz SD, Thomas M, LaVoi NM, Bauer KW, Neumark-Sztainer D.
20 Do youth sports prevent pediatric obesity? A systematic review and commentary.
21 *Curr Sports Med Rep*. 2011;10(6):360-70.

- 1 (34) Nilsson A, Andersen LB, Ommundsen Y, Froberg K, Sardinha LB, Piehl-Aulin K,
2 et al. Correlates of objectively assessed physical activity and sedentary time in
3 children: a cross-sectional study (The European Youth Heart Study). BMC Public
4 Health. 2009;9:322.
- 5 (35) NSW Department of Health. NSW Government Plan for Preventing Overweight and
6 Obesity in Children, Young People & their Families 2009 - 2011 [report]. 2009.
7 Available from:
8 http://www0.health.nsw.gov.au/pubs/2009/obesity_action_plan.html
- 9 (36) Pate RR, O'Neill JR. Youth sports programs: contribution to physical activity. Arch
10 Pediatr Adolesc Med. 2011;165(4):369-70.
- 11 (37) Patia-Spear L. Brain development and adolescent behavior. In: Coch D, Fischer
12 KW, Dawson G, editors. Human Behavior, Learning and the Developing Brain. New
13 York: The Guildford Press; 2007.
- 14 (38) Ridgers ND, Salmon J, Parrish AM, Stanley RM, Okely AD. Physical activity
15 during school recess: a systematic review. Am J Prev Med. 2012;43(3):320-8.
- 16 (39) Ridgers ND, Timperio A, Crawford D, Salmon J. Five-year changes in school
17 recess and lunchtime and the contribution to children's daily physical activity. Br J
18 Sports Med. 2012;46(10):741-6.
- 19 (40) Roemmich JN, Rogol AD. Physiology of growth and development. It's relationship
20 to performance in the young athlete. Clin Sports Med. 1995;14(3):483-502.

- 1 (41) Sacheck JM, Nelson T, Ficker L, Kafka T, Kuder J, Economos CD. Physical
2 activity during soccer and its contribution to physical activity recommendations in
3 normal weight and overweight children. *Pediatr Exerc Sci.* 2011;23(2):281-92.
- 4 (42) Sallis JF. Age-related decline in physical activity: a synthesis of human and animal
5 studies. *Med Sci Sports Exerc.* 2000;32(9):1598-600.
- 6 (43) Sherar LB, Cumming SP, Eisenmann JC, Baxter-Jones ADG, Malina RM.
7 Adolescent Biological Maturity and Physical Activity: Biology Meets Behavior.
8 *Pediatr Exerc Sci.* 2010;22(3):332-49.
- 9 (44) Steele RM, van Sluijs EM, Cassidy A, Griffin SJ, Ekelund U. Targeting sedentary
10 time or moderate- and vigorous-intensity activity: independent relations with
11 adiposity in a population-based sample of 10-y-old British children. *Am J Clin Nutr.*
12 2009;90(5):1185-92.
- 13 (45) Steele RM, van Sluijs EM, Sharp SJ, Landsbaugh JR, Ekelund U, Griffin SJ. An
14 investigation of patterns of children's sedentary and vigorous physical activity
15 throughout the week. *Int J Behav Nutr Phys Act.* 2010;7:88.
- 16 (46) Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical
17 Activity in the United States Measured by Accelerometer. *Med Sci Sports Exerc.*
18 2008;40(1):181-8.
- 19 (47) Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of accelerometer cut
20 points for predicting activity intensity in youth. *Med Sci Sports Exerc.*
21 2011;43(7):1360-8.

- 1 (48) Trost SG. State of the Art Reviews: Measurement of Physical Activity in Children
2 and Adolescents. *Am J Lifestyle Med.* 2007;1(4):299-314.
- 3 (49) US Department of Health and Human Services. Physical activity guidelines for
4 Americans [internet]. 2013. Available from:
5 <http://www.health.gov/PAguidelines/guidelines/chapter3.aspx>.
- 6 (50) Van Hoya AI, Fenton S, Krommidas C, Heuzé J-P, Quested E, Papaioannou A, et al.
7 Physical activity and sedentary behaviours among grassroots football players: A
8 comparison across three European countries. *International Journal of Sport and*
9 *Exercise Psychology.* 2013;11(4):341-50.
- 10 (51) Wickel EE, Eisenmann JC. Contribution of youth sport to total daily physical
11 activity among 6- to 12-yr-old boys. *Med Sci Sports Exerc.* 2007;39(9):1493-500.

12

13 Figure captions:

14 **Figure 1** The contribution of youth sport to daily weekend MVPA and VPA stratified by age
15 group and playing position.