

Effect of obstructive sleep apnoea on diabetic retinopathy and maculopathy

Leong, W B; Jadhakhan, F; Taheri, S; Chen, Y F; Adab, P; Thomas, G N

DOI:

[10.1111/dme.12817](https://doi.org/10.1111/dme.12817)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Leong, WB, Jadhakhan, F, Taheri, S, Chen, YF, Adab, P & Thomas, GN 2016, 'Effect of obstructive sleep apnoea on diabetic retinopathy and maculopathy: a systematic review and meta-analysis', *Diabetic Medicine*, vol. 33, no. 2, pp. 158-168. <https://doi.org/10.1111/dme.12817>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Supplementary materials:

**The association between obstructive sleep apnea on diabetic kidney disease:
a systematic review and meta-analysis**

Wen Bun Leong MBChB MRCP^{1,2}, Ferozkhan Jadhakhan MPH³, Shahrad Taheri
MBBS PhD FRCP^{1*,2*,4,5}, G Neil Thomas PhD^{6,7}, Peymané Adab MD⁶

1. School of Clinical and Experimental Medicine and Birmingham and Black Country NIHR CLAHRC, University of Birmingham
2. Specialist Weight Management Services, Heart of England NHS Foundation Trust, Birmingham UK
3. Primary Care Clinical Sciences, University of Birmingham, UK
4. Department of Medicine, Weill Cornell Medical College, New York USA and Doha Qatar.
5. Department of Medicine, King's College London, London, UK.
6. Public Health, Epidemiology and Biostatistics, University of Birmingham, UK
7. Institute of Public Health, Social and Preventive Medicine, Mannheim Medical Faculty, Heidelberg University, Mannheim, Germany

* Previous affiliations

Correspondence: Dr G Neil Thomas, Public Health, Epidemiology and Biostatistics, University of Birmingham B15 2TT, UK. Email: gneilthomas@yahoo.co.uk. Professor Shahrad Taheri, Weill Cornell Medical College, Education City, PO Box 24144, Doha, Qatar. Email: staheri@me.com.

Table 1: Eligibility criteria for systematic review

Inclusion criteria	Exclusion criteria
<i>Participants:</i> All adults with type 1 or type 2 diabetes mellitus (DM)	Children and adolescents (age below 18 years) Pregnant females Gestational diabetes mellitus Maturity onset diabetes of the young Latent autoimmune diabetes of adulthood No type 1 or type 2 DM
<i>Exposure:</i> Those with obstructive sleep apnea (OSA) or chronic intermittent hypoxemia (CIH)	Hypoxemia secondary to other respiratory disorders Chronic obstructive pulmonary disease Central sleep apnea Heart failure Asthma
<i>Comparator:</i> Adults without OSA or CIH	
<i>Outcomes:</i> Diabetic kidney disease (micro- and macro-albuminuria and chronic kidney disease assessed using glomerular filtration rate)	Renal impairment secondary to non-DM causes
<i>Study designs:</i> Cross sectional study Cohort study Case series	Non-human studies Single case reports Sample size < 10 participants Randomised controlled trials Intervention studies Experimental studies Systematic reviews/reviews Editorials Protocol papers Letters Guidelines/consensus statements
<i>Language:</i> All languages	

Search strategy:

Table 2: Search terms for MEDLINE, EMBASE and Cochrane Database

Search terms for diabetic kidney disease

1. Diabet\$.mp.
 2. Respiratory system/ or breathing disorder/ or snoring/ or hypoventilation/ or sleep/ or polysomnography/ or sleep apnea syndrome/ or breathing/ or sleep disordered breathing/ or sleep disorder/
 3. Hypoxemia/ or respiratory function disorder/ or anoxia/ or hypoxia/
 4. Somnolence/ or apnea/ or apnea hypopnea index/ or hypopnea.mp.
 5. 2 or 3 or 4
 6. 1 and 5
 7. Nephropathy.mp. or kidney disease/
 8. Albuminuria.mp. or proteinuria/ or albuminuria/
 9. \$albuminuria.mp.
 10. Renal failure.mp.
 11. Kidney function/
 12. Diabetic nephropathy/ or nephropa\$.mp. or chronic kidney failure/
 13. 7 or 8 or 9 or 10 or 11 or 12
 14. 6 and 13
-

The free text used for OpenGrey database were “obstructive sleep apnea” and “obstructive sleep apnoea”

The search terms for Zetoc database included “obstructive sleep apnea and diabetes”, “obstructive sleep apnoea and diabetes”, “apnea and diabetes” and “apnoea and diabetes”.

Table 3: Quality assessment form

Author:

Year:

Study ID:

Reviewer initials:

	Yes/No/ Unclear	Supporting evidence
Selection bias		
1. Does the study address an appropriate and clearly focused question?		
2. Does the study recruit consecutive patients?		
3. Are the cases and controls taken from comparable populations?		
4. Are the exclusion criteria the same for both cases and controls?		
5. Are the participants representative of the patient population?		
6. If applicable, is the control group comparable to cases (consider suitability, recruitment and baseline characteristics)?		
7. Is it clear that controls are not cases (in case control study)?		
Overall Judgement (Weak / Moderate /Strong)		
Respiratory measurement		
1. Did the study use a suitable measurement for OSA? (PSG/oximetry measured in standard, valid and reliable way)		
2. Is the scoring of the respiratory measures based on guidelines/consensus guidelines eg. AASM guidelines?		
3. Is there a clear definition of OSA?		
Overall Judgement (Weak / Moderate / Strong)		
Blinding		
1. Did the study blind the assessor performing the sleep analysis?		
Overall Judgement (Weak / Moderate / Strong)		
Study methods		
1. Are the attrition rates or missing data clearly documented?		
2. Did the study document the reasons for drop outs or missing data?		
3. Is this a retrospective or prospective design?		
Overall Judgement (Weak / Moderate / Strong)		
Analysis		
1. Are all the outcomes reported?		
2. Did the study adjust for confounding variables?		
Overall Judgement (Weak / Moderate /Strong)		
Overall judgement for the study (Weak, Moderate, Strong)		

Table 4: Criteria used for the diagnosis of obstructive sleep apnea and diabetic kidney disease assessment as reported by the included studies

Study	Definition of apnea/hypopnea or oxygen desaturation (OD)	OSA diagnosis	Min recording	DKD test	DKD diagnosis
Buyukaydin 2012 ¹	AASM criteria*	AHI≥5 events/hr	NA	24 hour urine albumin	Albuminuria ≥ 30mg/day
Furukawa 2013 ²	OD ≥3%	ODI≥5 events/hr	4 hours	Early morning urine ACR	Micro-albuminuria ≥3.4mg/mmol Macro-albuminuria ≥ 34.0 mg/mmol
Kosseifi 2010 ³	NA	NA	NA	NA	Micro-albuminuria
Laaban 2009 ⁴	Apnea = increased variation in suprasternal pressure with no airflow for ≥10 seconds; hypopnea = ≥50% reduction in airflow with ≥4% desaturation for ≥ 10 seconds	AHI≥5 events/hr	NA	24 hour urine albumin	Micro-albuminuria ≥ 30mg/day
Langrand 2014 ⁵	NA	NA	NA	NA	NA
Leong 2014 ⁶	AASM criteria*	AHI≥5 events/hr	4 hours	eGFR	eGFR <60ml/1.73m ² based on MDRD and CKD-EPI equations
Schober 2011 ⁷	Apnea = 80% reduction in airflow ≥ 10 seconds; hypopnea = 50-80% reduction in airflow with ≥ 4% desaturations for ≥ 10 seconds	AHI≥15 events/hr	NA	NA	NA
Storgaard 2014 ⁸	Apnea = 80% reduction in airflow ≥ 10 seconds; hypopnea = 50-80% reduction in airflow with ≥ 4% desaturations for ≥ 10 seconds	AHI≥5 events/hr	4 hours	24 hour urine albumin	Micro-albuminuria 30-300 mg/day Macro-albuminuria > 300mg/day (2 out of 3 samples)
Tahrani 2013 ⁹	AASM criteria*	AHI≥5 events/hr	4 hours	eGFR & urine ACR	MDRD eGFR <60ml/1.73m ² ACR >3.4 mg/mmol
Tanaka 2009 ¹⁰	OD ≥3%	ODI≥5 events/hr	NA	Creatinine (mg/dl)	NA
Zhang 2014 ¹¹	NA	AHI≥5 events/hr	NA	NA	NA
Zhang 2015 ¹²	Apnea = 80% reduction in airflow ≥	AHI≥5 events/hr	NA	Urine ACR or medical	ACR >300mg

10 seconds; hypopnea = 50-80%
reduction in airflow with $\geq 4\%$
desaturations for ≥ 10 seconds

history

OD = oxygen desaturation, min = minimum, OSA = obstructive sleep apnea, hr = hour, NA = not available, DKD = diabetic kidney disease, eGFR = estimated glomerular filtration rate, MDRD = Modification of Diet in Renal Disease, CKD-EPI = Chronic Kidney Disease Epidemiology Collaboration, ACR = albumin-creatinine ratio.*AASM = American Academy of Sleep Medicine criteria: apnea = complete cessation of airflow for ≥ 10 seconds; hypopnea = $\geq 30\%$ reduction in airflow $\geq 4\%$ drop in oxygen desaturation for ≥ 10 seconds.

Table 5: Summary of the results for diabetic kidney disease

Diabetic kidney disease	
OSA (based on AHI)	<p>10 studies (n=2927)^{1, 3-9, 11, 12} Adjusted: (+) 3 studies^{6, 9, 11}; (-) 0 studies</p> <p>Unadjusted : (+) 1 studies⁴; (-) 6 studies^{1, 3, 5, 7, 8, 12} Pooled OR (7 studies^{1, 4, 6-9, 12} – see Figure 1, supplementary materials) 1.59. 95% CI: 1.16 to 2.18 I²= 26.8%</p>
OSA (based on ODI)	<p>2 studies (n=1317)^{2, 10} Adjusted: (+) 2 study^{2, 13}; (-) 0 studies Pooled OR (2 studies^{2, 10} – see Figure 3, supplementary materials) 2.00. 95% CI: 1.36 to 2.94 I²= 0.0%</p> <p>Pooled OR (4 studies^{2, 9-11}, combined AHI & ODI as OSA diagnosis – see Figure 2, main manuscript) 1.73, 95% CI: 1.13 to 2.64, I²= 69.3%</p> <p>Unadjusted: No data</p>
%TST<90	<p>2 studies (n=158)^{5, 6} Adjusted: (+) 1 study⁶; (-) 0 study</p> <p>Unadjusted: (+) 1 study⁵; (-) 0 study</p>
Mean O₂	<p>2 studies (n=158)^{5, 6} Adjusted: (+) 0 study; (-) 1 study⁶</p> <p>Unadjusted: (+) 1 study⁵; (-) 0 study</p>
Minimum O₂	<p>3 studies (n=491)^{3, 6, 9} Adjusted: (+) 0 study; (-) 2 studies^{6, 9}</p> <p>Unadjusted: (+) 1 study³; (-) 0 study</p>

(+): reported significant association; (-): reported no significant associations; shaded cells indicate that no study reported data for the exposure-outcome combination. DR = diabetic retinopathy; OSA = obstructive sleep apnoea; AHI = apnoea-hypopnoea index; ODI = oxygen desaturation index; %TST<90 = percentage time spent under 90% oxygen saturation; O₂ = oxygen saturation, n = total number of participants

Meta-analysis

Three studies^{3, 5, 11} did not report on their results therefore a meta-analysis was carried out for seven studies^{1, 4, 6-9, 12} which reported unadjusted ORs on the association between OSA (defined using AHI) and DKD. The pooled estimates showed significant association (pooled OR 1.59, 95% CI: 1.16 to 2.18, $I^2=26.8\%$, Figure 1) using random effects analysis. The funnel plot of these studies^{1, 4, 6-9, 12} suggest an imbalance of small studies with positive results.




Figure 1: Forest plot of the association between obstructive sleep apnea and diabetic kidney disease using results from studies which reported unadjusted odds ratios and 95% confidence intervals. Random effects analysis performed using Stata 13. Fixed effects model yielded similar results (pooled OR 1.60, 95% CI: 1.25 to 2.05, $I^2=26.8\%$)




Figure 2: Funnel plot on studies which reported unadjusted odds ratios.




Figure 3: Forest plot of the **association between obstructive sleep apnea** diagnosed using ODI and diabetic kidney disease using results from studies which reported adjusted odds ratios and 95% confidence intervals.

References:

1. Buyukaydin B, Akkoyunlu ME, Kazancioglu R, et al. The effect of sleep apnea syndrome on the development of diabetic nephropathy in patients with type 2 diabetes. *Diabetes research and clinical practice* 2012;98:140-3.
2. Furukawa S, Saito I, Yamamoto S, et al. Nocturnal intermittent hypoxia as an associated risk factor for microalbuminuria in Japanese patients with type 2 diabetes mellitus. *European journal of endocrinology / European Federation of Endocrine Societies* 2013;169:239-46.
3. Kosseifi S, Bailey B, Price R, Roy TM, Byrd Jr RP, Peiris AN. The association between obstructive sleep apnea syndrome and microvascular complications in well-controlled diabetic patients. *Military medicine* 2010;175:913-6.
4. Laaban JP, Daenen S, Leger D, et al. Prevalence and predictive factors of sleep apnoea syndrome in type 2 diabetic patients. *Diabetes & Metabolism* 2009;35:372-7.
5. Langrand C, Glerant J, Gormand F, Guerin J, Moulin P. Association between obstructive sleep apneas and complications of type 2 diabetes. *Diabetologia* 2014;57:S499.
6. Leong WB, Nolen M, Thomas GN, Adab P, Banerjee D, Taheri S. The impact of hypoxemia on nephropathy in extremely obese patients with Type 2 diabetes mellitus. *J Clin Sleep Med* 2014;10:773-8.
7. Schober AK, Neurath MF, Harsch IA. Prevalence of sleep apnoea in diabetic patients. *Clinical Respiratory Journal* 2011;5:165-72.
8. Storgaard H, Mortensen B, Almdal T, Laub M, Tarnow L. At least one in three people with Type 2 diabetes mellitus referred to a diabetes centre has symptomatic obstructive sleep apnoea. *Diabetic medicine : a journal of the British Diabetic Association* 2014;31:1460-7.
9. Tahrani AA, Ali A, Raymond NT, et al. Obstructive sleep apnea and diabetic nephropathy: a cohort study. *Diabetes care* 2013;36:3718-25.
10. Tanaka SI, Akanuma Y, Ohashi Y. What is the prevalence of sleep apnea syndrome in Japanese patients with type II diabetes? JEDAS study. *Diabetes* 2009;58.
11. Zhang R, Ji L, Zhang P. Prevalence and associated factors of obstructive sleep apnea in hospitalized patients with type 2 diabetes in China. *Diabetes* 2014;63:A639.
12. Zhang R, Guo X, Guo L, Lu J, Zhou X, Ji L. Prevalence and associated factors of obstructive sleep apnea in hospitalized patients with type 2 diabetes in Beijing, China 2. *Journal of diabetes* 2015;7:16-23.
13. Goto K, Watanabe S. Large-billed crows (*Corvus macrorhynchos*) have retrospective but not prospective metamemory. *Anim. Cogn.* 2012;15:27-35.