

Therapeutic use of selective synthetic ligands for retinoic acid receptors

Brown, Geoffrey; Marchwicka, Aleksandra ; Cunningham, Alan; Marcinkowska, Ewa

DOI:

[10.1080/13543776.2016.1205586](https://doi.org/10.1080/13543776.2016.1205586)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Brown, G, Marchwicka, A, Cunningham, A & Marcinkowska, E 2016, 'Therapeutic use of selective synthetic ligands for retinoic acid receptors: a patent review', *Expert Opinion on Therapeutic Patents*, vol. 26, no. 8, pp. 957-971. <https://doi.org/10.1080/13543776.2016.1205586>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is an Accepted Manuscript of an article published by Taylor & Francis in *Expert Opinion on Therapeutic Patents* on 11th July 2016, available online: <http://www.tandfonline.com/doi/full/10.1080/13543776.2016.1205586>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Therapeutic use of selective synthetic ligands for retinoic acid receptors: a patent review

Aleksandra Marchwicka^{1*}, Alan Cunningham^{2*}, Ewa Marcinkowska¹ and Geoffrey Brown^{3#}.

¹ Laboratory of Protein Biochemistry, Faculty of Biotechnology, University of Wrocław, Joliot-Curie 14a, 50-383 Wrocław, Poland

² Institute of Immunology and Immunotherapy, College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

³ Institute of Clinical Sciences, College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

* These authors contributed equally to this work

Corresponding author: Dr Geoffrey Brown, tel. 0121 414 4082, g.brown@bham.ac.uk

Keywords: RARs, RXRs, retinoids, therapy, leukaemia, skin diseases

Introduction: Differentiation therapy using all-*trans* retinoic acid (ATRA) revolutionised the treatment of acute promyelocytic leukaemia to the extent this leukaemia is one of the most curable with ATRA and anthracycline-based chemotherapy providing cure rates above 80% (reviewed in [1]). Isotretinoin is used to treat chronic acne. Here, we examine the information described in recent patents and the extent to which new findings are leading towards extending retinoid-based differentiation therapy to other cancers and the development of new therapies for other disorders.

Areas covered: A search has been undertaken of the literature and of worldwide patents, filed during 2014 to the present time, in regard to synthetic agonists and antagonists of retinoic acid receptors and novel compositions for the delivery of these agents.

Expert opinion: New potential therapeutic applications have been described, including lung, breast and head and neck cancers, T cell lymphoma and neurodegenerative, metabolic, ophthalmic, muscle and inflammatory disorders. Recent patents have described the means to maximise retinoid activity. Two decades of efforts to extend retinoid-based therapies have been disappointing and new synthetic retinoids, target diseases and modes of delivery may well resolve this long standing issue.

Highlights

- There are highly selective agonists for all three RAR subtypes and antagonists except for RAR β .
- The tissue distribution of RAR subtypes is variable and functional redundancy is not the case.
- New patents describe the prospects of extending ATRA-driven differentiation therapy of acute promyelocytic leukaemia to other cancers.
- New patents have examined the prospects of broadening the therapeutic uses of retinoids to neurodegenerative, metabolic, ophthalmic, muscle and inflammatory disorders.

- New drug composition and delivery methods are important to improving the efficacy of retinoids.

1. Introduction

Natural retinoids are the ligands to different subtypes of retinoic acid receptors (RARs) and retinoid X receptors (RXRs). They are metabolically activated derivatives of vitamin A, which is delivered to the body by dietary sources. Active metabolites include the all-*trans*-, 9-*cis*- and 13-*cis*-retinoic acids (ATRA, 9cRA and 13cRA, respectively). The predominant natural retinoid is ATRA, which is present in most tissues and in blood serum. The one less abundant but detectable in almost all tissues is 13cRA. 9cRA is more difficult to detect, and the only organ in which currently available methods detect this metabolite is pancreas [2]. Retinoic acids play crucial roles in embryonic development and as to the behavior of adult tissues. In both these circumstances, retinoids are key modulators of cell growth, differentiation and apoptosis through their signaling pathways [3]. The different subtypes of RAR, their tissue distribution and the availability of novel synthetic analogues of retinoids that selectively agonize and antagonize a RAR subtype underlie the precise therapeutic use of retinoids. As outlined below, the tissue expression of subtypes is variable and functional redundancy is not the case in regard to the major subtypes.

2. Actions of retinoids *via* nuclear receptors

The biological activities of retinoids are mediated in target cells through two different classes of nuclear receptors: RARs and RXRs which are encoded by separate sets of genes [4]. There are three subtypes of RARs and of RXRs which are designated as α , β and γ . Several isoforms are generated by alternative splicing or alternative transcriptional start sites. In 2006, the International Union of Pharmacology recommended official names for these receptors and their subtypes. The agreed names for RAR α , RAR β and RAR γ are NR1B1,

NR1B2 and NR1B3, respectively, and for RXR α , RXR β and RXR γ are NR2B1, NR2B2 and NR2B3, respectively. The trivial, and more frequently used, names have been used throughout this manuscript. RARs and RXRs differ as to their ligand specificity. ATRA binds to and activates RARs and has a similar affinity for all three subtypes. 9cRA binds to and activates all three RARs and RXRs with different affinities [5]. The dual activation of RARs and RXRs by 9cRA is related to its flexible structure which allows conformational adaptation to the different binding pockets [6].

Like most members of the nuclear receptor superfamily, RARs are ligand activated transcription factors. They form heterodimers with RXRs and function as ligand-inducible transcription factors by binding to DNA sequences called retinoic acid response elements (RARE) within the promoter region of a target gene [5]. Activity of the heterodimer is controlled by RAR once ligand is bound; however RXR ligand can increase the transcriptional efficiency of the heterodimer [7]. The various isoforms of RAR and RXR create numerous receptor combinations. Moreover, the ability of RXRs to form heterodimers with numerous other nuclear receptors and to modulate a wide range of specific genes adds another level of complexity [8]. RARs can either repress or activate transcription [9,10]. Some of them act as repressors of transcription in the absence of the ligand, and upon binding of ligand the transcriptional repressors are released from the receptor complex and the interface for co-activators becomes accessible. Histone acetylase and RNA polymerase belong to the co-activation (CoA) complex, and, therefore, the chromatin structure becomes loose and the transcription of the target gene can start [11].

Recent observations suggest that, in addition to RARs, the retinoic acid signaling pathway involves the peroxisome proliferator-activated receptor β/δ (PPAR β/δ) [12]. PPAR β/δ is a

subtype of the PPARs which are ligand-activated nuclear receptors that are stimulated by small lipophilic ligands [13]. PPAR β/δ is expressed ubiquitously, with high expression observed in the brain, adipose tissue, skeletal muscle and skin [14]. Similar to RARs, activated PPAR β/δ forms a heterodimer with RXR and binds to specific PPAR response elements (PPREs) in the promoter regions of target genes [15]. Therefore, RARs and PPAR β/δ can activate different set of target genes and exert opposing activities. It has been shown that RARs trigger differentiation, cell cycle arrest and apoptosis [16], while PPAR β/δ promotes cell survival and proliferation [17,18]. A dual action of retinoic acid is related to different partitioning inside the cell between the two receptors. This process is regulated by the intracellular lipid binding proteins: cellular retinoic acid binding protein II (CRABP-II), and fatty acid binding protein 5 (FABP5) which deliver retinoic acid to RARs and PPAR β/δ , respectively. Thus, the cellular retinoic acid response depends on the CRABP-II/FABP5 ratio [19].

3. Tissue distribution of receptors

There are important differences as to the tissue distributions of RARs and RXRs. In humans, RXR β is ubiquitously expressed; RXR α is mainly expressed in the liver, lung, muscle, kidney, epidermis, and intestine and it is the major subtype in skin; and RXR γ is found in the brain, cardiac and skeletal muscle. As for the RARs, RAR α has a widespread expression pattern. In contrast, RAR β expression is prevalent in neural tissues and hardly detectable in skin; and RAR γ is expressed predominantly in the skin [20,21]. The distribution of RAR subtypes and their isoforms has been studied in different species using a variety of techniques. An overview of the distribution of RARs in different organs is presented in Table 1 and details of these studies are provided in the following sections. Besides physiological roles in normal tissues, retinoid receptors also play a role in the development of diseases

including cancers. Such is often due to mutations, chromosomal translocations, change to the level of expression, aberrant post-translational modifications, and epigenetic changes. These events result in altered function leading to disruption of homeostasis [22]. The abnormalities in retinoid signaling have been related mostly to dysregulation of RAR α and RAR β [23].

4. Subtypes and isoforms of retinoic acid receptors and their tissue distribution

4.1. Retinoic acid receptor α

The *RARA* gene is located on chromosome 17 and is composed of 10 exons and two promoters that give rise to the two isoforms RAR α 1 and RAR α 2 [24]. RAR α 1 has a broad tissue distribution, including the liver, spleen, kidney, prostate, spinal cord, cerebral cortex, uterus, ovary, testis and breast, and is considered as a canonical isoform [25]. In contrast, RAR α 2 is expressed in a limited number of tissues, for example, the intestine, lung and liver, and appears in the absence of ligand to be a more potent inhibitor of cell differentiation than RAR α 1. Its role in maintaining cells in an undifferentiated stem cell state [26] is seen in multiple myeloma whereby RAR α 2 expression is linked to primary multiple myeloma and drug resistance in this disease [27].

Cells of the hematopoietic system express mostly the RAR α and RAR γ subtypes and each has a specific modulatory role [17]. Although, disruption of any one of these two genes does not alter hematopoiesis, RAR α deficient mice demonstrated an impaired response to retinoids which leads to the accumulation of more immature granulocytes in their bone marrow after vitamin A treatment. Thus, RAR α can modulate granulopoiesis in response to retinoids [28]. Moreover, the double mutant mice RAR $\alpha^{-/-}$ RAR $\gamma^{-/-}$ die *in utero* and, therefore, it is hard to conclude whether RARs are needed for adult bone marrow hematopoiesis. However,

disruption of both genes affects granulocyte differentiation potential. The double mutant $RAR\alpha 1^{-/-}RAR\gamma^{-/-}$ cells derived from foetal liver were found to be blocked at the myelocyte/metamyelocyte stage of myelopoiesis [29]. On the other hand, $RAR\alpha^{-/-}RAR\gamma^{-/-}$ double mutant cells differentiate faster in response to G-CSF and SCF. These inconclusive results may be related to effects on different $RAR\alpha$ isoforms and implicate distinctions between the functions of $RAR\alpha 1$ and $RAR\alpha 2$. Oren and co-workers have suggested that $RAR\alpha 2$ may more effectively inhibit differentiation, and this becomes the predominant phenotype in $RAR\alpha 1^{-/-}$ cells [30]. Nevertheless, Zhu and co-workers demonstrated for murine progenitor cells that $RAR\alpha$ upregulation is required for optimal differentiation of primitive myeloid cells to granulocytes [31] and activity of $RAR\alpha$ is altered, by chromosomal translocations, in all cases of acute promyelocytic leukaemia (APL). A characteristic of this type of leukemia is the leukaemic cells are blocked at the promyelocyte stage of granulocyte differentiation [32]. The chromosomal translocation that fuses the PML gene to the $RAR\alpha$ gene gives rise to the protein PML- $RAR\alpha$ which acts as a constitutive transcriptional repressor [31]. However, a pharmacological dose of ATRA is able to dissociate co-repressors from the PML- $RAR\alpha$ bound to DNA to allow activation of transcription and, therefore, the differentiation of APL cells into mature granulocytes [33]. Resistance of leukemia cells to the differentiating effect of ATRA has also been correlated with aberrant or deficient phosphorylation of $RAR\alpha$ [34]. Appropriate phosphorylation of $RAR\alpha$ is important to retinoid signaling and activated RARs provoke signaling via non-genomic pathways by activating p38MAPK and its downstream target mitogen and stress-activated kinase 1 (MSK1). These events are required for cell differentiation. Additional disruption of $RAR\alpha$ has been observed in cells from acute myeloid leukemia (AML) patients whereby histones that associate with the *RARA2* promoter show a decrease in their overall acetylation and a

decreased level of dimethylation. These findings reveal that epigenetic changes to the landscape of *RARA* gene are involved in AML pathogenesis [33].

As to the involvement of $RAR\alpha$ in carcinomas, some estrogen receptor (ER)-negative breast cancer cells are resistant to retinoic acid and these cells have a reduced level of expression of $RAR\alpha$; overexpression of $RAR\alpha$ can restore ATRA-driven growth inhibition. $RAR\alpha$ can also participate in estrogen-mediated proliferation as $RAR\alpha$ has been shown to be induced by estrogens and shares a subset of binding regions with the estrogen receptor and, therefore, can be part of the estrogen receptor transcriptional complex [22]. As to ovarian cancer, the level of expression of $RAR\alpha$ has been correlated with ATRA-driven growth inhibition of ovarian cancer cell lines, but not for primary tumors [35]. $RAR\alpha$ can act also as an oncogene. This occurs in hepatocellular carcinoma whereby in the absence of the corepressor transcriptional intermediary factor 1 α (TIF1 α) oncogenesis correlates with the deregulation of retinoic acid signaling [36].

Recently it has been shown that retinoic acid is necessary for the function of the brain and new discoveries point to a crucial role in synaptic plasticity, learning and memory behaviors. $RAR\alpha$, $RAR\beta$ and $RAR\gamma$ have been detected in the adult brain. $RAR\alpha$ has a widespread distribution with high levels in the hippocampus, cerebellum and cortex. In contrast, $RAR\beta$ has a more restricted distribution that includes high levels in the striatum and the spinal cord, and $RAR\gamma$ is found at a low level in the hippocampus [37]. Recent findings suggest that age-related neuron loss, impairment of memory and cognition and even some neurodegenerative disorders are related to the disruption of retinoic acid signaling [37].

4.2. Retinoic acid receptor β

The *RARB* gene is located on chromosome 3 [38]. It is composed of 13 exons and 4 promoter regions which give rise to five distinct isoforms [39]. Expression of RAR β is rapidly induced by ATRA whereas expression of RAR α and RAR γ is sustained within cells at relative constant levels. In keeping with this observation, a RARE has been identified in the promoter region of RAR β 2 [40] and RAR α /RXR dimers bind to resulting in the expression of RAR β 2 [41]. Differences in the usage of *RARB1* and *RARB2* promoters and alternative splicing give rise to the major RAR β isoforms β 1, β 2, and β 4, and additional isoforms, such as RAR β 5 and RAR β 1', have been identified [42]. RAR β plays a key role in multiple diseases [5] and the various RAR β isoforms have different affinities for ATRA (at least with regard to RAR β 2 and RAR β 4) and different biological functions. For example, the RAR β 2 protein is a tumor suppressor whereas RAR β 4 has oncogenic properties [42].

Unlike RAR α , the tissue distribution of RAR β is more restricted and it is prevalent in epithelial tissues [40] and neural tissue and hardly detectable in skin [23]. RAR β 1 is considered to be a foetal isoform and plays a crucial role during development [39]. Whereas RAR β 1 is largely undetectable in adult tissues aberrant expression has been observed in several lung-cancer derived cell lines [38]. The recently identified RAR β 1' isoform, which arises from an alternative splicing of RAR β 1, is expressed in normal lung tissue and bronchial epithelial cells and the level of expression is suppressed in human lung cancers that are resistant to ATRA. Transfection of ATRA-resistant lung cancer cell lines and bronchial epithelial cells with RAR β 1' restored ATRA-sensitivity [43]. These data suggest that RAR β 1' plays a critical role in mediating the biological effects of retinoids in relation to carcinogenesis and may function as a tumor suppressor which is distinct from the observed function of RAR β 2 [38,44].

The RAR β 2 isoform is the major ATRA-inducible isoform of RAR [42] and it is considered as a canonical isoform of RAR β [43]. Loss of RAR β 2 activity, especially relating to gene hyper-methylation, has been observed in many different types of cancers, including head, neck [45] colon [46] non-small lung cancer [47], cervical cancer [44], breast [48] and prostate cancer [49]. Other studies have shown that transcriptional deregulation can silence RAR β 2 expression through decreased levels of co-activators, the presence of co-repressors or epigenetic mechanisms such as histone deacetylation. Expression of RAR β 2 also depends on the cellular level of retinoids as to the ATRA-inducible nature of this RAR [42]. Experiments *in vitro* as well as *in vivo* support the hypothesis that the ATRA growth inhibitory action is lost with impaired RAR β 2 expression. Berard and co-workers observed a higher incidence of pulmonary tumors in a truncated RAR β 2 mouse model. The endogenous RAR β 2 message level was found to be reduced in transgenic lung tissue and further reduced in the tumours [50]. Moreover, disruption of RAR β 2 in the teratocarcinoma cell line F9 impaired retinoic acid-mediated growth inhibition and differentiation [51]. On the other hand, endogenous reactivation of the RAR β 2 gene, by chromatin remodeling drugs, in breast cancer cell lines and xenograft tumors restored retinoic acid-dependent growth inhibition [52]. All of the above strongly suggest that RAR β 2 acts as tumor suppressor and loss of its expression may be an early and common event during malignant progression [41].

The RAR β 4 isoform is a splice variant of RAR β 2 and has a distinct functional attribute [53]. It lacks a DNA binding domain, and is unable to activate the transcription of target genes. Therefore, it has a dominant negative role [54]. ATRA-resistance of breast cancer cells is associated with down-regulation of RAR β 2 [55] and over-expression of RAR β 4. Findings from studies of breast cancer cells support the viewpoint that RAR β 2 is a potent inhibitor of

cell proliferation and that RAR β 4 interferes with RAR β 2-mediated growth suppression upon treatment of cells with ATRA [56].

The RAR β 5 isoform is similar to RAR β 4 and is a splice variant of RAR β 2 that lacks the domains A and B and part of domain C. RAR β 5 is able to form a functional hetero-complex with RXRs, but, as is the case for RAR β 4, RAR β 5 cannot bind to DNA and activate transcription of target genes [57]. RAR β 5 was initially found in epithelial cells but it is also expressed in breast epithelial cells and benign, premalignant and tumor cell lines. Moreover, it is preferentially expressed in estrogen-negative breast cancer cells that are resistant to the anti-proliferative action of retinoids [57].

4.3. Retinoic acid receptor γ

The *RARG* gene is located on chromosome 12. The major portion of the RAR γ protein, including the DNA and ligand binding domains, is encoded by seven exons that are identical for the RAR γ 1 and RAR γ 2 isoforms. These isoforms differ only in their N-terminal regions, the N-terminal region of RAR γ 2 shows high homology with that of RAR β and transcription of RAR γ 2 is regulated by its own promoter [58]. Lehmann and colleagues have shown that RAR γ 2, like RAR α 2 and RAR β 2, is activated by retinoids and RAR γ 1 is not able to drive ligand-dependent transcription. The poor transactivation function of RAR γ and ability to compete for DNA binding suggest that a major function of RAR γ 1 is to suppress gene activation by other RARs [59].

RAR γ 1 is predominantly expressed in skin where RAR γ 2 expression is low [60]. Both RAR α and RAR γ show differential expression throughout the epidermal layers whereby expression of RAR α and RAR γ is much higher in the spinous and granular layers in comparison to the

basal layer of epidermal cells [61]. In the case of keratinocytes, RAR γ is considered to be a tumor suppressor gene, as RAR γ was found to be absent from oral keratinocytes obtained from head and neck cancers. In addition, the incidences of premature skin aging and skin cancer, induced by ultra-violet radiation, have been correlated to a dramatic decrease in the level of RAR γ , due to proteasomal degradation of the receptor [22]. RAR γ can also act as an oncogene. In hepatocellular carcinoma, overexpression of RAR γ correlates with an increased cell survival through its altered subcellular localization. In this case, there is strong cytoplasmic localization of RAR γ which interacts with the phosphatidylinositol 3-kinase (PI3K) regulatory subunit p85 α leading to the activation of the PI3K/Akt pathway, which is one of the major survival pathways in cancer cells [62].

Though RAR γ has been found to be mainly expressed in skin, RAR γ is viewed as a critical regulator of the balance between haematopoietic stem cells (HSC) being able to maintain their stem cell status and these cells embarking on differentiation to produce mature blood cells [22]. The γ -knockout mouse has a reduced number of HSCs and loss of RAR γ also abrogated the capacity of ATRA to potentiate the maintenance of HSC in culture [63]. Agonizing RAR γ appears to promote self-renewal and/or proliferation of HSCs and, as such, opposes the ligand-driven action of RAR α to drive cells to differentiate. HSCs are still present in the knockout mouse and, like RAR α , the role of RAR γ is modulatory. Further evidence to support the notion that RAR γ is important to allowing cells to maintain their pluripotency comes from studies of the generation of induced pluripotent stem cells (iPSC) from somatic cells. The efficiency by which these cells can be generated can be increased by the addition of RAR γ to the Yamanaka cocktail of transcription factors used to generate iPSC [64].

5. Selective retinoic acid receptor ligands

Selectivity of ligand binding is conferred by the structure of the receptor's ligand binding domain (LBD). In the case of all RAR subtypes, LBD sequences are conserved as they differ only by three amino-acid residues. These differences are responsible for the different affinities of natural retinoids to RAR subtypes, and they allow the prospect of synthetic retinoids that are selective towards a particular subtype [65]. In the case of nuclear receptors, binding of the natural ligand to LBD induces conformational change which facilitates the binding of CoAs [66]. LBDs are composed of 12 α -helices and two short β -strands. After binding of the activating ligand the most mobile of all helices, helix H12 seals the ligand-binding cavity and creates the interface for binding of the transcriptional CoAs [65]. Some nuclear receptors, for example RAR α , act as transcriptional repressors in their un-liganded (apo) form. This is due to the short C-terminal part of H10, which adopts a β -strand conformation and creates a surface for interaction with transcriptional co-repressors (CoR) [67]. Upon binding of agonist, this region adopts a helical structure (H11), and promotes displacement of H12 into an active position. However, some synthetic ligands bind to RARs with high affinity but fail to stabilize helix H12 in an active position. These ligands are classified as partial agonists or antagonists, depending on their ability to prevent CoA recruitment. Antagonists not only destabilize the position of helix H12, but also prevent natural ligands from binding to LBD [66]. Some RAR antagonists, which are particularly efficient in stabilizing the β -strand conformation and favour CoRs binding, are defined as inverse agonists [67,68]. The possibility of fine-tuning the responses of particular RAR subtype and that isoforms are differentially expressed in tissues allows the therapeutic effects of synthetic retinoids to be directed against the required organs, and, accordingly, limit unwanted side-effects.

6. Therapeutic Applications of Retinoids

Longstanding interests as to development of the retinoid-based therapies have focussed attention to some extent on dysregulation of RAR-mediated signalling in leukaemia and other cancers and that skin cells express RARs and the prospect of treating a variety of skin disorders. In these two areas, patented developments within the retinoid field have delivered significant clinical benefits. Of particular importance is the standard use of ATRA and isotretinoin to treat APL and chronic acne, respectively. Recent patented findings bring about improvements to these existing applications and anticipate a broadening of the potential therapeutic use of retinoids as outlined below. The patents examined in this review are listed in Table 2 and structures of compounds are shown in Figure 1.

6.1. Dermatological Conditions

Retinoids are efficacious as to the treatment of dermatological conditions. Isotretinoin (13*cis*RA), an orally administered drug, is successfully used to treat severe forms of acne such as acne vulgaris, for which antibiotic and other retinoid treatments have failed. However, 13*cis*RA causes dry skin and is teratogenic. This retinoid can be given to women of child bearing age but contraception has also to be given 1 month before treatment, during the treatment and for one month after. Two contraception methods should ideally be introduced. Depression has also been associated to a low level with taking 13*c*RA [69]. However, this remains controversial [70] and recent studies have contradicted this association [71]. Taken together the side effects and patient compliance are restricting factors as to use. For some acne sufferers, less toxic retinoids are efficacious. Tazarotene (AGN190168) is a cream composition of a RAR β agonist that was developed by Allergan and alleviates acne in many patients. Retinoids are also a common ingredient to many cosmetic creams. Further applications of retinoids in dermatology include the use of acitretin, an oral retinoid, in the treatment of ichthyosis, lichen planus and psoriasis [72]. Acitretin is teratogenic and,

therefore, contraception has to be given to women of child bearing age 1 month before treatment, during the treatment and for 2 years after. Additionally, the RXR agonist bexarotene (LGD-1069) was licenced by the FDA in 2000 for use as an oral and topical treatment for cutaneous T-cell lymphomas and its associated lesions [73]. This compound can also bind to and activate RARs. Non-hormonal contraception is required in the case of LGD-1069 because medicament-activation of metabolic enzymes decreases the efficacy of hormonal contraception. A concern is that cases of hypothyroidism have been reported in patients given LGD-1069 [74,75]. The various retinoid-associated side effects lead to careful consideration of patient suitability, compound selection and regular monitoring of patients receiving treatment.

There has been little recent advancement with regard to new synthetic retinoids that are RAR subtype -selective for dermatological treatments. However, there have been advances regarding retinoid boosters. Granger and colleagues had demonstrated the ability of boosters to enhance the conversion of retinol and retinyl esters to retinoic acid in drug compositions, to enhance drug action [76-85]. More recently, Granger has described compositions of retinoid boosters that optimise the synergistic dermatological effects and has provided the respective concentrations that are required to maximise retinoid activity [86]. Interestingly, one new combination of tazarotene and dapsone (5% w/w), a neutrophil chemotaxis suppressant, resulted in a 13% greater reduction in acne associated comedonal lesions as compared to tazarotene alone. Additionally, combinations of tazarotene with either clindamycin (1% w/w) or benzyl peroxide (5% w/w) provided a 20% reduction in comedonal lesions in individuals.

6.2. Cancer

The most clinically significant application of retinoids to date is undoubtedly the use of ATRA to treat APL by overcoming the block to cell differentiation in this disease. What was once an untreatable illness is no longer so [87]; ATRA is the first in-line treatment for APL and often complemented with maintenance chemotherapy in the event of relapse or following refractoriness to ATRA. Considering this success and that RARs play an important role in regulating cell differentiation, proliferation and survival [88,89], there are ongoing efforts to extend ATRA differentiation therapy to other cancers. A limiting factor to use of ATRA to treat other cancers is an appreciable level of toxicity which can lead to fatal retinoic acid syndrome [90]. Moreover, extending the mechanism of ATRA's action in APL to other cancers is uncertain as to success in APL relates to the specific presence of the PML-RAR α fusion protein that blocks cell differentiation (see above and [32]). As to both APL and other cancers there is a need to develop retinoids that avoid the toxicities associated with ATRA and RAR subtype -selective analogues might provide the answer.

Recent findings indicate the use of retinoids in other leukaemias. Churchman and colleagues reported recently that alterations to the IKZF1 gene are linked to a stem cell-like phenotype and increased cell adhesiveness in BCR-ABL1-associated acute lymphoblastic leukaemia pre-B cells [91]. A consequence of these alterations was reduced responsiveness to tyrosine kinase inhibitor (TKI) therapy. A reversal of the stem cell phenotype could be achieved by disrupting cell clustering. Four hundred and eighty-three compounds were tested and revealed that ATRA, 9cRA, 13cRA and LGD-1069 are potent inhibitors of cellular aggregation, as observed by an abrogation of the formation of spheres in Arf^{-/-} BCR-ABL1 IK6-expressing pre-B cells. The retinoids and LGD-1069 also selectively induced expression of IKZF1 target genes and reduced colony forming potential, with LGD-109 significantly increasing cell responsiveness to TKI therapy.

There is the potential use of retinoids that are RAR subtype-selective to broaden retinoid treatment of cancer as evidenced by abnormal RAR signalling contributing to the pathogenesis of various carcinomas. Yan and colleagues have identified elevated levels of expression of RAR γ in hepatocellular carcinoma cell lines and primary tumours and the growth stimulatory effect of such was alleviated by treating these cells with ATRA [62]. Also, RAR γ has been shown to mediate ATRA-induced growth arrest and apoptosis of neoplastic mouse papilloma cell lines. Already, RAR subtype-selective retinoids have demonstrated their suitability for the treatment of malignancies other than APL and some are non-toxic [92,93]. One group has examined the possible use of retinoids to treat head and neck cancer (HNC). HNC comprises of cancers of the lips, larynx, pharynx, oral cavity, nasal cavity and treatment largely relies on surgery with the serious associated risks of perturbations to speech and the ability to swallow [94]. Previously, the RXR agonist LGD-1069 has shown potential efficacy in treating human T-cell lymphoma and lung cancer [92,93,95], and RAR γ agonists in suppressing tumour growth in mouse epidermal keratinocytes [96]. Gudas and colleagues have tested these agents individually and in combination in C57BL/6 mice bearing nitroquinoline-1-oxide (4-NQO)-induced oral cancers and significant benefit was observed [97,98]. Notably, gene expression changes induced by 4-NQO were prevented by LGD-1069 and the RAR γ agonist. Additionally, the agents did not increase triglycerides and suppressed reactive oxygen species production; Gudas suggested that these effects aid inhibition of tongue carcinogenesis. As to tumour migration, a combination of the above agents led to down-regulation of mRNA levels for a number of extracellular matrix proteases (MMPs) as well as the protein levels of MMP9, which point to a potential strategy for preventing metastasis. The inventors additionally used a tongue carcinogenesis model to show that LGD-1069 and CD-1530 inhibited HIF1 α signalling, a

pathway that regulates carcinogenesis, tumour development and migration [99] and overexpression of HIF1 α , which is known to relate to poor prognosis in HNCs [100]. Further indicators of effectiveness included a reduction in the level of β -catenin, which is normally increased in oral squamous cell carcinoma [101,102], fewer cancer stem cells within the oral cavity and reduced tongue tumour development in general.

In vitro results have indicated the potential efficacy of retinoids in treating breast cancer, but this has still to translate to an effective treatment for patients [103]. In contrast, histone deacetylase inhibitors (HDACi) have demonstrated efficacy in pre-clinical studies of breast cancer [104,105]. Additionally, HDACi have been found to synergise with retinoids to suppress the growth of breast cancer cells [106]. Miller and colleagues have created a hybrid molecule that is a derivative of TTNN (6-(5, 6, 7, 8-tetrahydro-5, 5, 8, 8-tetramethyl-2-naphthalenyl)-2-naphthalenecarboxylic acid, see Figure 1), which agonises both RAR β and RAR γ and inhibits histone deacetylase [107]. Hybrid 3 has been shown to have anti-proliferative effects *in vitro* against three breast cancer cell lines; MCF-7, SkBr3 and MDA-MB-231. Interestingly, the molecule was also shown to be active against the BT-20 breast cancer line which is double-negative for expression of ER and HER2 and insensitive to ATRA. Minimal effects were observed against non-tumour and normal mammary epithelial cells creating hope of a potential new treatment for breast cancer which appears not to affect normal breast cells.

6.3. Neurodegenerative Disorders

The pathology of Alzheimer's disease (AD) has been attributed to the deposition of amyloid plaque in neurological tissues [108]. The plaques comprise of amyloid β (A β) peptides [109,110], most of which are either A β 1-40 or A β 1-42 [111]. Critical to the creation of A β

peptide is the α -, β - and γ -secretase family of cell surface proteolytic enzymes. Sequential processing of amyloid precursor protein (APP) by β - and γ -secretases generates the pathologic A β peptides, whereas APP interaction with α -secretase, prior to β -secretase, yields a non-pathologic peptide. A disintegrin and metalloproteinase 10 (ADAM10) is the predominant proteolytic enzyme having α -secretase activity [112]. Increasing ADAM10 levels is, therefore, of interest as a therapeutic approach to AD. Retinol and its derivatives are known regulators of A β peptide formation and stimulate an increase in α -secretase activity [113]. A recent innovative combination of a byrostatin and ATRA, incorporated into a microsphere composition, has yielded an increase in α -secretase production in SH-SY5Y neuroblastoma cells as compared to bryostatin-1 treatment alone [114]. This novel finding points to the potential use of RAR-selective retinoids in combination with a bryostatin-1 for the treatment AD to mediate elevation of ADAM10.

There is evidence of a defect in retinoid transport in the brains of Alzheimer's patients [115] and the RAR agonist Am80 has been shown to reduce the level of A β in the brain of 5 month old APP23 mice [116]. Kawara and co-workers have pursued the avenue followed by Gudas and co-workers in the case of HNC [97] and examined the benefit to Alzheimer's patients of an RAR agonist used in combination with an RXR agonist [117]. Previously, LGD-1069 was found to enhance apolipoprotein E-dependent A β clearance alone [116] and the RAR agonist Am80 in addition to the RXR agonist HX630 reduced the level of A β significantly and strikingly reversed the defects in memory and spatial learning. As to the mechanism of repair, likely effectors include increased expression of the A β degrading enzymes, insulin degrading enzyme (IDE) and nuclear export protein (NEP).

Cell based therapies have been proposed as a non-drug method of promoting repair within the brain and neural tissues of individuals affected by neurodegenerative disorders [118].

However, there is limited availability of healthy brain tissue for obtaining neural stem cells and embryonic and iPSC are seen as an alternative source. Agonizing RAR γ is known to enhance the frequency of reprogramming of somatic cells to iPSC [64]. The use of iPSC to repair neuronal tissue is dependent on the means to drive iPSC towards neuronal development. The embodiments of a new patent describe the use of ATRA to induce human trophoblast stem (hTS) cells to differentiate into neural cells, creating the potential for *in vitro* testing of drug efficacy and safety and the development of cellular therapies for neurodegenerative conditions [119]. Post-treatment of hTS cells with 10 μ M ATRA, dopaminergic neurons, glutamine neurons, serotonergic neurons or GABA (γ -aminobutyric acid) neurons expressing the neural markers neurofilament, nestin and glial fibrillary acidic protein were obtained. An embryonic or iPSC based therapy has not been licenced for use to date, but this development implies a contribution that retinoids might provide to neural cell therapies and more broadly to the field of regenerative medicine.

6.4. Metabolic Disorders

The consequences of a high fat diet and/or vitamin A deficiency can be severe. Heart, kidney, liver, testes and pancreatic disease, as well as cancer, stroke, and diabetes are some of the common complications. The results contained in a recent patent position selective RAR β agonists as a potential medicament for such ailments [120]. Following administration of retinoids capable of agonizing RAR β , such as; AC261066, AC55649, LE-135, AGN190168, CD-271, CD-666, 9cRA, BMS641 and AGN191183 significant improvements to indicators of the aforementioned conditions were observed. Notably, in murine models of diabetes and other pancreatic diseases, insulin and glucagon sensitivities were either maintained or improved, and β -cell degeneration was suppressed. In the case of liver disease, decreases in alpha smooth muscle actin and hepatic reactive oxygen species as well as suppression of

hepatic stellate cell activation were observed, highlighting an ability to modulate the inflammatory response. The latter was further indicated by a reduction in inflammatory markers, such as monocyte chemoattractant protein 1 and TNF α . Additionally, RAR β agonist treatment was found to increase lecithin:retinol acyltransferase (LRAT) and decrease sterol regulatory element binding protein 1c levels in the liver. Of particular note, the highly specific RAR β agonist AC261066 led to reduced liver steatosis, indicating a preventative role in fat accumulation. The inventors also found that agonism of RAR β could restore vitamin A signalling in vitamin A deficient organs such as the liver. Another interesting finding was triglyceride levels were not found not to be elevated suggesting a safe way of targeting high fat diet associated conditions. One of the long term considered risks with the therapeutic use of retinoids has been their tendency to elevate serum triglyceride levels [121-123], leading to a risk of heart disease. While this patent describes LE-135 as a RAR β agonist it should be noted that a previous report has described the compound as having RAR β antagonist activity [124].

6.5. Graft versus host disease

Graft versus host disease (GVHD) is a complication of allogeneic haematopoietic stem cell transplantation in which donor T cells within the graft recognise host antigens as foreign resulting in T cell activation. This condition can be fatal, with the three year survival rates for grade 1 GVHD being 58% and 30% above grade IV [125]. ATRA regulates T cell differentiation in a bimodal manner. Early on post-antigen stimulation of T cells, ATRA signalling suppresses CD4⁺ve T cell differentiation towards T regulatory cells and provokes differentiation towards T helper 17 (Th17) and, thus, an inflammatory response. In contrast, ATRA signalling at a late stage during T cell responses suppresses the Th17 pro-inflammatory response. Given that vitamin A metabolism is upregulated during GVHD, a heightened level of ATRA-driven RAR α signalling exacerbates GVHD. That RAR α signalling is required for

acute GVHD [126] means that antagonism of RAR α is an interesting approach to ameliorating the side effects of or preventing GVHD. The embodiments of a new patent describe the use of the RAR selective antagonists, specified in US patents 5952345 and 5958954, to treat GVHD [127]. Inhibiting ATRA signalling in donor T cells attenuated GVHD lethality and importantly allowed the graft versus leukaemia effect to remain. Attenuation of ATRA signalling in GVHD patients or in patients immediately prior to transplant may, therefore, prove a useful means to manage and/or prevent GVHD. Indicated in the patent is that either a RAR α or RAR γ antagonist may be used, alone or in combination with an RXR agonist, to modulate the responsiveness of T cell donor cells. Given the bimodal action of selective retinoids in promoting and suppressing T cell inflammatory responses, a great deal of care will be required to ensure safe manipulation of the T cell response.

6.6. Ophthalmic Disorders

Keratoconjunctival disorders are inflammatory conditions that affect the cornea and conjunctiva. Defects to either can negatively impact on the other and the damage can be serious. Keratoconjunctival disorders follow long delays as to the recovery from ophthalmic conditions such as dry eye (sicca), keratitis, infection or corneal ulcers due to a secondary injury. Degradation of the stromal element of the corneal parenchymal tissue and particularly of collagen type I is a key element of the pathogenesis. The delay in repair and degradation to collagen results in scar formation which can obstruct vision. This places the prevention/repair of collagen degradation as a key target for therapeutic agents.

ATRA has been shown to promote corneal regeneration, albeit weakly and by a mechanism that is as yet unclear. The development of selective RAR agonists that can be used to efficiently treat keratoconjunctival disorders and which are safe could have major benefits.

Kimura and colleagues have observed that the RAR γ agonists R-677, CD-437 and BMS961 display a significant capacity to suppress collagen type I degradation when tested against rabbit keratinocytes. These agonists were also found to be effective in suppressing the collagen contraction during corneal cicatrization and conjunctival cicatrization [128].

6.7. Inflammatory Conditions

The precise mechanisms of alcohol induced liver disease (ALD) remain undetermined. However, the secretion of inflammatory cytokines, oxidative stress and lipid peroxidation toxicity of acetaldehyde are known to perpetuate an inflammatory reaction that results in cell apoptosis and tissue fibrosis [129-131]. The three stages of the disease include fatty liver disease (FLD), alcohol induced hepatitis (AIH), and finally liver cirrhosis [132]. While FLD is reversible with cessation of alcohol intake, inflammation in AIH leads to fibrosis and damage to areas normally involved in chemical detoxification, leading to irreversible scarring [133]. Finally, cirrhosis occurs often leading to the requirement for a liver transplant in severe cases. The prevalence of immune cells such as liver sinusoidal endothelial cells, Kupffer cells, dendritic cells, natural killer (NK) cells and natural killer T (NKT) cells in the liver [134] and the central role of inflammation in alcohol induced liver damage [135] lead to modulation of the immune response as an obvious therapeutic target in ALD. NKT cells are capable of promoting and suppressing the immune response and they represent an interesting means of regulating inflammation in ALD. In a 2014 patent, RAR selective retinoids and/or sulfatides were employed in a mouse model of ALD inflammation to suppress the pro-inflammatory activity of the type 1 NK cells and to activate the type 2 NKT cells [136]. The retinoids tested as to their capacity to inhibit the activity of the type 1 NKT cells included AGN190168, AGN190129 ATRA, retinol, 9cRA, 13cRA, AM580, AC55649, CD-1530, etretinate and acitretin. The investigations showed that injection of isotretinoin during or before the period of alcohol consumption was protective of liver damage. In particular, mice

given a high dose of alcohol experienced increased levels of alanine transaminase (ALT), while mice given a high dose of alcohol and ATRA showed no elevation of ALT. Additionally, histological examination for steatohepatitis in the liver revealed no evidence of fatty liver disease in the mice given ATRA.

Osteoarthritis (OA) results from breakdown to the cartilage lining the joints and to the underlying bone. Inflammation of the synovium generally ensues which exacerbates the existing pain from bone grinding [137]. The effects of retinoids are known to contribute to pain [138] and retinoids are involved in the pathological processes of OA [139]. The use of antagonists of RARs has been proposed as a means to alleviate OA-associated pain. To avoid any potential toxic effects of pan-antagonists, the inventors of a new patent have used pyrazole analogs as RAR γ selective antagonists to treat pain in OA [140]. Agonists were tested in the knee joints of rats in which the inflammatory agent monosodium iodoacetate had been administered resulting in an acute phase reaction. Using a capacitance test, the level of pain measured was found to be significantly inhibited in the rats treated with the analogues. The inventors may therefore have developed a method that could help alleviate pain generally, or specifically in osteoarthritis patients for whom other available treatments such as non-steroidal anti-inflammatory drugs have major side effects [141-143].

6.8. Muscle Damage

The causes of muscle injury are broad and common contributors include physical injury, infection, hypoxia due to vascular disruption and muscular dystrophy. Iwamoto and Pacifici have described the promotion of muscle repair by either the systemic or local administration of RAR γ agonists, or by pre-treatment of stem cells with RAR γ agonists prior to transplantation to the defect site [144]. The RAR γ agonists were of the group consisting; CD-

271, CD-394, CD-437, CD-1530, CD-2247, BMS270394 and BMS189961. Muscle defects were generated in mice and the systemic or local administration of RAR γ was performed at varying time points following the insult. The findings included that CD-1530 greatly increased repair to the defects, as compared to control mice in which the repair was largely fibrous scar tissue. Further support to the role of RAR γ in muscle repair is that agonism of RAR γ failed to promote acceleration of repair in γ -null mice. Of the RAR γ agonists tested, BMS270394 was found to be the most effective. As to the stem cells, pre-treatment of mesenchymal stromal cells with a number of RAR γ agonists was performed for periods of between 12 hours and 72 hours, and treated cells were administered at varying time points following muscle injury. Treatment on days 5, 7 and 10 following injury was found to be optimal for repair. The identification of treated mesenchymal stromal cells in the repair site was possible by DsRed labelling of the cells followed by fluorescent stereomicroscopy of histological sections. Overall it was found that local and systemic administration of the RAR γ agonist BMS270394 enhanced muscle repair, as did mesenchymal stromal cells pre-treated with this agonist for 72 hours and cells administered between days 5 and 9 following injury.

7. Drug Composition and Delivery Methods

Appropriate delivery methods are essential to the therapeutic application of selective retinoids. A prominent feature of recent retinoid patents has been in regard to drug composition and administration. Traditionally retinoids have been delivered either topically or orally. While these delivery methods still dominate, alternative applications are being developed. One such example has been in relation to the use of expanded polytetrafluoroethylene (ePTFE) artificial grafts to treat critical limb ischaemia. A complication of stenosis is neointimal hyperplasia and Ameer and colleagues have developed a wrap or gel comprising of a biocompatible polymeric matrix and a selective retinoid which

is placed around a section of a vascular implant [145]. The retinoid diffuses through the implant and into the vascular wall and has been seen to stop the proliferation of smooth muscle cells and neointimal hyperplasia and up-regulate anti-thrombogenic genes and the production of nitric oxide. This represents a promising method of improving the failure rate of ePTFE grafts in critical limb ischemia, particularly when healthy autologous vein availability is limited.

Retinoids can cause skin irritation when used topically [146]. Formulations of retinoids contained within liposomes and solid lipid nanoparticles have been found to be clinically equivalent to retinoid gels and the side effects are significantly reduced [147,148]. Despite these results, issues with composition stability have limited their development. Patent [149] describes a novel composition of retinoids contained within microcapsules that reduces irritation and improves the chemical and physical stability of the composition. A double phase retinoid release was also achieved, with the benefit of a reduced initial exposure of the skin to retinoids and the maintenance of an effective therapeutic concentration. The employed retinoids included ATRA, 13cRA, acitretin, AGN191183, CD-271, AGN190168, retinal and etretinate. Another approach that has been used to reduce irritation is the incorporation of a retinoid within an aqueous gel suspension. Difficulties in preparing a homogeneous retinoid distribution have hindered the use of this technology. However, hydrophobic silica is suited to overcoming this limitation [150]. Hydrophobic silica was found to be chemically and physically stable and when tested on the flanks of Gottingen mini-pigs, Draize scale test results indicated that the composition was non-irritant.

Another invention describes a formulation of retinoids contained within oleosomes. These are oily liquid globules which are coated in a layer of lipophilic surfactant, hydrophilic surfactant

and an anionic surfactant [151]. Retinoids employed in the composition included 3''-tert-butyl-4'-(2-hydroxyethoxy)-4''-pyrrolidin-1-yl-[1,1';3',1'']-terphenyl-4-carboxylic acid. Following application of 20µl of the formulation to the ears of BALB/C mice for 7 days, indicators of inflammation were examined by clinical observation and measurement of the ear thickness. Additional testing of the various formulations included epidermal penetration kinetics. A chemically and physically stable slow release formulation, named composition 1, provided preferential localisation within the epidermis and controlled retinoid release, and importantly maintained a therapeutic concentration.

8. Expert Opinion

The successful use of ATRA to treat APL demonstrated for the first time differentiation therapy of a malignancy that was previously incurable. An overarching importance of this approach to treating cancer is the prospect of milder treatment, alone or when coupled with gentler chemotherapy, which is tolerated by very old patients. Twenty three percent of the UK population is projected to be aged ≥ 65 by 2034, older patients are often excluded from new and aggressive chemotherapy trials and common and intractable fears from toxicity often lead to no treatment in very old age as the preferred/advised option. Two decades of endeavours have still to make provisions of retinoid-based differentiation therapies for other cancers, particularly carcinomas that are prevalent or difficult to treat in old age. Current efforts to find a means of extending differentiation therapy are, therefore, important to meeting a 21st century societal need. As emphasised below, choosing the right synthetic retinoid and targeting the right malignancy, or other disorder, are crucial as different synthetic retinoids will be effective for different diseases.

Advances towards extending retinoid-based differentiation therapy to a variety of cancers focus on the investigation of novel synthetic retinoids that specifically agonise or antagonise a particular retinoid receptor subtype(s). However, highly specific and selective synthetic retinoids have been available for quite some time [152], including antagonists [152,153] and, in particular, a RAR α subtype specific agonist [154]. In keeping, the use of novel synthetic retinoids as anti-cancer agents, as opposed to extending the use of ATRA, has been of interest for a decade, as brought to attention in 2001 by Altucci and Gronemyer [16]. In the case of APL, agonising RAR α , as opposed to all RARs by ATRA, is sufficient to differentiate APL cells and the use of an agonist of RAR α might well avoid ATRA-provoked retinoic acid syndrome which can be fatal. However, an agonist of RAR α is very unlikely to replace the use of ATRA to treat APL. This therapy is highly successful and clinical trial of an RAR α agonist would be restricted to a relatively small number of patients who have failed 1st and 2nd line treatments. Therefore, it is encouraging to see the use of retinoids to treat other cancers is being explored. The choice of the right synthetic retinoid is crucial as, for example, a RAR γ agonist and baxarotene have been shown to have a beneficial effect against n4-NQO-induced oral epidermal cancers in mice and a hybrid molecule which selectively agonises RAR β and RAR γ and inhibits histone deacetylase has been shown to be effective *in vitro* against breast cancer cell lines. Identifying the right RAR subtype-selective retinoid may provide new ways forward to differentiation therapy for malignancies other than APL.

ATRA has long been known to drive differentiation of keratinocytes. Administration of a retinoid topically provides a safe form of delivery and, for example, *Tazorac*® cream, containing the RAR $\beta\gamma$ agonist tazarotene, is a safe and effective treatment for adult and teen acne. An interesting way forward to the use of the localised release of retinoids is a patent that describes the benefit of placing a wrap/gel containing a selective retinoid around a

vascular implant to improve the success rate by preventing the proliferation of smooth muscle cells. Again the choice of synthetic analogue is crucial: agonizing RAR γ is beneficial to muscle repair as a recent patent describes a method for muscle regeneration by means of local or systemic administration of a RAR γ agonist. New methods of delivery and boosters to maximise activity, as describe in recent patents, are also important to these possible clinical developments.

Cancer and skin disorders have for quite a number of years been seen as targets for retinoid therapies. Albeit, retinoids are known to affect a wide variety of cell types and some recent patents extend the disease targets of interest in terms of retinoid-based therapies. Applications of retinoids to inflammatory, neurodegenerative, metabolic and ophthalmic disorders have been patented. The findings indicate a broadening of the possible therapeutic uses of retinoids. Moreover and to return to the theme of old age, retinoids may have a future in ensuring a health old age by preventing neurodegeneration and the shift in the immune system in older people towards an inflammatory profile, termed inflammaging. In this case, a chronic production of inflammatory cytokines appears to remodel the system. Whether this can be controlled by the systemic administration of an appropriate and safe retinoid is particularly interesting. All in all, the prospect of extending the current therapeutic uses of retinoids should be viewed as most promising in view of ongoing investigations of new synthetic analogues, the combined use of retinoids and other agents and a range of target clinical disorders.

Conflict of interest

The authors declare there are no conflicts of interest.

Acknowledgements

This project has received funding from the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreement no 315902. One author AC gratefully acknowledges receipt of a Marie Curie Research Fellowship. GB, EM, AM and AC are participants within the Marie Curie Initial Training Network DECIDE (Decision-making within cells and differentiation entity therapies). AM is the recipient of the grant PRELUDIUM No 2013/11/N/NZ3/00197 from National Science Centre in Poland. AC is a Marie Curie Research Associate. We thank Dr M. Chodynski, Pharmaceutical Research Institute, Warsaw for kindly drawing the structures for figure 1.



References

1. Lo-Coco F, Cicconi L, and Breccia M. *Current standard treatment of adult acute promyelocytic leukaemia*. Br J Haematol 2016; 172:841-54 * This paper provides an up to date review of the use of ATRA to treat APL.
2. Kane MA. *Analysis, occurrence, and function of 9-cis-retinoic acid*. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids 2012; 1821:10-20
3. Hayden LJ, Hawk SN, Sih TR, et al. *Metabolic conversion of retinol to retinoic acid mediates the biological responsiveness of human mammary epithelial cells to retinol*. Journal of cellular physiology 2001; 186:437-47
4. Zusi FC, Lorenzi MV, and Vivat-Hannah V. *Selective retinoids and rexinoids in cancer therapy and chemoprevention*. Drug discovery today 2002; 7:1165-74
5. Chambon P. *A decade of molecular biology of retinoic acid receptors*. FASEB journal : official publication of the Federation of American Societies for Experimental Biology 1996; 10:940-54 ** This paper provides a comprehensive overview of the biology of RARs.
6. Dollé P and Niederreither K. *The Retinoids: Biology, Biochemistry, and Disease*. John Wiley & Sons, 2015
7. Minucci S, Leid M, Toyama R, et al. *Retinoid X receptor (RXR) within the RXR-retinoic acid receptor heterodimer binds its ligand and enhances retinoid-dependent gene expression*. Molecular and cellular biology 1997; 17:644-55

8. Lane MA and Bailey SJ. *Role of retinoid signalling in the adult brain*. Progress in neurobiology 2005; 75:275-93
9. Hauksdottir H, Farboud B, and Privalsky ML. *Retinoic acid receptors beta and gamma do not repress, but instead activate target gene transcription in both the absence and presence of hormone ligand*. Molecular endocrinology (Baltimore, Md.) 2003; 17:373-85
10. Nagy L, Thomázy VA, Saydak MM, et al. *The promoter of the mouse tissue transglutaminase gene directs tissue-specific, retinoid-regulated and apoptosis-linked expression*. Cell death and differentiation 1997; 4:534-47
11. Aranda A and Pascual A. *Nuclear hormone receptors and gene expression*. Physiological reviews 2001; 81:1269-304
12. Shaw N, Elholm M, and Noy N. *Retinoic acid is a high affinity selective ligand for the peroxisome proliferator-activated receptor beta/delta*. J Biol Chem 2003; 278:41589-92
13. Oliveira AC, Bertollo CM, Rocha LT, et al. *Antinociceptive and anti-inflammatory activities of fenofibrate, an agonist of PPAR alpha, and pioglitazone, an agonist of PPAR gamma*. Eur J Pharmacol 2007; 561:194-201
14. Peters JM, Lee SS, Li W, et al. *Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta)*. Mol Cell Biol 2000; 20:5119-28
15. Desvergne B and Wahli W. *Peroxisome proliferator-activated receptors: nuclear control of metabolism*. Endocr Rev 1999; 20:649-88
16. Altucci L and Gronemeyer H. *The promise of retinoids to fight against cancer*. Nat Rev Cancer 2001; 1:181-93 **This paper provides an overview of the potential use of retinoids to treat cancer.
17. Adhikary T, Brandt DT, Kaddatz K, et al. *Inverse PPARbeta/delta agonists suppress oncogenic signaling to the ANGPTL4 gene and inhibit cancer cell invasion*. Oncogene 2013; 32:5241-52
18. Tan NS, Michalik L, Noy N, et al. *Critical roles of PPAR beta/delta in keratinocyte response to inflammation*. Genes Dev 2001; 15:3263-77
19. Schug TT, Berry DC, Shaw NS, et al. *Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors*. Cell 2007; 129:723-33
20. Germain P, Chambon P, Eichele G, et al. *International Union of Pharmacology. LXIII. Retinoid X receptors*. Pharmacological reviews 2006; 58:760-72
21. Huang P, Chandra V, and Rastinejad F. *Retinoic acid actions through mammalian nuclear receptors*. Chemical reviews 2014; 114:233-54
22. Duong V and Rochette-Egly C. *The molecular physiology of nuclear retinoic acid receptors. From health to disease*. Biochimica et biophysica acta 2011; 1812:1023-31
23. Sun S-Y and Lotan R. *Retinoids and their receptors in cancer development and chemoprevention*. Critical Reviews in Oncology/Hematology 2002; 41:41-55
24. Brand NJ, Petkovich M, and Chambon P. *Characterization of a functional promoter for the human retinoic acid receptor-alpha (hRAR-alpha)*. Nucleic acids research 1990; 18:6799-806
25. de The H, Marchio A, Tiollais P, et al. *Differential expression and ligand regulation of the retinoic acid receptor alpha and beta genes*. The EMBO journal 1989; 8:429-33
26. Leroy P, Krust A, Zelent A, et al. *Multiple isoforms of the mouse retinoic acid receptor alpha are generated by alternative splicing and differential induction by retinoic acid*. The EMBO journal 1991; 10:59-69

27. Wang S, Tricot G, Shi L, *et al.* *RARalpha2 expression is associated with disease progression and plays a crucial role in efficacy of ATRA treatment in myeloma.* Blood 2009; 114:600-7
28. Kastner P. *Positive and negative regulation of granulopoiesis by endogenous RARalpha.* Blood 2001; 97:1314-20 * This paper provides an opinion on the regulation of cell differentiation, particularly granulopoiesis, by RARs.
29. Labrecque J, Allan D, Chambon P, *et al.* *Impaired granulocytic differentiation in vitro in hematopoietic cells lacking retinoic acid receptors alpha1 and gamma.* Blood 1998; 92:607-15
30. Oren T, Sher JA, and Evans T. *Hematopoiesis and retinoids: development and disease.* Leuk Lymphoma 2003; 44:1881-91
31. Zhu J, Heyworth CM, Glasow A, *et al.* *Lineage restriction of the RARalpha gene expression in myeloid differentiation.* Blood 2001; 98:2563-7
32. Chen SJ, Wang ZY, and Chen Z. *Acute promyelocytic leukemia: from clinic to molecular biology.* Stem cells (Dayton, Ohio) 1995; 13:22-31
33. Schenk T, Stengel S, and Zelent A. *Unlocking the potential of retinoic acid in anticancer therapy.* British journal of cancer 2014; 111:2039-45
34. Bruck N, Vitoux D, Ferry C, *et al.* *A coordinated phosphorylation cascade initiated by p38MAPK/MSK1 directs RARalpha to target promoters.* The EMBO journal 2009; 28:34-47
35. Soprano DR, Qin P, and Soprano KJ. *Retinoic Acid Receptors and Cancers.* Annual Review of Nutrition 2004; 24:201-21
36. Khetchoumian K, Teletin M, Tisserand J, *et al.* *Loss of Trim24 (Tif1alpha) gene function confers oncogenic activity to retinoic acid receptor alpha.* Nat Genet 2007; 39:1500-6
37. Sodhi RK and Singh N. *Retinoids as potential targets for Alzheimer's disease.* Pharmacology, biochemistry, and behavior 2014; 120:117-23
38. Houle B, Pelletier M, Wu J, *et al.* *Fetal isoform of human retinoic acid receptor beta expressed in small cell lung cancer lines.* Cancer research 1994; 54:365-9
39. Toulouse A, Morin J, Pelletier M, *et al.* *Structure of the human retinoic acid receptor beta 1 gene.* Biochimica et biophysica acta 1996; 1309:1-4
40. van der Leede BJ, Folkers GE, Kruyt FA, *et al.* *Genomic organization of the human retinoic acid receptor beta 2.* Biochemical and biophysical research communications 1992; 188:695-702
41. Alvarez S, Germain P, Alvarez R, *et al.* *Structure, function and modulation of retinoic acid receptor beta, a tumor suppressor.* The international journal of biochemistry & cell biology 2007; 39:1406-15
42. Xu X-C. *Tumor-suppressive activity of retinoic acid receptor-beta in cancer.* Cancer letters 2007; 253:14-24
43. Petty WJ, Li N, Biddle A, *et al.* *A novel retinoic acid receptor beta isoform and retinoid resistance in lung carcinogenesis.* Journal of the National Cancer Institute 2005; 97:1645-51
44. Ivanova T, Petrenko A, Gritsko T, *et al.* *Methylation and silencing of the retinoic acid receptor-beta2 gene in cervical cancer.* 2002; 2:2-4
45. Youssef EM, Lotan D, Issa JP, *et al.* *Hypermethylation of the retinoic acid receptor-beta(2) gene in head and neck carcinogenesis.* Clin Cancer Res 2004; 10:1733-42
46. Youssef EM, Estecio MR, and Issa JP. *Methylation and regulation of expression of different retinoic acid receptor beta isoforms in human colon cancer.* Cancer Biol Ther 2004; 3:82-6

47. Muñiz-Hernández S, Hernández-Pedro N, Macedo-Pérez OE, *et al.* *Alterations in Retinoic Acid Receptors in Non-Small Cell Lung Cancer and Their Clinical Implications.* Journal of Cancer Therapy 2015; 06:648-64
48. Widschwendter M, Berger J, Hermann M, *et al.* *Methylation and Silencing of the Retinoic Acid Receptor- 2 Gene in Breast Cancer.* JNCI Journal of the National Cancer Institute 2000; 92:826-32
49. Nakayama T, Watanabe M, Yamanaka M, *et al.* *The role of epigenetic modifications in retinoic acid receptor beta2 gene expression in human prostate cancers.* Lab Invest 2001; 81:1049-57
50. Berard J, Laboune F, Mukuna M, *et al.* *Lung tumors in mice expressing an antisense RARbeta2 transgene.* FASEB J 1996; 10:1091-7
51. Faria TN, Mendelsohn C, Chambon P, *et al.* *The targeted disruption of both alleles of RARbeta(2) in F9 cells results in the loss of retinoic acid-associated growth arrest.* J Biol Chem 1999; 274:26783-8
52. Sirchia SM, Ren M, Pili R, *et al.* *Endogenous reactivation of the RARbeta2 tumor suppressor gene epigenetically silenced in breast cancer.* Cancer Res 2002; 62:2455-61
53. Nagpal S, Zelent A, and Chambon P. *RAR-beta 4, a retinoic acid receptor isoform is generated from RAR-beta 2 by alternative splicing and usage of a CUG initiator codon.* Proceedings of the National Academy of Sciences of the United States of America 1992; 89:2718-22
54. Chen LI, Sommer KM, and Swisshelm K. *Downstream codons in the retinoic acid receptor beta -2 and beta -4 mRNAs initiate translation of a protein isoform that disrupts retinoid-activated transcription.* The Journal of biological chemistry 2002; 277:35411-21
55. Swift CB, Hays JL, and Petty WJ. *Distinct functions of retinoic acid receptor beta isoforms: implications for targeted therapy.* Endocrine, metabolic & immune disorders drug targets 2008; 8:47-50
56. Sommer KM, Chen LI, Treuting PM, *et al.* *Elevated retinoic acid receptor beta(4) protein in human breast tumor cells with nuclear and cytoplasmic localization.* Proceedings of the National Academy of Sciences of the United States of America 1999; 96:8651-6
57. Nagpal S, Saunders M, Kastner P, *et al.* *Promoter context- and response element-dependent specificity of the transcriptional activation and modulating functions of retinoic acid receptors.* Cell 1992; 70:1007-19
58. Lehmann JM, Hoffmann B, and Pfahl M. *Genomic organization of the retinoic acid receptor gamma gene.* Nucleic Acids Res 1991; 19:573-78
59. Lehmann JM, Zhang XK, and Pfahl M. *RAR gamma 2 expression is regulated through a retinoic acid response element embedded in Sp1 sites.* Molecular and cellular biology 1992; 12:2976-85
60. Husmann M, Lehmann J, Hoffmann B, *et al.* *Antagonism between retinoic acid receptors.* Molecular and cellular biology 1991; 11:4097-103
61. Xu XC, Wong WY, Goldberg L, *et al.* *Progressive decreases in nuclear retinoid receptors during skin squamous carcinogenesis.* Cancer research 2001; 61:4306-10
62. Yan TD, Wu H, Zhang HP, *et al.* *Oncogenic potential of retinoic acid receptor-gamma in hepatocellular carcinoma.* Cancer Res 2010; 70:2285-95
63. Purton LE, Dworkin S, Olsen GH, *et al.* *RARgamma is critical for maintaining a balance between hematopoietic stem cell self-renewal and differentiation.* J Exp Med 2006; 203:1283-93 * This paper provides insight to the role of RAR γ .

64. Wang W, Yang J, Liu H, *et al.* *Rapid and efficient reprogramming of somatic cells to induced pluripotent stem cells by retinoic acid receptor gamma and liver receptor homolog 1.* Proc Natl Acad Sci U S A 2011; 108:18283-8
65. Gehin M, Vivat V, Wurtz JM, *et al.* *Structural basis for engineering of retinoic acid receptor isotype-selective agonists and antagonists.* Chem Biol 1999; 6:519-29
66. Alvarez S, Bourguet W, Gronemeyer H, *et al.* *Retinoic acid receptor modulators: a perspective on recent advances and promises.* Expert Opin Ther Pat 2011; 21:55-63
67. le Maire A, Teyssier C, Erb C, *et al.* *A unique secondary-structure switch controls constitutive gene repression by retinoic acid receptor.* Nat Struct Mol Biol 2010; 17:801-7
68. Germain P, Gaudon C, Pogenberg V, *et al.* *Differential action on coregulator interaction defines inverse retinoid agonists and neutral antagonists.* Chem Biol 2009; 16:479-89
69. Azoulay L, Blais L, Koren G, *et al.* *Isotretinoin and the risk of depression in patients with acne vulgaris: a case-crossover study.* J Clin Psychiatry 2008; 69:526-32
70. D'Erme AM, Pinelli S, Cossidente A, *et al.* *Association between isotretinoin and mood changes: myth or reality? An updated overview.* Int J Dermatol 2013; 52:499-500
71. Gnanaraj P, Karthikeyan S, Narasimhan M, *et al.* *Decrease in "Hamilton Rating Scale for Depression" Following Isotretinoin Therapy in Acne: An Open-Label Prospective Study.* Indian J Dermatol 2015; 60:461-4
72. Ormerod AD, Campalani E, and Goodfield MJ. *British Association of Dermatologists guidelines on the efficacy and use of acitretin in dermatology.* Br J Dermatol 2010; 162:952-63 * This paper provides a guideline to the use of retinoids to treat dermatological conditions.
73. Henney JE. *New drug for refractory cutaneous t-cell lymphoma.* JAMA 2000; 283:1131-31
74. Sherman SI, Gopal J, Haugen BR, *et al.* *Central hypothyroidism associated with retinoid X receptor-selective ligands.* N Engl J Med 1999; 340:1075-9
75. Smit JW, Stokkel MP, Pereira AM, *et al.* *Bexarotene-induced hypothyroidism: bexarotene stimulates the peripheral metabolism of thyroid hormones.* J Clin Endocrinol Metab 2007; 92:2496-9
76. Burger AR, Iwata K, Granger SP, *et al.* *Skin care compositions containing certain cyclic aliphatic unsaturated compounds and retinol or retinyl ester.* US5759556A (1998)
77. Burger AR, Zhang KH, Granger SP, *et al.* *Skin care compositions containing geranyl geraniol and retinol or retinyl esters.* US5756109A (1998)
78. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing an amide of a hydroxy fatty acid and a retinoid.* US5747051A (1998)
79. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing fatty acid amides, azoles, and retinol or retinyl ester.* US5716627A (1998)
80. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing fatty acid amides and retinol or retinyl ester.* US5811110A (1998)
81. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing dimethyl imidazolidinone and retinol or retinyl ester.* EP0745375B1 (1996)
82. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing fatty acid amides and retinol or retinyl ester.* US5599548 (1997)
83. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing an n-substituted fatty acid amide and retinol or retinyl ester.* US5955092A (1999)

84. Corey J, Dorogi PL, Meyers AJ, *et al.* *Cosmetic composition with a retinol fatty acid ester.* US5885595A (1999)
85. Granger SP, Rawlings AV, and Scott IR. *Skin care compositions containing melinamide and a retinoid.* US5693330A (1997)
86. Granger SP, Scott IR, Donovan RM, *et al.* *Method for Treating Skin with Retinoids and Retinoid Boosters.* US20140050676A1 (2014) **This patent describes new methods of retinoid delivery and boosters to improve retinoid efficacy.
87. Lengfelder E, Saussele S, Weisser A, *et al.* *Treatment concepts of acute promyelocytic leukemia.* Crit Rev Oncol Hematol 2005; 56:261-74 * This paper provides an overview of the concepts to successful treatment of APL.
88. Altucci L, Rossin A, Raffelsberger W, *et al.* *Retinoic acid-induced apoptosis in leukemia cells is mediated by paracrine action of tumor-selective death ligand TRAIL.* Nat Med 2001; 7:680-6
89. Clarke N, Jimenez-Lara AM, Voltz E, *et al.* *Tumor suppressor IRF-1 mediates retinoid and interferon anticancer signaling to death ligand TRAIL.* EMBO J 2004; 23:3051-60
90. Patatanian E and Thompson DF. *Retinoic acid syndrome: a review.* J Clin Pharm Ther 2008; 33:331-8
91. Churchman ML, Low J, Qu C, *et al.* *Efficacy of Retinoids in IKZF1-Mutated BCR-ABL1 Acute Lymphoblastic Leukemia.* Cancer Cell 2015; 28:343-56 ** This paper describes the possibility of extending the use of retinoids to the treatment of BCR-ABL1 acute lymphoblastic leukaemia
92. Dragnev KH, Ma T, Cyrus J, *et al.* *Bexarotene plus erlotinib suppress lung carcinogenesis independent of KRAS mutations in two clinical trials and transgenic models.* Cancer Prev Res (Phila) 2011; 4:818-28
93. Gniadecki R, Assaf C, Bagot M, *et al.* *The optimal use of bexarotene in cutaneous T-cell lymphoma.* Br J Dermatol 2007; 157:433-40
94. Olson ML and Shedd DP. *Disability and rehabilitation in head and neck cancer patients after treatment.* Head Neck Surg 1978; 1:52-8
95. Dragnev KH, Petty WJ, Shah SJ, *et al.* *A proof-of-principle clinical trial of bexarotene in patients with non-small cell lung cancer.* Clin Cancer Res 2007; 13:1794-800
96. Chen CF, Goyette P, and Lohnes D. *RARgamma acts as a tumor suppressor in mouse keratinocytes.* Oncogene 2004; 23:5350-9
97. Gudas LJ, Tang XH, Osei-Safro K, *et al.* *Combination therapy for head and neck cancer.* WO2015138354A1 (2015)
98. Tang X-H, Osei-Sarfo K, Urvalek AM, *et al.* *Combination of bexarotene and the retinoid CD1530 reduces murine oral-cavity carcinogenesis induced by the carcinogen 4-nitroquinoline 1-oxide.* Proceedings of the National Academy of Sciences of the United States of America 2014; 111:8907-12
99. Semenza GL. *Hypoxia-inducible factors in physiology and medicine.* Cell 2012; 148:399-408
100. Eckert AW, Lautner MH, Schutze A, *et al.* *Co-expression of Hif1alpha and CAIX is associated with poor prognosis in oral squamous cell carcinoma patients.* J Oral Pathol Med 2010; 39:313-7
101. Ueda G, Sunakawa H, Nakamori K, *et al.* *Aberrant expression of beta- and gamma-catenin is an independent prognostic marker in oral squamous cell carcinoma.* Int J Oral Maxillofac Surg 2006; 35:356-61
102. Pannone G, Bufo P, Santoro A, *et al.* *WNT pathway in oral cancer: epigenetic inactivation of WNT-inhibitors.* Oncol Rep 2010; 24:1035-41

103. Freemantle SJ, Spinella MJ, and Dmitrovsky E. *Retinoids in cancer therapy and chemoprevention: promise meets resistance*. *Oncogene* 2003; 22:7305-15
104. Rao R, Balusu R, Fiskus W, *et al.* *Combination of pan-histone deacetylase inhibitor and autophagy inhibitor exerts superior efficacy against triple-negative human breast cancer cells*. *Mol Cancer Ther* 2012; 11:973-83
105. Fortunati N, Catalano MG, Marano F, *et al.* *The pan-DAC inhibitor LBH589 is a multi-functional agent in breast cancer cells: cytotoxic drug and inducer of sodium-iodide symporter (NIS)*. *Breast Cancer Res Treat* 2010; 124:667-75
106. Emionite L, Galmozzi F, Grattarola M, *et al.* *Histone deacetylase inhibitors enhance retinoid response in human breast cancer cell lines*. *Anticancer Res* 2004; 24:4019-24
107. Miller WH, Gleason J, and Mader S. *Hybrid Molecule Having Mixed Retinoic Acid Receptor Agonism and Histone Deacetylase Inhibitory Properties*. US20140051760A1 (2014) * This patent describes a new hybrid molecule with activity against RARs and histone deacetylase inhibitory properties.
108. Murphy MP and LeVine H. *Alzheimer's Disease and the β -Amyloid Peptide*. *Journal of Alzheimer's disease : JAD* 2010; 19:311
109. Glenner GG and Wong CW. *Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein*. *Biochem Biophys Res Commun* 1984; 120:885-90
110. Masters CL, Simms G, Weinman NA, *et al.* *Amyloid plaque core protein in Alzheimer disease and Down syndrome*. *Proc Natl Acad Sci U S A* 1985; 82:4245-9
111. Lerner AJ, Gustaw-Rothenberg K, Smyth S, *et al.* *Retinoids for treatment of Alzheimer's disease*. *Biofactors* 2012; 38:84-9
112. Endres K and Fahrenholz F. *Regulation of alpha-secretase ADAM10 expression and activity*. *Exp Brain Res* 2012; 217:343-52
113. Koryakina A, Aeberhard J, Kiefer S, *et al.* *Regulation of secretases by all-trans-retinoic acid*. *FEBS J* 2009; 276:2645-55
114. Castor TP. *Combination therapeutics and methods for the treatment of neurodegenerative and other diseases*. WO2014085494 (2014) * This patent examines the possibility of treating neurodegenerative disorders with retinoids.
115. Goodman AB and Pardee AB. *Evidence for defective retinoid transport and function in late onset Alzheimer's disease*. *Proc Natl Acad Sci U S A* 2003; 100:2901-5
116. Cramer PE, Cirrito JR, Wesson DW, *et al.* *ApoE-directed therapeutics rapidly clear beta-amyloid and reverse deficits in AD mouse models*. *Science* 2012; 335:1503-6
117. Kawahara K, SUENOBU M, and Shudo K. *Médicament de traitement de la maladie d'alzheimer*. WO20141999905A1 (2014)
118. Jung YW, Hysolli E, Kim KY, *et al.* *Human induced pluripotent stem cells and neurodegenerative disease: prospects for novel therapies*. *Curr Opin Neurol* 2012; 25:125-30
119. Lee J, Lee TT, Lee U, *et al.* *Generation of neural stem cells from human trophoblast stem cells*. CN103561751 (2014)
120. Gudas LJ, Benoit Y, Perez R, *et al.* *Methods of treating diseases associated with high fat diet and vitamin a deficiency using retinoic acid receptor agonists*. WO2014113695A1 (2014)
121. Barth JH, Macdonald-Hull SP, Mark J, *et al.* *Isotretinoin therapy for acne vulgaris: a re-evaluation of the need for measurements of plasma lipids and liver function tests*. *Br J Dermatol* 1993; 129:704-7
122. Cisneros FJ, Gough BJ, Patton RE, *et al.* *Serum levels of albumin, triglycerides, total protein and glucose in rats are altered after oral treatment with low doses of 13-cis-*

- retinoic acid or all-trans-retinoic acid*. Journal of Applied Toxicology 2005; 25:470-78
123. Stoll D, Binnert C, Mooser V, *et al*. *Short-term administration of isotretinoin elevates plasma triglyceride concentrations without affecting insulin sensitivity in healthy humans*. Metabolism 2004; 53:4-10
 124. Li Y, Hashimoto Y, Agadir A, *et al*. *Identification of a novel class of retinoic acid receptor beta-selective retinoid antagonists and their inhibitory effects on AP-1 activity and retinoic acid-induced apoptosis in human breast cancer cells*. J Biol Chem 1999; 274:15360-6
 125. Sorrow ML, Martin PJ, Storb RF, *et al*. *Pretransplant comorbidities predict severity of acute graft-versus-host disease and subsequent mortality*. Blood 2014; 124:287-95
 126. Hall JA, Cannons JL, Grainger JR, *et al*. *Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor alpha*. Immunity 2011; 34:435-47
 127. Blazar B, Aoyama K, and Chandraratna R. *Treatment of graft-versus-host disease disorders using rar antagonists*. US20140194517A1 (2014) * This patent examines the potential use of RAR antagonists to treat graft versus host disease.
 128. Kimura K. *Therapeutic agent for keratoconjunctive disorders*. EP2918290A1 (2015)
 129. Lieber CS. *Alcoholic fatty liver: its pathogenesis and mechanism of progression to inflammation and fibrosis*. Alcohol 2004; 34:9-19
 130. Situnayake RD, Crump BJ, Thurnham DI, *et al*. *Lipid peroxidation and hepatic antioxidants in alcoholic liver disease*. Gut 1990; 31:1311-17
 131. Bataller R and Brenner DA. *Liver fibrosis*. Journal of Clinical Investigation 2005; 115:209-18
 132. Menon KVN, Gores GJ, and Shah VH. *Pathogenesis, Diagnosis, and Treatment of Alcoholic Liver Disease*. Mayo Clinic Proceedings 2001; 76:1021-29
 133. Gramenzi A, Caputo F, Biselli M, *et al*. *Review article: alcoholic liver disease--pathophysiological aspects and risk factors*. Aliment Pharmacol Ther 2006; 24:1151-61
 134. Crispe IN. *The Liver as a Lymphoid Organ*. Annual Review of Immunology 2009; 27:147-63
 135. Nagy LE. *The Role of Innate Immunity in Alcoholic Liver Disease*. Alcohol Research : Current Reviews 2015; 37:237-50
 136. Chaturvedi VK. *Prevention and treatment of inflammatory conditions*. US20140187504A1 (2014)
 137. Hutton CW. *Osteoarthritis: the cause not result of joint failure?* Ann Rheum Dis 1989; 48:958-61
 138. Pehlivan Y, Kisacik B, Sayiner ZA, *et al*. *Inflammatory back pain in patients treated with isotretinoin*. J Rheumatol 2011; 38:2690
 139. Davies MR, Ribeiro LR, Downey-Jones M, *et al*. *Ligands for retinoic acid receptors are elevated in osteoarthritis and may contribute to pathologic processes in the osteoarthritic joint*. Arthritis & Rheumatism 2009; 60:1722-32
 140. Bleisch TJ, Coates DA, Hughes NE, *et al*. *Substituted pyrazole analogs as RAR antagonists*. US20140275049A1 (2014)
 141. Masso Gonzalez EL, Patrignani P, Tacconelli S, *et al*. *Variability among nonsteroidal antiinflammatory drugs in risk of upper gastrointestinal bleeding*. Arthritis Rheum 2010; 62:1592-601
 142. Capone ML, Tacconelli S, Rodriguez LG, *et al*. *NSAIDs and cardiovascular disease: transducing human pharmacology results into clinical read-outs in the general population*. Pharmacol Rep 2010; 62:530-5

143. Trelle S, Reichenbach S, Wandel S, *et al.* *Cardiovascular safety of non-steroidal anti-inflammatory drugs: network meta-analysis.* BMJ 2011; 342:c7086
144. Iwamoto M and Pacifici M. *Composition and method for muscle repair and regeneration.* US20140303223A1 (2014)
145. Ameer GA, Kibbe M, and Webb A. *Controlled and Localized Release of Retinoids to Improve Neointimal Hyperplasia.* US20150071984A1 (2015) * This patent examines the potential use of retinoids to improve vascular grafts.
146. Geria AN, Lawson CN, and Halder RM. *Topical retinoids for pigmented skin.* J Drugs Dermatol 2011; 10:483-9
147. Shah KA, Date AA, Joshi MD, *et al.* *Solid lipid nanoparticles (SLN) of tretinoin: potential in topical delivery.* Int J Pharm 2007; 345:163-71
148. Patel VB, Misra A, and Marfatia YS. *Topical liposomal gel of tretinoin for the treatment of acne: research and clinical implications.* Pharm Dev Technol 2000; 5:455-64
149. Djedour A. *Microcapsules containing retinoids, method of preparing same, and pharmaceutical compositions containing same.* US20150190372A1 (2015)
150. Duprat A. *Aqueous-gel-type topical compositions in the form of a homogenous suspension of an active principle of the class of retinoids containing at least one hydrophobic silica.* US20150150974A1 (2015)
151. Djedour A. *Dermatological composition comprising oleosomes and retinoids, process for preparing the same and use thereof.* US20150147403A1 (2015)
152. Johnson AT, Klein ES, Gillett SJ, *et al.* *Synthesis and characterization of a highly potent and effective antagonist of retinoic acid receptors.* J Med Chem 1995; 38:4764-7 * This paper outlines the development of potent antagonists of RARs.
153. Klein ES, Pino ME, Johnson AT, *et al.* *Identification and functional separation of retinoic acid receptor neutral antagonists and inverse agonists.* J Biol Chem 1996; 271:22692-6
154. Teng M, Duong TT, Klein ES, *et al.* *Identification of a retinoic acid receptor alpha subtype specific agonist.* J Med Chem 1996; 39:3035-8

Figure 1 Structures of selective synthetic ligands for retinoic acid receptors

(B) shows the structures of RAR selective agonists and antagonists employed in patents. For comparison the structures of ATRA, 9cRA and 13cRA (A) and RXR selective agonists (C) are shown.

Table 1 Tissue distribution of the RAR subtypes and isoforms.


RAR isoform	Tissue	Technique used	Organism
RAR α 1	<i>Overexpressed in</i> haematopoietic cell lines; <i>also expressed in:</i> kidney, prostate, spinal cord, cerebral cortex hepatoma-derived cell line, liver, spleen, uterus, ovary, haematopoietic cell lines	Northern Blot assay	human
RAR α 1	Skin – spinous and granular layer	Non-radioactive in situ hybridization	human
RAR α 1	Brain, skin, intestine, muscle, heart, lung, liver, kidney	Northern Blot assay	mouse
RAR α 2	Intestine, lung, liver	Northern Blot assay	mouse
RAR β 1	Fetal (kidney, lung, skin) <i>Undetectable in</i> heart, colon, muscle, lung, spleen from adult	RNase protection assay	human
RAR β 1'	ATRA-resistant human lung cancer cell lines	PCR, immunoblotting	human
RAR β 2	epithelial tissues, cancer lung tissue neural tissue	RNAase protection assay, Southern blot analysis and Northern Blot assay	human
RAR β 2	<i>High:</i> Kidney, prostate, spinal cord, cerebral cortex hepatoma-derived cell line <i>Average:</i> liver, spleen, uterus, ovary, <i>Undetectable in:</i> haematopoietic cell lines	Northern Blot assay	human
RAR β 4	Breast cancer	Northern Blot assay, RT-PCR	human
RAR β 5	normal, premalignant, and malignant breast epithelial cells	RT-PCR, Western Blot	human
RAR γ	Skin – spinous and granular layer	Non-radioactive in situ hybridization	human
RAR γ 1	Skin - the predominant isoform	Northern Blot assay	mouse
RAR γ 2	Skin – very low level (<5%)	Northern Blot assay	mouse

Table 2 Examples of retinoid compounds employed in patents from 2014 to the present time.


Number & Patent	Chemical Name	Compound Name
US20140050676A1 “Method for treating skin with retinoids and retinoid boosters”	2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	AGN100335/Tretinoin (all- <i>trans</i> -retinoic acid), pan-RAR agonist
WO2015138354A1 “Combination therapy for head and neck cancer”	4-(6-Hydroxy-7-tricyclo[3.3.1.1 ^{3,7}]dec-1-yl-2-naphthalenyl)benzoic acid	CD-1530, RAR γ agonist
US20140051760A1 “Hybrid molecule having mixed retinoic acid receptor agonism and histone deacetylase inhibitory properties”	6-(5,5,8,8-tetramethyl-6,7-dihydronaphthalen-2-yl)naphthalene-2-hydroxamic acid	SR3957/TTNN hybrid 3 molecule, RAR agonist
WO2014085494 “Combination therapeutics and methods for the treatment of neurodegenerative and other diseases”		A retinoid derived from the group; AGN100335, retinol, retinol acetate, retinol palmitate, 13- <i>cis</i> -retinoic acid, and LGD-1069
WO2014199905A1 “Medicament for treatment of alzheimer's disease”	4-[[[(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)amino]carbonyl]benzoic acid	Am80, RAR α agonist
	4-(7,8,9,10-Tetrahydro-7,7,10,10-tetramethylbenzo[<i>b</i>]naphtho[2,3- <i>f</i>][1,4]thiazepin-12-yl-benzoic acid	HX630, RXR pan agonist
CN103561751 “Generation of neural stem cells from human trophoblast stem cells”	2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	AGN100335/Tretinoin (all- <i>trans</i> -retinoic acid), pan-RAR agonist
WO2014113695A1 “Methods of treating diseases associated with high fat diet and vitamin a deficiency using retinoic acid receptor agonists”	4-[4-(2-Butoxyethoxy)-5-methyl-2-thiazolyl]-2-fluorobenzoic acid	AC261066, RAR β 2 agonist
	4'-Octyl-[1,1'-biphenyl]-4-carboxylic acid	AC55649, RAR β agonist
	4-[(5,6,7,8-Tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)carboxamido]benzoic acid	AM580, RAR α agonist
US20140194517A1 “Treatment of graft-versus-host disease disorders using RAR antagonists”		RAR α and/or RAR γ antagonists (compounds not specified)
EP2918290A1 “Therapeutic agent for keratoconjunctive disorders”	(E)-4-(2-{3-[(1H-pyrazole-1-yl)methyl]-5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalene-2-yl}vinyl)benzoic acid	R-667, RAR γ agonist
	6-[3-(1-adamantyl)-4-hydroxyphenyl]-2-naphthalene acid	CD-437, RAR γ agonist
	3-fluoro-4-[2-hydroxy-2-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalene-2-yl)acetylamino]benzoic acid	BMS961, RAR γ agonist
US20140187504A1 “Prevention and treatment of inflammatory conditions”	6-[2-(3,4-Dihydro-4,4-dimethyl-2 <i>H</i> -1-benzothiopyran-6-yl)ethynyl]-3-pyridinecarboxylic acid ethyl ester	AGN190168/Tazarotene, RAR β γ agonist
US20140275049A1 “Substituted pyrazole analogues as RAR antagonists.”		Compound names not specified, RAR γ antagonists
US20140303223A1 “Composition and method for	3-Fluoro-4-[(R)-2-hydroxy-2-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-naphthalen-	BMS270394, RAR γ agonist

muscle repair and regeneration”	2-yl)-acetylamino]-benzoic acid	
	4-(6-Hydroxy-7-tricyclo[3.3.1.1 ³ ,7]dec-1-yl-2-naphthalenyl)benzoic acid	CD-1530, RAR γ agonist
	6-(4-Hydroxy-3-tricyclo[3.3.1.1 ³ ,7]dec-1-ylphenyl)-2-naphthalenecarboxylic acid	CD-437, RAR γ agonist
	6-[3-(1-adamantyl)-4-methoxyphenyl]-2-naphthoic acid	CD-271, RAR γ agonist
US201500719841 “Controlled and localised release of retinoids to improve neointimal hyperplasia”	2E,4E,6E,8E)-3,7-dimethyl-9-(2,6,6-trimethylcyclohexen-1-yl)nona-2,4,6,8-tetraenoic acid	AGN100335/Tretinoin (all- <i>trans</i> -retinoic acid), pan-RAR agonist
US20150190372A1 “Microcapsules containing retinoids, method of preparing same, and pharmaceutical compositions containing same”	3''-tert-Butyl-4'-(2-hydroxyethoxy)-4''-pyrrolidin-1-yl-[1,1';3',1'']-terphenyl-4-carboxylic acid	Company name not specified, selectively activates RAR γ relative to the subtypes α and β Protected in Patent Application WO2006066978
US20150150974A1 “Aqueous-gel-type topical compositions in the form of a homogenous suspension of an active principle of the class of retinoids containing at least one hydrophobic silica”	3''-tert-Butyl-4'-(2-hydroxyethoxy)-4''-pyrrolidin-1-yl-[1,1';3',1'']-terphenyl-4-carboxylic acid	Company name not specified, selectively activates RAR γ relative to the subtypes α and β . Protected in Patent Application WO2006066978
US20150147403A1 “Dermatological composition comprising oleosomes and retinoids, process for preparing the same and use thereof”	3''-tert-butyl-4'-(2-hydroxy-ethoxy)-4''-pyrrolidin-1-yl-[1,1';3',1'']-terphenyl-4-carboxylic acid	Company name not specified, selectively activates RAR γ relative to the subtypes α and β . Protected in patent application WO2006066978


A



1. all-*trans*-retinoic acid




2. 9-*cis*-retinoic acid




3. 13-*cis*-retinoic acid


B




4. AM80




5. AM580




6. TTNN Hybrid 3




7. AC261066




8. AC55649




9. CD-1530




10. AGN190168




11. CD-271




12. CD-437



13. R-667




14. BMS961




15. BMS270394

C



16. LGD-1069



17. HX630