

A systematic quality assessment of Environmental Impact Statements in the oil & gas industry

Anifowose, Babatunde; Lawler, Damian; van der Horst, Dan; Chapman, Lee

DOI:

[10.1016/j.scitotenv.2016.07.083](https://doi.org/10.1016/j.scitotenv.2016.07.083)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Anifowose, B, Lawler, D, van der Horst, D & Chapman, L 2016, 'A systematic quality assessment of Environmental Impact Statements in the oil & gas industry', *Science of the Total Environment*, vol. 572, pp. 570–585. <https://doi.org/10.1016/j.scitotenv.2016.07.083>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1. Introduction

1.1 Importance of the Problem

Much of the world economy is underpinned by the international oil and gas industry which is moving into a new phase of unconventional resources and competition from renewables (e.g. see Torres et al., 2016; Moustafa, 2016). However, environmental impacts of oil and gas activities on air, water, soil and ecosystems have been well documented (e.g. Lawler, 2005; Skierszkan et al., 2013; Anifowose and Odubela, 2015; Barcelo and Bennett, 2016; Landis et al., 2016).

On the international scene, Lawler (2005) found that poorly defined environmental changes, relevance and data quality issues were the major problems facing water resource management in oil-rich Azerbaijan and Georgia. A study of contaminants from Canada's Alberta heavy oil revealed some metal (i.e. arsenic, cadmium, nickel, vanadium) enrichment in soil samples within 20 km of oil facilities in the Cold Lake area (Skierszkan et al., 2013). In the US state of Colorado where active shale wells as of March 2015 are about 53,288, it is estimated that a minimum of 500 million m³ of water is required for hydraulic fracturing; and in 2013, up to 600 spills of produced water chemicals were reported (Barcelo and Bennett, 2016). Though, health impacts studies linking environmental health hazards with shale gas activities lack methodological rigour (Werner et al., 2015). New data have suggested that macroinvertebrate communities in north-central Arkansas are impacted by different levels of gas activity thereby prioritising the need for quantitative analyses of cumulative freshwater-impacts from oil and gas projects (Johnson

et al., 2015). In Cyprus, a study on liquefied natural gas (LNG) and pipeline network found key environmental impacts to include the release of particulate matter, odour/smell, noise and declining soil conditions as well as job opportunities (Papadopoulou and Antoniou, 2014).

In Nigeria, the case study country for this paper, an examination of 200 locations, 122km of pipelines and health records of 5,000 community members found significant environmental and health impacts following inadequate maintenance and decommissioning of oil and gas facilities in Ogoniland (UNEP, 2011). Giwa et al., (2014) report that communities adjacent to gas flaring sites in Nigeria often utilise the heat generated from the flare to dry farm produce and cloths; and to roast fish and maize. This increases the exposure of local people to noxious gases and other by-products emitted through the flaring processes which can result in environmental impacts and health problems including asthma, cancer, blood disorder and bronchitis, amongst others (see Davoudi et al., 2013). Nigeria has a long history of oil exploitation, spillage and pollution, particularly in the Niger Delta (UNDP, 2006; UNEP, 2011; Webb, 2011; Anifowose et al., 2012a). In addition, human injury and fatalities resulting from pipeline incidents are common here (e.g. Onuoha, 2007; Jasper, 2009; Aroh et al., 2010; Anifowose et al., 2012a). Ma'anit (2011) and Webb (2011) estimated that 9-13 million barrels of oil were spilt in the Niger Delta over the past 50 years (equivalent to one Exxon Valdez oil spill every year).

Environmental Impact Assessment (EIA) is a proactive methodical process that investigates and predicts the potential direct, indirect and cumulative impacts of proposed project activities on environmental receptors, ideally from project initiation to decommissioning, and offers mitigation strategies.

Produced as part of an EIA process, the Environmental Impact Statement (EIS) is a key document for reporting anticipated impacts of oil and gas projects, their mitigation and management plans. In most countries, the EIA process is part of the project permit or project approval procedure stipulated by the relevant authorities. Financial institutions like the World Bank, European Bank for Reconstruction and Development and the International Finance Corporation also require the submission of a detailed EIS as part of environmental due diligence for project financing (e.g. Lawler and Milner, 2005).

Over the years, significant awareness of environmentally sound processes and sustainable development have been promoted by EIA practice in large-scale infrastructural projects (Gilbuena et al., 2013; Cesar et al., 2014).

However, EIS appraisal studies show that quality is not always satisfactory (e.g. European Commission, 2009). Lawrence (1997) and Cashmore (2004) found that EIA practice has evolved without coherent conceptual theoretical and methodological foundation. Backlund (2009) stated that the quality of impact assessments in the EU suffered from applications of overly simple methodologies, and incomplete assessment of environmental impacts.

Eilperin (2010) and the National Commission (2011) found that major oil and

gas projects (e.g. BP's Macondo well drilling) were exempted from environmental impact analysis.

The effectiveness of the EIA system can be evaluated against the quality of EISs (Heinma and Poder, 2010), and/or regulatory compliance, adequacy of information and methodology, presentation of information and communication, objectivity, fairness and transparency (HMSO, 1996; Glasson et al., 1997; EC, 2001). A systematic quality review of EISs involves the sampling and methodical evaluation of several approved project EIS documents, using a set of review criteria (see section 2.1). Such quality reviews are common in non-oil and gas project sectors such as road construction, power and dam installations, mining activities and green-field developments. Performance review of EISs can help to strengthen quality control within EIA systems (e.g. Lee and Colley, 1992; UNEP, 2002; European Commission, 2009) especially when evidence-based methods are used (Backlund, 2009). There is a strong link between EIA process and EIS quality (Zhang et al., 2013).

Therefore, if the full strengths of EIA processes are to be realised, we suggest here that critical independent periodic reviews of the quality of EIA report (i.e. EIS) samples are essential. This should identify strengths and weaknesses which can be disseminated to encourage and share best practices in oil and gas developments. However, such reviews for the oil and gas industry are still rare today.

1.2 *Research Gap*

Several EIS quality review studies have been reported for non-oil and gas project sectors (see Table 1). Following periodic EIS quality review (at country or sector-level), a common feature of these studies is the hypothesis that project EISs tend to show improvement in quality over time.

However, the only known evaluation of EIS quality for oil and gas projects is the useful study of Barker and Jones (2013), commissioned by the then UK Department for Business Enterprise and Regulatory Reforms (DBERR). This focused solely on UK offshore petroleum production. Based on literature search from publicly accessible databases, our study here is believed to be only the second study which focuses on EIS quality for the oil and gas sector, and the only study anywhere which includes both onshore and offshore projects. In addition, none of the above cited studies in Table 1 apparently utilised inferential statistical analysis to evaluate their suggested EIS quality improvement. Hence, our study presented here.

1.3 Study Aims and Objectives

This paper has three unique aims. First, we address this international research gap by presenting a systematic analysis of the quality of EISs for both onshore and offshore oil and gas projects. Second, to our knowledge, this is the first assessment of its kind to examine the effectiveness with which decommissioning impacts are assessed within EISs of both oil and gas projects. Third, and again novel to our knowledge, this paper tests the

hypothesis that EIS quality of onshore and offshore oil and gas projects improves over time by using the 1998 to 2008 era (Table 2).

The study objectives are to:

- a. Develop an EIS quality review model for oil and gas projects, based partly on a review of methodologies adopted in previous studies of EIS quality review (see Table 3);
- b. Apply this model to assess the quality of 19 EISs produced between 1998 and 2008 covering *both* onshore and offshore projects (Table 2);
- c. Test the hypothesis that EIS quality for hydrocarbon projects in Nigeria has improved over the 1998-2008 period; and
- d. Make recommendations for future EIA practice in the Nigerian oil and gas industry, and in similar international contexts.

1.4 Environmental Regulation in the Nigerian Oil and Gas Industry

1.4.1 EIA Governance Structure

The evolution of EIA policy, practice and environmental awareness in Nigeria is closely tied to the oil and gas industry (Olokesusi, 1998; Ogunba, 2004). Understandably, when commercial oil and gas exploration began in the 1950s (Steyn, 2009), only simple environmental protection protocols and policies existed (Anifowose et al., 2014). To date, there are two main EIA governance structures in Nigeria's oil and gas industry; and we summarise below their evolution.

First, the inadequacies of reactive environmental control measures (e.g. the Petroleum Act of 1969), and the need for Nigeria to progress towards sustainable environmental practices, led to the establishment of the Federal Environmental Protection Agency (FEPA) by Decree 58 of 1988. FEPA had responsibility for the overall protection and sustainable use of Nigeria's environmental resources until its functions were transferred to the newly created Federal Ministry of Environment (FME) in 1999. Afterwards, FEPA ceased to exist and the FME was further empowered to formulate and review the National Policy on the Environment. Meanwhile, Nigeria's EIA decree 86 was enacted in 1992 and it mostly mirrors the US NEPA Act of 1969.

Second, the Petroleum Act of 1969 is the major regulatory framework in Nigeria that empowers the Department of Petroleum Resources (DPR) – a parastatal of the Federal Ministry of Petroleum Resources – as the responsible agency for environmental protection and pollution management in the oil and gas industry. The DPR also possesses the statutory responsibility for ensuring 'compliance to petroleum laws, regulations and guidelines' including those relating to health, safety and environment in conformity with national and international standards.

EIA consultants / preparers are often registered by both the DPR and FME. Any oil company or proponent seeking to undertake large scale project(s) in Nigeria's oil and gas industry has to seek EIA permit or approval from both the FME and DPR. This is in line with extant regulations such as the EIA decree 86 of 1992 and the Petroleum Act of 1969. This requirement appears to have inadvertently contributed to the widely reported frictions in environmental

regulatory duties in Nigeria's oil and gas industry. For example, these two agencies, though set up for a common goal, have been adjudged duplicative, overlapping and at times conflicting in their responsibilities (Olokesusi, 1998; Ogunba, 2004). However, in theory this double review could make for a more robust process since both agencies are expected to approve or disapprove projects based on the quality of EISs. Multijurisdictional approaches are not limited to Nigeria. For example in Canada both Federal and Provincial authorities have legislated EIA requirements which subject projects to multiple EIA processes (Leboeuf et al., 2010).

1.4.2 Other Regulatory Instruments

Apart from the EIA decree 86 of 1992 and the Petroleum Act of 1969, there are about 16 other regulatory controls that support EIA practice in Nigeria's oil and gas industry (see Agha et al., 2002). The major instruments are operated as follow:

1. *Federal Ministry of Environment (FME)*: The sectoral Guidelines for Oil and Gas Industry Projects (FEPA, 1995). These outline approaches for conducting EIAs in compliance with the requirements of the Nigerian EIA Decree 86 of 1992 which commits Nigeria to international best practices (Ameyan, 2008).
2. *Department of Petroleum Resources (DPR)*: The Environmental Guidelines and Standards for the Petroleum Industry in Nigeria (EGASPIN, 2002). Between 2000 and 2002, the DPR upgraded its EIA process to allow a systematic integration with project engineering

design and continuous monitoring of especially the implementation of designed mitigation (Agha et al., 2002).

3. The States and the Local Government Authorities are encouraged by the amended FEPA decree 59 of 1992 to establish environmental protection bodies. This has further complicated the EIA process, leading to State Environmental Protection Agencies (SEPAs) seeking to be involved at key EIA stages rather than just at the final review stage.

Indeed, Eneh (2011) has reported rivalries and jealousies at different levels of government while Emeseh (2012) found that conflict exists amongst federal and state parastatals over unclear jurisdiction on oil activities.

There are different versions of the Petroleum Industry Bill (PIB) and it is currently being replaced by the Petroleum Industry Governance Bill 2016 which passed its first reading in Nigeria's national assembly in April. The PIB sought to establish for the oil and gas industry a comprehensive legal framework which includes the protection of health, safety and environment (PIB, 2012).

To improve the enforcement of environmental regulations in Nigeria, an Act establishing the National Environmental Standards and Regulations Enforcement Agency (NESREA) was signed into law in 2007. Sections 7(g), (h), (j), (k), (l) and 8 (k), (l), (m), (n), (s) clearly exclude oil and gas from the purview of the Act but section 7(c) appears to have a contrary view. The

removal of the phrase 'oil and gas' from section 7(c) has been advocated by many (e.g. Ladan, 2012) as it appears to have been added in error.

2. Materials and Methods

2.1 Adaptation of the EIS quality review model

Past EIS quality review studies evaluated key performance criteria such as the completeness of information contained in each EIS; compliance with local, national and international standards; veracity and acceptability of approach(es) to impact analyses; and assessment of alternative options amongst others. In the past 20 years, these performance criteria have evolved as major yardsticks for grading the quality of EISs and the most widely used model criteria are Lee and Colley (Lee et al., 1999); the European Commission checklist (EC, 2001); and the IAU Oxford Brookes (HMSO, 1996). More details about these can be found in Table 3. We modelled our study on the Lee et al. (1999) framework because it is comprehensive, robust and widely used (see Table 1). To help adapt the Lee et al. (1999) model, we:

- (a) assessed the methods and structure of oil and gas project EISs housed at the European Bank for Reconstruction and Development (EBRD) in London;
- (b) ran a pilot application of the Lee et al. (1999) model on three representative Nigerian EISs and reviewed the results;
- (c) adapted and enhanced the Review Areas, Categories and Sub-Categories, including adding a new Review Area (RA) on 'Decommissioning/abandonment' (Fig. 1, Table 4) as informed by the pilot run. This addition to the EIS quality review model is important given the

environmental problems following the abandonment of oil facilities in Ogoniland Nigeria (UNEP, 2011), and also because international best practices for oil and gas project EIAs now include decommissioning (see FEPA, 1995; World Bank, 1999; DPR, 2002; Lawler, 2003; 2005; Ekins et al., 2006). Rising environmental standards make decommissioning a major challenge for oil companies (Strutt et al., 2006; Boothroyd et al., 2016).

- (d) reconciled the model with the basic requirements of Nigerian policy frameworks including the two main national environmental regulatory instruments guiding the oil and gas industry (section 1.4); and,
- (e) informed the model from international best practices, including HMSO (1996), Glasson et al. (1997), and World Bank Environmental Assessment Sourcebook (1999) for Energy and Industry Projects.

A summary of all modifications to the Lee et al. (1999) model is provided in Table 5.

2.2 EIS submissions and sample selection

Before EGASPIN in 1991, there were fewer than ten environmental study reports in the Nigerian oil and gas industry, including two pre-project EIAs and five post-impact assessment reports (Agha et al., 2002). However, between 1990 and 2001 about 130 EIA studies/EISs were recorded for the upstream sector alone as shown in Agha et al. (2002). During a field visit to Nigeria in April 2009, 19 EISs (~150-300 pages each) were selected: 14 through customized stratified random sampling from an EIA repository, and five directly from some Energy companies. The customised stratified random

sampling was done by picking EISs from the four directions of the repository. The EISs were not arranged or listed in any specified order therefore a gauged top, middle and lower strata were identified. Then simple random samples were taken from each stratum at the top, middle and lower piles.

Though there is hardly any oil or gas field development project without pipelines built into it, the customised stratified sampling approach ensured that the final sample contained pipeline projects and others alike. This approach took advantage of a *priori* knowledge of data access situation including how the EISs were emplaced, thereby improving the efficiency of the sampling design (e.g. see Claggett et al. 2010, p.338-9). Under such situation, a simple random sampling or any other sampling technique was not considered as useful as this customised stratified approach. Seawright and Gerring (2008) stated that choosing small sample in a completely random manner without prior stratification can lead to serious problems.

This sample represents approximately 9% of the estimated total submission at that time, and each sampled Project EIS is listed in Table 2. The depth and richness offered by this single-unit research design combined with the comparability of the 19 cases provide some useful trade-off regarding the limited representativeness of the study (Gerring, 2004). However, it is widely recognised that a truly representative case is difficult to identify (Seawright and Gerring, 2008). Nevertheless, large scale research with higher sample sacrifices depth for reliability (Gorard, 2006).

2.3 *Applying the adapted Lee et al. (1999) model*

We followed the ‘advice for reviewers’ in Lee et al. (1999, p.31) which contain the recommended systematic steps in the review process. The expanded version of the model is shown in Fig. 1. Figure 2 provides a schema of the process followed in applying the revised model to the 19 EISs listed in Table 2. The scoring of each EIS is based on how the criteria in Table 4 are adhered to or performed; and this scoring follows a similar approach used in the evaluation of design and safety engineering documents (e.g. see Strutt et al., 2006). The grading and scoring criteria used here are given in Table 6 and follow a similar principle to the tiered-classification (e.g. Excellent to Very Poor) reported in Boriani et al., (2010). The scoring approach is, no doubt, constrained by ontological and epistemological underpinning typical of many research studies. Table 6 explains the justification for the allocation of grades to EIS components or tasks. The enhanced result collation sheet (supplementary_folder_1) contains a translation script of the numerical equivalents to the grading symbols (A to F) (Table 6).

Typically, a ‘satisfactory’ EIS is awarded a grade of A, B or C depending on the extent to which the performance criteria (Table 4) have been met. For example, an EIS can be deemed ‘satisfactory’ if it clearly addresses the points below, amongst other things:

- evident prediction of impacts as deviation from baseline condition
- states confidence limits and uncertainties in data, and demonstrates quality control and quality assurance – these are rarely done in EISs

- provides clear evaluation criteria, technical and environmental justification for project alternatives
- balanced view e.g. considering not just negative impacts but also the positive ones and vice versa
- adherence to extant policy requirements (section 1.4) in the oil and gas industry; and application of international standards like the ISO 14000 series and World Bank Assessment Sourcebook for Energy Projects
- rigorous analyses and quantitative impact modelling (e.g. oil spill trajectory simulation and hydrodynamic modelling, sediment plume dispersion modelling, pipeline leak modelling using historical data) where necessary – less often done in EISs
- anticipates the level of success for the proffered mitigation measures
- field data collection, where relevant, adequately covers seasonal and inter-annual variability.

On the other hand, an ‘unsatisfactory’ EIS is awarded a grade of D, E or F depending on the extent to which the performance criteria have not been met. For example, an EIS can be deemed ‘unsatisfactory’ if it has some or all of the following, amongst others:

- lack of evidence to suggest that impacts have been defined as deviation from baseline condition;
- little or no indication of confidence limits, uncertainties or data gaps;
- the absence of quantitative modelling where this would have been beneficial;

- application of simple/basic impact prediction methods where more advanced methods would be most appropriate;
- none declaration of residual impacts;
- lack of justification for any unmitigated impacts;
- limited adherence to laid down policies and protocols on key project aspects such as decommissioning;

Strict privacy restrictions on access to these confidential EISs in this developing nation meant that only one reviewer could be used in this study. However, similar 'one-reviewer' approaches have been used widely and successfully e.g. Guilanpour and Sheate (1997), Gray and Edward-Jones (2003), and part of Peterson (2010). In common with usual practice, six (~32%) of the 19 EISs were re-reviewed four weeks after their initial review (Fig. 2). Re-reviews of EISs generally show repeatable results (e.g. Guilanpour and Sheate, 1997). For the occasional difference in grades here, EISs were re-reviewed once more to produce a final grade.

The Lee et al. (1999) model, as shown in Fig. 1, is hierarchical and consists of Review Areas (RAs), Review Categories (RCs) and Review Sub-Categories (RSCs). The quality assessment of RSCs determines the grading of RCs and RCs in turn determine the grading of the RAs. The five RAs represent the major areas of EIA activities considered necessary for oil and gas projects while the RCs are categories of activities that must be accomplished under each RA. The RSCs are the detailed basic level tasks that must be accomplished within each RC. Details of RAs, RCs and RSCs are in Table 4.

Assessment of these hierarchical topics and the overall EIS scores are then recorded using the revised score sheet shown in supplementary_folder_1.

2.4 Testing the hypothesis of EIS quality improvement over time

A Mann-Whitney U test at $p = 0.05$ was used to test the hypothesis that EIS quality has improved over time, based on the samples in Table 2. To do this, we compared the earlier half of the sampled EISs (1998-2004: $n = 10$) with the later half (2004-2008: $n = 9$). The difference in quality between the two periods was measured by the number of EISs which achieved at least 'satisfactory' grades in the EIS overall, and in all individual Review Areas.

Apart from the 14 EISs selected through customised stratified sampling, the five EISs retrieved from other sources (section 2.2) are not expected to significantly influence the statistical analysis here. Normally, the probabilities generated from statistical tests are estimates and the alpha (α) level (e.g. 0.05) for retaining or rejecting a null hypothesis has no mathematical or empirical relevance (Gorard, 2006). Effectively, statistical analyses rely on personal judgement in a way not different from qualitative data analysis.

According to Field (2009), an attempt to understand the representativeness of a given sample and how much variability there is between sample means requires approximations of the standard error. This is because, in reality, collecting hundreds or thousands of samples to construct a sampling distribution is only better imagined. More so, the basis of inference in randomisation models is the random assignment of cases to groups (e.g. lower, middle and upper), hence random sampling from certain population with specified distribution is not necessary (Ernst, 2004). Therefore,

inferences should be limited to the cases in the study. It is impossible to completely eliminate errors and bias in conducting research (Gorard, 2006; Field, 2009) and in rare occasions where errors are known, calculating their precise impact is often not feasible (Gorard, 2006).

3. Results

3.1 Overall Assessment for the Environmental Impact Statements

A key finding is that almost half (47%) of the 19 sampled EISs for the Nigerian oil and gas industry were rated as unsatisfactory in quality (i.e. grades D to F) (Fig. 3A). About three-quarters of the unsatisfactory EISs were graded D and the rest graded E. Ten EISs were satisfactory (i.e. grades A to C). Just 26% of these were graded 'Very Good' (all rated as B). Two EISs (11%) were scored as Poor (grade E) (Figure 3A). No EISs were graded in the extreme categories of A and F.

3.2 Quality of individual Review Areas (RAs)

Table 7 shows a full matrix of results for all the Review Areas (RAs) and Review Categories (RCs). Figure 3B presents grading for each Review Area across the 19 EISs. Clearly, the most successful area is RA 4 (Communication of Results), where satisfactory grades were achieved in 95% of the EISs. This is followed by RA 1 (Project Description/Baseline Environment) and RA 3 (Alternatives and Mitigation) where satisfactory ratings were returned for 63% and 58% of EISs respectively. The 'Identification and Evaluation of Key Impacts (RA 2)' – one of the most

important aspects of EIA – was found to be unsatisfactory in 58% of the sampled EISs. Importantly, the Decommissioning/Abandonment aspect of EISs (RA 5), a newly added Review Area (see Fig. 1), was scored as ‘unsatisfactory’ in 15 of the 19 EISs (see section 3.2.5).

3.2.1 Review Area 1 – Project Description and Baseline Environment

Analysis of the five Review Categories within RA 1 (Table 7) shows that ‘*Environment Description*’ was handled most successfully of all categories (and rated satisfactory in 95% of EISs), followed by ‘*Description of the Project Development*’ (79%). The ‘*Baseline Environment Condition*’ was satisfactorily described in 63% of the EISs, while ‘*Site Description*’ was satisfactory in 42% of them. ‘*Wastes and Residuals*’ were satisfactorily addressed in just 37% of the EISs.

3.2.2 Review Area 2 - Identification and Evaluation of Key Impacts (during site preparation, construction, operation)

Review Area 2 is crucial, but the ‘*Definition of Impacts as Deviation from Baseline Condition*’ (RC 2.1) was the weakest category in 74% of the EISs (Table 7). Also, 63% of the EISs unsatisfactorily addressed ‘*Impact Prediction*’ with only 11% of the EISs scoring very good grades while ‘*Impact Significance*’ was unsatisfactory in 68% of them. However, ‘*Impact Identification*’ was satisfactorily addressed in 74% and ‘*Scoping*’ was satisfactorily addressed in 68% of the EISs.

3.2.3 Review Area 3 - Alternatives and Mitigation

Table 7 shows that the *'Alternatives'* RC was generally handled satisfactorily in 95% of EISs, and 68% received very good grades. However, the *'Scope and Effectiveness of Mitigation'* category was scored as unsatisfactory in 63% of the 19 EISs. Also, *'Commitment to Mitigation'* was satisfactory in 63% of them, though this does not guarantee actual adherence to mitigation measures after project approval.

3.2.4 Review Area 4 - Communication of Results

The *'Layout'* and *'Presentation'* categories were satisfactory for all EIAs, with grades falling between A and C, although about only half (58%) of the former could be classed as very good (Table 7). In only 53% of the EISs was the *'Emphasis'* RC scored as 'Very Good'. Most EISs (89%) included satisfactory sections for the *'Non-technical Summary'*, and 79% were 'Very Good' (Table 7).

3.2.5 Review Area 5 – Decommissioning, Closure or Abandonment

In the Category of *'Addressing the Positives'* only 21% of EISs were rated as satisfactory (Table 7). This is despite the potential for oil and gas engineering structures to be recycled, or serve as artificial reefs, such as the 'rigs-to-reef' program in the Gulf of Mexico (Schroeder and Love, 2004) where about 93% of the 3000 decommissioned platforms has been removed completely and 7% already made into artificial reefs (Vedachalam et al., 2015). Similarly, the *'Addressing the Negatives'* category was scored as unsatisfactory in 79% of the EISs.

3.3 *Change in EIS quality over time*

Mann-Whitney *U* test on the number of 'Satisfactory' EISs does not provide statistically significant evidence (at $p < 0.05$) of an improvement in the overall quality of EISs between the earlier period (1998-2004; $n = 10$) and the later period (2004-2008; $n = 9$). The result was similarly non-statistically significant when the central break-point of the sample was adjusted to reduce the earlier period to 9 EISs. Moreover, the same test on the number of 'Satisfactory' individual *Review Areas* shows no evidence of an increase in quality over time. However, for the case with 9 EISs in the 1998-2004 period, a statistically significant increase in quality over time emerged for just one *Review Area*, namely 'Alternatives and Mitigation' ($p = 0.019$). In both cases of overall EIA quality and *Review Areas*, there is an assumption that any observed difference is partly explained by sampling variability (e.g. Sullivan and Feinn, 2012).

4. **Discussion**

4.1 *Implications of the overall results*

The trends on Figure 3 show that undertaking EIS quality review for oil and gas projects could be beneficial by unveiling areas needing adequate attention. For example, the non-oil and gas UK study by Glasson et al. (1997) revealed that EIA preparers and other stakeholders have learnt from experience. The overall results in Fig. 3 may not be unconnected with, amongst others, the absence of full scoping, unfocused assessment, inadequate baseline data and poor coverage of mitigation methods as found by Gray and Edwards-Jones (1999). In South Africa, a similar study

suggested that the quality of EISs in the North-West province is at par with international standards though not without some shortcomings (Sandham and Pretorius, 2008). Barker and Jones (2013) found that a number of EISs were unsatisfactory and cited a concern that the EIA process may have been driven by regulatory compliance instead of best practices. The overall EIS quality review results for this Nigerian study, as demonstrated in Fig. 3, is not any different. However, it is unclear if an empirically positive correlation exists between a country's national policy implementation compliance and its resultant environmental performance.

4.2 Drilling down into the quality of individual Review Areas

Interestingly, the level of satisfactory grades recorded for RAs 1 and 4 are reasonably consistent with findings of past EIS quality review studies for non-oil and gas projects (e.g. McMahon, 1996; Barker and Wood, 1999; Sandham and Pretorius, 2008; Sandham et al., 2008a; Kabir et al., 2010; Sandham et al., 2010). We discuss below the assessments of Review Categories within each individual Review Area.

4.2.1 Project Description and Baseline Environment

'*Environment Description*' and '*Description of Project Development*' are more straightforward tasks (i.e. more descriptive than analytical and less technical) for EIA teams and this explains the high ratings for these categories. For the '*Wastes and Residuals*' category, however, most EISs mentioned likely waste types but gave little or no information on the quantities of wastes to be generated or the timing and nature of waste generation. Also, the methods by

which waste quantity was derived, and their uncertainties and confidence limits were often unsatisfactorily addressed. Clearly, communication of uncertainty is important in environmental impact assessment (e.g. Wardekker et al., 2008; Butt et al., 2014).

Majority of the EISs demonstrated some evidence of biological and chemical data collection covering the wet and dry seasons. However, very little physical data (e.g. hydrological and hydraulic) were presented even when substantial water bodies were clearly at risk e.g. at pipeline river crossings (see Anifowose et al., 2012b; Anifowose et al., 2014; Lawler and Wilkes, 2015). In addition, most sampled EISs did not describe the 'Baseline Environment' as it could be expected to evolve should the project *not* proceed – despite 'baseline' being crucial in environmental risk assessment (Butt et al., 2014). Unsurprisingly therefore, this problem also resurfaced under impact identification and prediction (section 3.2.2). As for '*Site Description*', more than 50% of sampled EISs lacked a clear land-use map of the project environment. This is likely to hinder the study of the impacts of scale (spatial or detail) on the accuracy of impact prediction (e.g. João, 2002).

4.2.2 Identification and Evaluation of Key Impacts

With respect to RC 2.1, the absence of predictive modelling of impacts in most EIAs, as also previously noted by Ogunba (2004), weakens a score. Only five of the 19 EISs applied some predictive modelling (e.g. atmospheric dispersion of released gases; numerical simulation of sediment plume and oil spill; land-subsidence modelling due to hydrocarbon withdrawal etc.). Similar problems,

including non-testable and non-auditable predictions, have also been reported in the UK (Floater 2002), parts of Europe (Glasson et al. 2005), Australia (Buckley 1991, Warnken and Buckley 1998) and Canada (Bernard et al. 2001, Noble and Storey 2005). Impacts in the sampled EISs were not evidently determined as deviation from a baseline condition, therefore the ability to adequately predict future environmental impacts is compromised. This led to unsatisfactory grades in the *'Impact Prediction'* RC (Table 7). This supports similar findings in the Barker and Jones (2013) study and a British forest sector study by Gray and Edward-Jones (2003).

The majority of sampled EISs relied on 'expert' opinion and past experience in assessing impact significance, and lack a convincing or objective rationale. This largely makes the process subjective, as Sadler (1996) found in an international study of EIA effectiveness. Russo (1999) posited that over-reliance on experience is a major problem facing the conduct of energy sector EIA because the environment is constantly changing. Also, Glasson et al. (2005) argued that expert opinion, though useful, cannot provide totally defensible and flawless foundations for EIA impact prediction and significance evaluation. However, sophisticated predictive techniques may provide alternatives to this problem, but such advanced methodologies could restrict active participation of key stakeholders in the EIA process (Weston, 2004; Glasson et al., 2005).

4.2.3 *Alternatives and Mitigation*

For oil and gas exploration, '*Alternative*' options arguably are restricted in location but several options exist for the process technologies, especially extraction, and oil and gas transportation. Some of the EISs focused more on economic than environmental issues when examining alternatives.

A key point, however, is that residual impacts and uncertainties of '*Mitigation Effectiveness*' were hardly addressed. Such confirmation of '*Mitigation Effectiveness*' requires, of course, detailed post-project monitoring, although this can be difficult to achieve (e.g. Noble and Storey, 2005; Nasen et al., 2011). There is some international recognition of the need for better implementation of mitigation measures for energy-related projects (e.g. E&P Forum/UNEP, 1997; Russo, 1999; Khadka and Khanal, 2008; Klevas et al., 2009). Indeed, Wawryk (2002) and Marazza et al., (2010) suggest that Environmental Management Systems (EMS) (*an environmental protocol often voluntarily adopted by multinational oil companies*) could help here given their focus on change over time and the monitoring of mitigation plans and environmental impacts. Perhaps, implementation of mitigation is where the rather controversial NESREA Act (see Ladan, 2012; Ambituuni et al., 2014) detailed in section 1.4 could become useful. Fortunately, Nigeria's DPR has already enhanced its EIA process to allow a systematic integration with project engineering design and continuous monitoring of the implementation of designed mitigation. This, with similarly robust complements from the FME, could mean that the exclusion of oil and gas activities from NESREA Act is apt.

4.2.4 Communication of Results

Review Area 4 was the strongest part of the EISs examined, perhaps because it is a simpler task and less technical unlike impact prediction. Nevertheless, many environmental baseline chapters were disproportionately lengthy, yet none of the EISs presented clear chapter summaries. Also, many of the EISs cited largely old references (e.g. 1960s-1980s) relative to their submission dates.

4.2.5 Dealing with project decommissioning impacts

This was the worst performed of the five Review Areas (see Table 7; Fig. 3B). The project decommissioning sections in 79% of the EISs did not apparently follow requirements set out by, for example, FEPA (1995). Very few EISs provided clear outlines of how contaminated environmental receptors would be remediated or replaced. Instead, most EISs only made reference to the relevant sections of the DPR EGASPIN (2002) or its earlier version; or to FME FEPA (1995) with promises of adherence. This is unfortunate, given the UNEP (2011) report which demonstrated that inadequate handling of oil and gas asset decommissioning contributed to environmental problems in Ogoniland – a similar point reinforced by Schroeder and Love (2004) in the Gulf of Mexico. Similarly, a recent study found high concentrations of soil gas methane above decommissioned oil and gas wells in the UK and suggested that this could be due to well integrity failure where facilities have been inappropriately decommissioned (Boothroyd et al., 2016). Clearly, asset decommissioning in Nigeria (and possibly elsewhere) needs more thorough

attention in EISs including a step-change that involves the integration of EMS and project-level HSE Management Systems (see Wawryk, 2002).

The rather weakly assessed decommissioning impacts is not surprising given the misconception about project decommissioning in EIAs. Russo (1999) stated that there are uncertainties about the necessity for an anticipation of possible environmental impacts of decommissioning in EIAs of energy projects since such project lifespans are substantial. But a potential solution to this problem is to treat EISs as 'living documents' which will allow updates to relevant aspects (e.g. decommissioning) as technology and science evolve. In the Desire Oil Plc. exploratory drilling project in the UK (Palframan, 2010; p.3), the EIS through its Environmental Management Plan (EMP) was integrated with the ISO 14001 EMS and the HSE Management System. Recent studies have emphasised the significance of EIA-EMS integration as a fundamental step-change towards addressing such project environmental concerns (e.g. Palframan, 2012; Perdicoulis et al., 2012; Raissiyan and Pope, 2012).

4.3 Discussing the hypothesis of EIS quality improvement (1998-2002)

The lack of statistically significant improvement in EISs quality is surprising. It was hypothesised that EIS quality would rise over time as per the previous studies (sections 1.2, 1.3), and driven by at least four temporal changes like: (a) emerging environmental/EIA legislation e.g. the EIA Decree of 1992 and EGASPIN (2002); (b) emerging analytical methods and improved Information Technology through time, including software availability, enhanced instrumental/remote sensing platforms (e.g. NigeriaSat-1) for data collection;

(c) greater availability of longer and more representative datasets of enhanced quality through time; and (d) increased experience with EIA analytical and reporting methods within oil and gas companies.

This result is intriguing, given the improvement over time suggested for non-oil and gas project EISs, albeit without statistical analysis (e.g. see Table 1).

Based on our study sample (section 2.2), one explanation for the lack of statistically significant improvement over time in Nigerian oil and gas EISs is perhaps the absence of independent monitoring/periodic EIS quality review.

With this, independent review feedback into the EIA system can enhance the practices and processes of EIA (Lee and Brown, 1992; Glasson et al., 1997).

However, the increase in quality over time observed in the 'Alternatives and Mitigation' Review Area could have been driven by the fact that 'alternatives' are mostly restricted to specific locations (e.g. hydrocarbon reservoir sites).

And alternative processes and designs adopted are dependent upon evolving technologies. Our analyses and key findings did not support the popular hypothesis regarding EIS quality improvement over time. Nevertheless, our study reported herein, and those of the non-oil and gas projects cited above, provide some empirical basis for potential evolution of a theory of EIS quality change over time.

Finally, though not all EISs may be suitable for full online access due to commercial confidentiality, technical reasons and security concerns, studies of EIS quality review should be encouraged for the benefit of the industry and on-line availability of EIS is consistent with a drive towards greater

transparency and public accountability. Notwithstanding alleged criticism by some government regulators (Webb, 2011), we were pleased to find that since 2011, some Nigerian oil and gas project EISs have become available online.

5. Conclusion

In this paper we present the first ever study of EISs quality for both onshore and offshore oil and gas projects. As further novel elements, the study tests the hypothesis of improvement over time and critically reviews the degree to which project decommissioning impact is assessed within EISs.

The main study findings are: (a) project description and communication of results are main areas of strength; (b) of the 19 sampled EISs, almost half (47%) were scored as unsatisfactory; (c) environmental impact prediction and decommissioning were among the areas requiring enhanced attention; (d) Mann-Whitney tests suggest that there is no evidence that EISs for the latter period (2004-2008) were of a higher quality than those of the earlier period (1998-2004) – a finding that is contrary to common expectation.

Based on the findings of this study, we argue for National Environmental Policy in Nigeria to mandate a systematic, independent, periodic quality review of EISs every 3 to 5 years in the oil and gas industry, for the explicit purpose of improvement in good practice. Secondly, project decommissioning should henceforth be explicitly addressed in all future EISs in line with regulatory frameworks.

Beyond the case study sector and country, this study provides a robust demonstration that a systematic quality review reveals important areas needing adequate attention in EISs. Furthermore, this review could be used to target the dissemination of good practice/s and verify if EIA approval processes are sufficiently robust and adhered to.

Periodic systematic quality review of EISs, as described in this article, is part of wider responsible actions that would encourage transparency, stewardship or accountability and integrity – all of which are key principles of environmental sustainability. It would be useful to test our findings and hypothesis in other oil and gas producing countries, and in other extractive industries which carry a significant environmental risk.

Acknowledgements

The work reported in this paper formed part of a 3-year PhD Research Scholarship at The University of Birmingham funded by the Federal Government of Nigeria through the Petroleum Technology Development Fund (2007 to 2010). We wish to acknowledge the support received from the Federal Ministry of Environment and the Department of Petroleum Resources during the lead-author's working visits to Nigeria. Also, the lead author enjoyed the support of the Library and Information Services unit at the European Bank for Reconstruction and Development, London, UK. We appreciate the feedback on an earlier version of this paper presented at the first Society of Petroleum Engineers European HSSE conference in 2011, Vienna, Austria. Thanks to Anne Ankcorn of GEES, The University of Birmingham, for helping with Figure 1. We are grateful for the constructive criticisms from three anonymous peer-reviewers of this article.

References

- Agha, G.U, Irrechukwu, D.O. and Zagi, M.M. 2002. Environmental Impact Assessment and the Nigerian Oil Industry: A Review of Experiences and Learnings. Proceedings of the Society of Petroleum Engineers' International Conference on Health, Safety and Environment in Oil and Gas Exploration and Production; Kuala Lumpur, Malaysia 20-22 March.
- Agha, G.U, Irrechukwu, D.O. and Zagi, M.M. 2004. The Development of Environmental Guidelines and Standards for the Petroleum Industry in Nigeria: A Systematic Approach and Future Challenges. Proceedings of the Society of Petroleum Engineers' 7th International Conference on Health, Safety and Environment in Oil and Gas Exploration and Production; Calgary, Canada 29-31 March.
- Ambituuni, A., Amezaga, J. and Emeseh, E. 2014. Analysis of safety and environmental regulations for downstream petroleum industry operations in Nigeria. *Journal of Environmental Development* 9, 43–60.
- Ameyan, O. 2008. Environmental Impact Assessment: Insight into the environmental impact regulatory process and implementation for qualifying projects. Paper presented at a One-Day seminar on preparing business in Nigeria for environmental challenges, organized by Manufacturers Association of Nigeria (MAN), Ikeja, Lagos; 4 November.
- Anifowose, B.A., Lawler, D.M., van der Horst, D. and Chapman, L.C. 2011. Assessing the Quality of Oil and Gas project EIS. Proceedings of the Society of Petroleum Engineers European Health, Safety and Environmental (HSE) Conference in Oil and Gas Exploration and Production, 22-24 February, held in Vienna, Austria.
- Anifowose, B.A., Lawler, D.M., van der Horst, D. and Chapman, L. 2012a. Attacks on oil transport pipelines in Nigeria: a quantitative exploration and possible explanation of observed patterns. *Applied Geography*, (32) 636-651.
- Anifowose B.A., Lawler DM, van der Horst D and Chapman L 2012b Application of Flow Duration Analysis in River Oil Spill Management Proceedings of the Thirty-fifth AMOP Technical Seminar on Environmental Contamination and Response, Environment Canada, Ottawa, ON, pp. 813-835
- Anifowose, B., Lawler, DM., van der Horst, D., Chapman, L. 2014. Evaluating interdiction of oil pipelines at river crossings using Environmental Impact Assessments. *AREA*, 46(1) 4-17
- Anifowose, B.A. and Odubela, M.T. 2015. Methane emissions from oil and gas transport facilities – exploring innovative ways to mitigate environmental consequence. *Journal of Cleaner Production* 92, 121-133
- Backlund, A. 2009. Impact assessment in the European Commission – a system with multiple objectives. *Environmental science & policy* 12, 1077 – 1087.
- Barcelo, D. and Bennett, J. 2016. Human health and environmental risks of unconventional shale gas hydrofracking. *Science of The Total Environment* 544(15), 1139–1140.
- Barker, A. and Wood, C. 1999. An evaluation of EIA system performance in Eight EU Countries. *EIA Rev* 19, 387-404.
- Barker, A. and Jones, C. 2013. A critique of the performance of EIA within the offshore oil and gas sector. *Environ.ImpactAssess.*43,31–39.
- Bernard, D.P., Hunsaker, D.P., and Marmorek, D.R. 2001. Tools for improving predictive capabilities of environmental impact assessments: structured hypotheses, audits and monitoring. In: Hildebrand SG, Cannon JB, editors. Environmental analysis: the NEPA experience. Boca Raton (FL): Lewis, p. 547- 564.

- Bierkens, J. and Geerts, L. 2014. Environmental hazard and risk characterisation of petroleum substances: A guided “walking tour” of petroleum hydrocarbons. *Environment International* 66, 182–193
- Boothroyd, I., Almond, S., Qassim, S., Worrall, F. and Davies, R. 2016. Fugitive emissions of methane from abandoned, decommissioned oil and gas wells. *Science of The Total Environment* 547(15), 461–469.
- Boriani, E., Mariani, A., Baderna, D., Moretti, C., Lodi, M. and Benfenati, E. 2010. ERICA: A multiparametric toxicological risk index for the assessment of environmental healthiness. *Environment International* 36, 665–674
- Buckley, R. 1991. Auditing the precision and accuracy of environmental impact predictions in Australia. *Environmental Monitoring and Assessment* 18:1-23.
- Burgess, P. and Banks, E. 2013. Shepway District Council Folkestone Seafrost Environmental Statement Review. Empress House, Hampshire, pp.60.
- Butt, T., Gouda, H., Baloch, M., Paul, P., Javadi, A. and Alam, A. 2014. Literature review of baseline study for risk analysis — The landfill leachate case. *Environment International* 63, 149–162
- Canelas, L., Almansa, P., Merchan, M. and Cifuentes, P. 2005. Quality of environmental impact statements in Portugal and Spain. *EIA Rev.* 25, 217-225
- Cashmore, M. 2004. The role of science in environmental impact assessment: process and procedure versus purpose in the development of theory. *EIA Rev.* 24, 403–426.
- Cesar, A., Lia, L., Pereira, C., Santos, A., Cortez, F., Choueri, R., De Orte, M. and Rachid, B. 2014. Environmental assessment of dredged sediment in the major Latin American seaport (Santos, São Paulo — Brazil): An integrated approach. *Science of The Total Environment* 497–498(1) 679–687.
- Claggett, P., Okay, J. and Stehman, S. 2010. Monitoring Regional Riparian Forest Cover Change Using Stratified Sampling and Multiresolution Imagery. *Journal of the American Water Res. Ass.* 46(2), 334-343. DOI: 10.1111/j.1752-1688.2010.00424.x.
- Davoudi, M., Rahimpour, M., Jokar, S., Nikbakht, F. and Abbasfard, H. 2013. The major sources of gas flaring and air contamination in the natural gas processing plants: A case study. *Journal of Natural Gas Science and Engineering* 13, 7-19
- DPR, Department of Petroleum Resources 2002. Environmental Guidelines & Standards for the Petroleum Industry in Nigeria (EGASPIN), Lagos, Nigeria.
- EC, European Commission 2001. Guidance on EIA. EIS review.
- European Commission 2009, DG ENV. Study concerning the report on the application and effectiveness of the EIA Directive (Final Report). COWI, Denmark, p.198.
- Eilperin, 2010. J. U.S. exempted BP's Gulf of Mexico drilling from environmental impact study. Available online: <http://www.washingtonpost.com/wp-dyn/content/article/2010/05/04/AR2010050404118.html>. Accessed on: 3 June 2010.
- Ekins, P., Vanner, R. and Firebrace, J. 2006. Decommissioning of offshore oil and gas facilities: A comparative assessment of different scenarios. *Journal of Environmental Management* 79(4), 420–438
- Emeseh, E., 2012. Mainstreaming enforcement for the victims of environmental pollution: towards effective allocation of legislative competence under a Federal Constitution. *Environ. Law Rev.* 14 (2012), 184–198.

- Eneh, C.O., 2011. Managing Nigeria's environment: the unresolved issues. *J. Environ. Sci. Technol.* 4 (3), 250–263
- E&P Forum/UNEP 1997. Environmental management in Oil and Gas exploration and production. Joint E&P Forum/UNEP Technical Publications, Paris.
- Ernst, M.D. 2004. Permutation Methods: A Basis for Exact Inference. *Statistical Science* 19(4), 676–685.
- Evans, J., Wood, G. and Miller, A. 2006. The risk assessment–policy gap: An example from the UK contaminated land regime. *Environment International* 32, 1066–1071
- FEPA, Federal Environmental Protection Agency 1995. Sub-sectoral Guidelines for EIA in Nigeria's Oil and Gas Industry. Lagos, Nigeria.
- Field, A. 2009. Discovering statistics using SPSS (3rd ed.). Los Angelis: Sage Pub.
- Floater, G.J. 2002. A risk model for predicting the effects of lake-side development on wildfowl populations. *Journal of Environmental Management* 66, 307-316.
- Fuller, K. 1999. Quality and quality control in environmental impact assessment. In: Handbook of environmental impact assessment, vol. 2, Petts, J., Ed. Oxford: Blackwell. p.55-82.
- Gerring, J. 2004. What Is a Case Study and What Is It Good for? *American Political Science Review* 98(2), 341-354
- Gilbuena Jr., R., Kawamura, A., Medina, R., Amaguchi, H., Nakagawa, N. and Bui, D. 2013. Environmental impact assessment of structural flood mitigation measures by a rapid impact assessment matrix (RIAM) technique: A case study in Metro Manila, Philippines. *Science of The Total Environment* 456–457(1), 137–147.
- Giwa, S., Adama, O. and Akinyemi, O. 2014. Baseline black carbon emissions for gas flaring in the Niger Delta region of Nigeria. *Journal of Natural Gas Science and Engineering* 20, 373-379
- Glasson, J., Therivel, R., Weston, J., Wilson, E. and Frost, R. 1997. EIA - Learning from Experience: Changes in the Quality of Environmental Impact Statements for UK Planning Projects. *Journal of Environmental Planning and Mgt.* 40(4), 451- 464.
- Glasson J, Therivel R, Chadwick A. 2005. Introduction to environmental impact assessment. 3rd ed. London: Routledge.
- Gorard, S. 2006. Towards a judgement-based statistical analysis. *British Journal of Sociology of Education* 27(1), 67-80.
- Gray, I, and Edwards-Jones, G. 1999. Environmental impact assessment in the Scottish forest sector. *Forestry* 72(1),1–10.
- Gray, I. and Edwards-Jones, G. 2003. A review of environmental statements in the British forest sector. *Impact Assessment and Project Appraisal* 21(4), 303–312.
- Guilanpour, K and Sheate, W. 1997. A systematic review of Tanzanian environmental impact statements. *Project Appraisal* 12(3), 138-150.
- Gwimbi, P. and Godwell, G. 2016. Benchmarking the effectiveness of mitigation measures to the quality of environmental impact statements: lessons and insights from mines along the Great Dyke of Zimbabwe. *Environment Development and Sustainability* 18(2), 527-546.

- Heinma, K. and Poder, T. 2010. Effectiveness of Environmental Impact Assessment system in Estonia. *EIA Rev.* 30(4), 272–277
- HMSO, Her Majesty's Stationary Office 1996. Changes in the quality of environmental statements for planning projects. Research Report by the Impact Assessment Unit, School of Planning, Oxford Brookes University. London: HMSO.
- João, E. 2002. How scale affects environmental impact assessment. *EIA Rev.* 22, 289-310
- Johnson, E., Austin, B., Inlander, E., Gallipeau, C., Evans-White, M. and Entekin, S. 2015. Stream macroinvertebrate communities across a gradient of natural gas development in the Fayetteville Shale. *Science of The Total Environment* 530–531(15), 323–332.
- Kabir, S.M.Z., Momtaz, S. and Gladstone, W. 2010. The quality of Environmental Impact Statement (EIS) in Bangladesh. Proceedings of 30th Annual Meeting of the IAIA, ICC Geneva, Switzerland, April 6-11.
- Kabir, S. and Momtaz, S. 2014. Sectorial variation in the quality of environmental impact statements and factors influencing the quality. *Journal of Environmental Planning and Management* 57(11), 1595-1611
- Khadka, R.B. and Khanal, A.B. 2008. Environmental management plan (EMP) for Melamchi water supply project, Nepal. *Environ Monit Assess*, 146:225–234
- Klevas, V., Streimikiene, D. and Kleviene, A. 2009. Sustainability assessment of the energy projects implementation in regional scale. *Renewable and Sustainable Energy Reviews* 13, 155–166.
- Landis, M., Kamal, A., Kovalcik, K., Croghan, C., Norris, G. and Bergdale, A. 2016. The impact of commercially treated oil and gas produced water discharges on bromide concentrations and modeled brominated trihalomethane disinfection byproducts at two downstream municipal drinking water plants in the upper Allegheny River, Pennsylvania, USA. *Science of The Total Environment* 542-Part A(15), 505–520
- Lawler, D.M. 2003. Safeguarding water resources in the FSU in an era of transition: establishing WIA within the EIA process. *European Water* 3(4), 28-35.
- Lawler, D.M. 2004. Rivers and Environmental Impact Assessment issues. Abstract at the International Geographical Union conference; Glasgow, 15-20 August.
- Lawler, D.M. 2005. Towards the implementation of Strategic Environmental Assessment (SEA): learning from EIA for water resources, In: Schmidt, M., Joao, E. and Albrecht, E. (Eds) *Implementing Strategic Environmental Assessment*, Springer-Verlag, p.495-511.
- Lawler, D. and Milner, A. 2005. Sakhalin II Pipeline Project: River Crossings Report - Initial Review, Report to AEA Technologies, 21 October 2005, 31pp + Tables and Figs.
- Lawler, D.M. and Wilkes, M.A. 2015. Towards Improved Fluvial Sediment Impact Assessment (FSIA) approaches within Environmental Impact Assessments. *Croatian Geographical Bulletin (Hrvatski Geografski Glasnik)*, 77/2, pp.7–31.
- Lawrence, D.P. 1997. The Need for EIA Theory-Building. *EIA Rev.* 17, 79-107.
- Lee, N. and Brown, D. 1992. Quality control in environmental assessment. *Project Appraisal* 7(1), 41-45.
- Lee, N. and Colley, R. 1992. Reviewing the Quality of Environmental Statements. Occasional paper No. 24 (2nd Edition), EIA Center, University of Manchester, p.1-55.

Lee, N., Colley, R., Bonde, R. and Simpson, J. 1999. Reviewing the Quality of Environmental Statements and Environmental Appraisals. Occasional Paper No. 55, EIA Center, University of Manchester, p.1-72.

Leboeuf, Y., Burnett, P. and Bromm, S. 2010. International Panel on Experiences and Lessons Learned in a Multi-jurisdictional Context - Canada, the United States and Australia. Available online: <http://www.iaia.org/iaia10/documents/pdfs/International%20Panel%20on%20Experiences%20and%20Lessons%20Learned%20in%20a%20Multi-Jurisdictional%20Context%20-%20Canada,%20the%20United%20States%20and%20Australia.pdf?AspxAutoDetectCookieSupport=1>. Accessed on: 31 October 2014.

Ma'anit, A. 2011. Oil spill exposes Shell's ticking timebomb. Available online: <http://www.guardian.co.uk/commentisfree/2011/aug/17/oil-spill-shell-timebomb>. Accessed on 18 August 2011.

Marazza, D., Bandini, V. and Contin, A. 2010. Ranking environmental aspects in environmental management systems: A new method tested on local authorities. *Environment International* 36, 168–179

McMahon, N. 1996. Quality of environmental statements submitted in Northern Ireland in relation to the disposal of waste on land. *Project Appraisal* 11(2), 85-94.

Moustafa, K. 2016. Oil, Earth mass and gravitational force. *Science of The Total Environment* 548–549(1), 479–482.

Nasen, L.C., Noble, B.F. and Johnstone, J.F. 2011. Environmental effects of oil and gas lease sites in a grassland ecosystem. *Journal of Environmental Management* 92, 195-204.

Noble, B. and Storey, K. 2005. Towards increasing the utility of follow-up in Canadian Environmental Impact Assessment. *EIA Rev* 25, 163-180

Ogunba, O.A. 2004. EIA systems in Nigeria: evolution, current practice and shortcomings. *EIA Rev.* 24(6), 643-660.

Olokesusi, A.O. 1998. Legal and Institutional Framework of Environmental Impact Assessment in Nigeria: An Initial Assessment. *EIA Rev.*18, 159-174.

Palframan, L. 2010. The Integration of Environmental Impact Assessment and Environmental Management Systems: Experiences from the UK. Proceedings of 30th Annual Meeting of the IAIA, April 6-11, ICC Geneva, Switzerland.

Palframan, L. 2012. EIA–EMS Link from the Waste Management Sector. In: Perdicoulis, A., Durning, B. and Palframan, L. (eds). *Furthering Environmental Impact Assessment: Towards a Seamless Connection between EIA and EMS*. Edward Elgar: London.

Papadopoulou, M. and Antoniou, C. 2014. Environmental impact assessment methodological framework for liquefied natural gas terminal and transport network planning. *Energy Policy* 68, 306–319

Perdicoulis, A., Durning, B. and Palframan, L. 2012. *Furthering Environmental Impact Assessment: Towards a Seamless Connection between EIA and EMS*. Edward Elgar: London.

Pérez-López, P., Jeffryes, C., Agathos, S., Feijoo, G., Rorrer, G. and Moreira, M. 2016. Environmental life cycle optimization of essential terpene oils produced by the macroalga *Ochtodes secundiramea*. *Science of The Total Environment* 542-Part A(15) 292–305.

Peterson, K. 2010. Quality of environmental impact statements and variability of scrutiny by reviewers. *EIA Rev.* 30(3), 169–176

Petts, J. 1999. Introduction to environmental impact assessment in practice: fulfilled potential or wasted opportunity?. In: J. Petts, Editor, Handbook of environmental impact assessment vol 2, Oxford: Blackwell, p. 3–9.

PIB, Petroleum Industry Bill 2012. An Act to provide for the establishment of a legal, fiscal and regulatory framework for the petroleum industry in Nigeria and for other related matters. Abuja, Nigeria.

Raissiyan, B. and Pope, J. 2012. EIA–EMS Link from the Oil and Gas Industry. In: Perdicoulis, A., Durning, B. and Palframan, L. (eds). Furthering Environmental Impact Assessment: Towards a Seamless Connection between EIA and EMS. Edward Elgar: London.

Russo, T. 1999. Environmental Impact Assessment for Energy Projects. In: Handbook of environmental impact assessment, vol. 2, Petts, J., Ed. Oxford: Blackwell, p. 351–376.

Sadler, B. 1996. International Study of the Effectiveness of Environmental Assessment - Environmental Assessment in a Changing World: Evaluating Practice to Improve Performance. IAIA and Canadian Environmental Assessment Agency, Canada.

Sandham, L.A, Hoffmann, A.R. and Retief, F.P. 2008a. Reflections on the quality of mining EIA reports in South Africa. *Journal of the Southern African Institute of Mining & Metallurg* 108:701-716.

Sandham, L.A, Moloto, M.J. and Retief, F.P. 2008b. The Quality of Environmental Impact Reports for projects with the potential of affecting wetlands in South Africa. *Water SA* 34,155–163.

Sandham, L.A. and Pretorius, H.M. 2008. A review of EIA report quality in the North West province of South Africa. *EIA Rev.* 28, 229–240.

Sandham, L.A, Carroll, T.H. and Retief, F.P. 2010. The contribution of EIA to decision making for biological pest control in South Africa – The case of *Lantana camara*. *Biological Control*; doi:10.1016/j.biocontrol.2009.12.010

Scholten, J. 1995. Reviewing EISs/EIA Reports. Report of the EIA Process Strengthening Workshop, IAIA, Canadian Environmental Assessment Agency and Australian Environmental Protection Agency. Canberra April.

Schroeder, D.M. and Love, M.S. 2004. Ecological and political issues surrounding decommissioning of offshore oil facilities in the Southern California Bight. *Ocean & Coastal Management* 47, 21-48.

Seawright, J. and Gerring, J. 2008. Case Selection Techniques in Case Study Research. *Political Research Quarterly* 61(2), 294-308

Skierszkan, E., Irvine, G., Doyle, J., Kimpe, L. and Blais, J. 2013. Is there widespread metal contamination from in-situ bitumen extraction at Cold Lake, Alberta heavy oil field?. *Science of The Total Environment* 447, 337–344.

Steyn, P. 2006. Shell International, the Ogoni People and Environmental Injustice in the Niger Delta, Nigeria: The Challenge of Securing Environmental Justice in an Oil-based Economy. In: Hood Washington Sylvania, Rosier Paul C, Goodall Heather (ed.). *Echoes from the Poisoned Well: Global Memories of Environmental Injustice*, Lanham, MD: Lexington Books, pp. 371-388.

Steyn, P. 2009. Oil Exploration in Colonial Nigeria, c. 1903-58, *The Journal of Imperial and Commonwealth History* 37(2), 249-274

- Strutt, J., Sharp, J., Terry, E. and Miles, R. 2006. Capability maturity models for offshore organisational management. *Environment International* 32, 1094–1105
- Sullivan, G.M. and Feinn, R. 2012. Using effect size – or Why the P value is Not Enough. *Journal of Graduate Medical Education: Editorial*, 279-282.
- Talime, L.A. 2011. *A Critical Review Of The Quality Of EIA Reports In Lesotho*. An unpublished MSc Thesis, University of Free State, Bloemfontein, pp. 91.
- The National Commission 2011. *Deep Water: The Gulf Oil Disaster and the Future of Offshore Drilling*. Report to the US President on the BP Deepwater Horizon Oil Spill and Offshore Drilling.
- Torres, L., Yadav, O. and Khan, E. 2016. A review on risk assessment techniques for hydraulic fracturing water and produced water management implemented in onshore unconventional oil and gas production. *Science of The Total Environment* 539(1), 478–493
- Tzoumis, K. and Finegold, L. 2000. Looking at the quality of draft environmental impact statements over time in the United States: Have ratings improved? *Environ Impact Assess Rev* 20, 557-578.
- UNEP, United Nations Environmental Programme 2002. Topic 9 – Review of EIA quality. *EIA Training Resource Manual, Second (eds.)*.
- UNEP, United Nations Environmental Programme 2011. *Environmental Assessment of Ogoniland*. Nairobi, KENYA.
- Vedachalam, N., Srinivasalu, S., Ramesh, R., Aarthi, A., Ramadass, G. and Atmanand, M. 2015. Review and reliability modeling of maturing subsea hydrocarbon boosting systems. *Journal of Natural Gas Science and Engineering* 25, 284-296
- Wardekker, J.A., van der Sluijs, J.P., Janssen, P.H.M, Kloprogge, P. and Petersen, A.C. 2008. Uncertainty communication in environmental assessments: views from the Dutch science-policy interface. *Environmental science & policy* 627 – 641.
- Warnken, J. and Buckley, R. 1998. Scientific quality of tourism Environmental Impact Assessment. *Journal of Applied Ecology* 35,1-8.
- Werner, A., Vink, S., Watt, K. and Jagals, P. 2015. Environmental health impacts of unconventional natural gas development: A review of the current strength of evidence. *Science of The Total Environment* 505(1), 1127–1141
- Wawryk, A.S. 2002. *International Environmental Standards in the Oil Industry: Improving the Operations of Transnational Oil Companies in Emerging Economies*. Archived Journal Submission of the Centre for Energy, Petroleum and Mineral Law and Policy, Dundee, UK.
- Webb, T. 2011. Delta gangs are to blame, says Shell. Pollution has scarred its reputation but the oil giant is fighting back, says Tim Webb in the first of two reports. *The Times (London)*, July 11, p.37.
- Weston, J. 2004. EIA in a Risk Society. *Journal of Environmental Planning and Management*, 4(2), 313–325.
- World Bank 1999. *Environmental Assessment Sourcebook*. ENVLW, The World Bank, 1818 H St. NW, Washington, D.C.
- Zhang, J., Kørnøv, L and Christensen. P. 2013. Critical factors for EIA implementation: Literature review and research options. *Journal of Environmental Management* 114, 148-157.