

# Haemodynamic reactions to acute psychological stress and smoking status in a large community sample

Phillips, Anna; Der, G; Hunt, K; Carroll, Douglas

DOI:

[10.1016/j.ijpsycho.2009.04.005](https://doi.org/10.1016/j.ijpsycho.2009.04.005)

## Document Version

Peer reviewed version

## Citation for published version (Harvard):

Phillips, A, Der, G, Hunt, K & Carroll, D 2009, 'Haemodynamic reactions to acute psychological stress and smoking status in a large community sample', *International journal of psychophysiology : official journal of the International Organization of Psychophysiology*, vol. 73, no. 3, pp. 273-278.  
<https://doi.org/10.1016/j.ijpsycho.2009.04.005>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005.  
<http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

## Haemodynamic reactions to acute psychological stress and smoking status in a large community sample

Running Head: Disability and reactivity

Anna C. Phillips<sup>1</sup>

Geoff Der<sup>2</sup>

Michaela Benzeval<sup>2</sup>

Douglas Carroll<sup>1</sup>

<sup>1</sup>School of Sport and Exercise Sciences, University of Birmingham, Birmingham, England

<sup>2</sup>MRC Social and Public Health Sciences Unit, University of Glasgow, Glasgow, Scotland

Number of text pages:

Number of tables:

Number of figures:

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005.  
<http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

Address correspondence to: Anna C. Phillips, PhD, School of Sport and Exercise Sciences, University of Birmingham, Birmingham B15 2TT, England, UK. Tel: (00) 44 121 414 4398, Fax: (00) 44 121 414 4121. E-mail: [A.C.Phillips@bham.ac.uk](mailto:A.C.Phillips@bham.ac.uk)

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

### **Abstract**

Exaggerated haemodynamic reactions to acute psychological stress have been implicated in a number of adverse health outcomes. This study examined, in a large community sample, the cross-sectional associations between haemodynamic reactivity and self-reported smoking status. Blood pressure and heart rate were measured at rest and in response to a 3-minute arithmetic stress task. Participants were classified as current, ex-, or non-smokers by their response to a simple prompt. Smokers had significantly smaller SBP and DBP reactions to acute stress than ex- and non-smokers; current and ex-smokers had lower HR reactivity. These effects remained significant following adjustment for a host of variables likely to be associated with reactivity and/or smoking. Although the act of smoking acutely increases haemodynamic activity, the present findings contribute to a growing body of literature showing that smokers have blunted reactivity to mental stress. They also support the hypothesis that blunted reactivity may be characteristic of a range of dependencies. The present results also suggest that smoking status needs to be considered in the design and analysis of stress reactivity studies.

Key words: Acute psychological stress; Blood pressure; Cardiovascular reactivity; Disability; Heart rate;

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

## 1. Introduction

The present analyses of data from a large community sample of men and women allowed us to examine the issue of whether the magnitude of haemodynamic reactions to a standard mental stress task, as well as resting haemodynamic activity, was linked to physical disability status contemporary with reactivity measurement and the change in physical disability prospectively over five years. In addition, the richness of the database permitted adjustment for a number of potential confounders.

## 2. Materials and Methods

### 2.1 Participants

Data were collected as part of the West of Scotland Twenty-07 Study. Participants were all from Glasgow and surrounding areas in Scotland, and have been followed up at regular intervals since the baseline survey in 1987 (Ford et al., 1994). The study was designed to examine the processes that produce and maintain socio-demographic differences in health (Macintyre, 1987). The data reported here are from two age cohorts (born in the 1930s and 1950s) at the third follow-up carried out in 1995 and fourth follow-up conducted in 2000. Physical disability was measured at both follow-ups. At the third follow-up, cardiovascular reactions to an acute psychological stress task were measured (Carroll et al., 2000, Carroll et al., 2003). Reactivity data were available for 1038 participants, comprising two distinct age cohorts; 614 (59%) 44-year olds, and 424 (41%) 63-year olds. There were 563 (54%) women and 475 (46%) men, and 506 (49%) were from manual and 529 (51%) from non-manual occupation households. Household occupational group data were not available for three participants. The sample was almost entirely Caucasian, reflecting the West-of-Scotland population from which it was drawn. Overall mean age at this follow-up was 51.9 (SD = 9.39) years. The exact mean (SD) ages of the middle aged and eldest cohorts were 44.1 (0.85), and 63.1 (0.67) years, respectively. Mean (SD) body mass index, calculated from measured height and weight was 26.3 (4.24) kg/m<sup>2</sup>. Local ethics committee approval was obtained for each phase of data collection and all the participants provided informed consent at each follow-up.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

## 2.2 Apparatus and procedure

Participants were tested in a quiet room in their homes by trained nurses. As part of a structured interview, household occupational group was classified as manual or non-manual from the occupation of the head of household, using the Registrar General's Classification of Occupations (1980). Head of household was usually the man. Participants were classified as never, ex-, or current smokers by their response to a simple question: "do you ever smoke tobacco now? I am thinking of a pipe, cigars, and your own roll-ups as well as cigarettes you might buy." If the answer was no, they were then asked: "did you ever used to smoke any sort of tobacco?" In addition, current smokers were asked whether they regarded themselves as a very light occasional smoker, a light but regular smoker, a moderate smoker, quite a heavy smoker, or a very heavy smoker. It was also determined whether they were currently taking antihypertensive medication. Participants then undertook an acute psychological stress task: the paced auditory serial addition test (PASAT), which has been shown in numerous studies to reliably perturb the cardiovascular system (Ring et al., 2002, Winzer et al., 1999) and to demonstrate good test-retest reliability (Willemsen et al., 1998). Participants were presented, via audiotape, with a series of single digit numbers and had to add sequential number pairs while retaining the second of the pair in memory for addition to the next number presented, and so on throughout the series. Answers were given orally and, if participants faltered, they were prompted to begin again with the next number pair. The number of correct answers was recorded as a measure of performance. The first sequence of 30 numbers was presented at a rate of one every four seconds, and the second sequence of 30 at one every two seconds. The complete task took three minutes, two minutes for the slower and one minute for the faster sequence. A brief practice was given to ensure that participants understood the demands of the task. Only those who registered a score on the PASAT were included in the analyses. Out of a possible score of 60, the mean score was 40.9 (SD = 9.03).

SBP, DBP and HR were measured by an Omron (model 705CP) semi-automatic sphygmomanometer recommended by the European Society of Hypertension (O'Brien et al., 2001). Following the interview, (at least an hour), there was then a formal 5-minute period of relaxed sitting, at the end of which a resting baseline reading of SBP, DBP, and HR was taken.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

Task instructions were then given, followed by the brief practice. Two further SBP, DBP, and HR readings were taken during the task, the first initiated 20 seconds into the task (during the slower sequence of numbers), and the second initiated 110 seconds later (at the same point during the faster sequence). For all readings, the nurses ensured that the participant's elbow and forearm rested comfortably on a table at heart level. The two task readings were averaged and the resting baseline value subsequently subtracted from the resultant average task value to yield reactivity measures for SBP, DBP, and HR for each participant.

### 2.3 Statistical analyses

Repeated measures ANOVAs, comparing baseline and mean task value, were undertaken to confirm that the PASAT perturbed cardiovascular activity. To examine the socio-demographic patterning of smoking,  $\chi^2$  was applied. Prior to testing the association between smoking and reactivity, the relationship between smoking and baseline cardiovascular level was examined using ANOVA. Analyses of reactivity (the difference between the mean task and the baseline value) were by ANCOVA, with SBP, DBP, and HR reactivity as the dependent variable, smoking status as the independent variable; and baseline cardiovascular level was entered as a covariate. Where significant effects emerged, further ANCOVAs were undertaken that, in addition to baseline level, adjusted for socio-demographic characteristics (age, sex, occupational group) (Carroll et al., 2000) and other potential confounders previously found to be associated with reactivity in this sample (body mass index, antihypertensive medication status, PASAT performance score, depression) (Carroll et al., 2008, Carroll et al., 2007). Slight variations in degrees of freedom reflect occasional missing data for some variables. Significant variations among smoking status groups were followed up by post hoc pairwise comparisons using the Newman-Keuls method. For both ANOVAs and ANCOVAs, partial  $\eta^2$  is reported as a measure of effect size.

## 3. Results

### 3.1 Smoking status

Five hundred and ninety three (36%) of the sample were current smokers, 338 (21%) were ex-smokers, and 715 (43%) reported never smoking. Data were unavailable for one participant. The

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

socio-demographic breakdown is presented in Table 1. The numbers of both current and never smokers declined with age, whereas the numbers who had given up increased,  $\chi^2(4) = 80.99, p < .001$ . Participants from manual occupational households were more likely to smoke than those from non-manual households,  $\chi^2(2) = 51.03, p < .001$ . The sex difference in smoking status was not statistically significant,  $\chi^2(2) = 5.67, p = .06$ .

[Insert Table 1 about here]

### 3.2 Cardiovascular reactivity to the stress task

The PASAT provoked significant increases in SBP,  $F(1,1646) = 1562.32, p < .001, \eta^2_p = .487$ , DBP,  $F(1,1646) = 1066.62, p < .001, \eta^2_p = .393$ , and HR,  $F(1,1646) = 1132.96, p < .001, \eta^2_p = .408$ . We have described the socio-demographic effects on reactivity in previous reports (Carroll et al., 2000, Carroll et al., 2007) and they are summarized in Table 2. Briefly, HR reactivity declined with age, with the youngest cohort exhibiting higher reactivity than the middle cohort who, in turn, showed higher reactivity than the eldest cohort. HR reactivity was also greater in participants from non-manual occupational status households. The youngest cohort showed smaller SBP and DBP reactions than the other two cohorts and women were less SBP and DBP reactive than men.

[Insert Table 2 about here]

### 3.3 Smoking and baseline cardiovascular activity

Baseline cardiovascular levels according to smoking status are presented in Table 3. Ex smokers had higher SBP,  $F(2,1643) = 9.57, p < .001, \eta^2_p = .012$ , and DBP,  $F(2,1643) = 8.45, p < .001, \eta^2_p = .010$ , baseline levels than current smokers ( $p < .001$  in each case) and those who never smoked ( $p < .001$  in each case). There was also a main effect of smoking status on resting HR,  $F(2,1643) = 4.02, p = .02, \eta^2_p = .005$ ; current smokers had higher resting HR than never smokers ( $p = .005$ ). These associations were re-examined in ANCOVAs, adjusting for age cohort, sex, occupational status, body mass index, and whether or not participants were taking antihypertensive



Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

medication, which 142 (7%) were. The effects of smoking status on baseline SBP and DBP were abolished,  $F < 1$  in both cases. However, the effect on baseline HR levels survived such adjustment,  $F(2,1630) = 5.28, p = .005, \eta^2_p = .006$ .

### 3.4 Smoking and cardiovascular reactivity

The summary statistics are presented in Table 3. Adjusting for baseline levels, there were main effects of smoking group on cardiovascular reactivity, for SBP,  $F(2,1642) = 10.85, p < .001, \eta^2_p = .013$ , DBP,  $F(2,1642) = 7.02, p = .001, \eta^2_p = .008$ , and HR,  $F(2,1642) = 11.07, p < .001, \eta^2_p = .013$ . Smokers had significantly smaller SBP and DBP reactions than ex smokers ( $p < .001$ , and  $p = .003$ , respectively) and those who had never smoked ( $p \leq .001$  in each case). Current smokers also had smaller HR reactions than never ( $p < .001$ ) but not ex smokers.

The analyses were repeated adjusting for, in addition to baseline level, age cohort, sex, occupational status, body mass index, whether or not participants were taking antihypertensive medication, and performance score on the PASAT. The latter is associated with cardiovascular reactivity in this sample (Carroll et al., 2008) and smokers performed more poorly than participants who had never smoked,  $F(2,1643) = 4.21, p = .02, \eta^2_p = .006$ ; the mean (SD) performance scores were 43.0 (9.26), 43.6 (9.29), and 44.5 (9.09), for current, ex, and never smokers, respectively. The effects reported above were still significant following adjustment:  $F(2,1628) = 10.27, p < .001, \eta^2_p = .012$ ,  $F(2,1628) = 5.32, p = .005, \eta^2_p = .006$ , and  $F(2,1628) = 8.98, p < .001, \eta^2_p = .011$ , for SBP, DBP, and HR reactivity respectively. We have already reported that reactivity was negatively associated with depressive symptomatology in this sample (Carroll et al., 2007). Symptoms of depression were measured at the third follow up using the Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983), a well-recognised assessment tool. Depression scores were also higher in smokers than participants who had never smoked,  $F(2,1605) = 9.55, p < .001, \eta^2_p = .012$ ; the mean (SD) scores were 4.0 (2.95), 3.7 (2.87), and 3.3 (2.75), for current, ex, and never smokers, respectively. Accordingly, in another series of ANCOVAs, we adjusted additionally for depression score. Although the smoking status effects were attenuated, they remained statistically significant:  $F(2,1590) = 6.93, p = .001, \eta^2_p = .009$ ,  $F(2,1590) = 4.89, p = .008, \eta^2_p = .006$ , and  $F(2,1590) = 3.87, p = .02, \eta^2_p = .005$ , for SBP, DBP,

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

and HR reactivity, respectively. The models with all these covariates accounted for 13%, 12%, and 16% of the variation in SBP, DBP, and HR reactivity, respectively.

[Insert Table 3 about here]

### 3.5 Smoking regularity and cardiovascular reactivity

The cardiovascular reactivity of smokers who described themselves as very light occasional smokers (N = 87) was compared to that of the rest of the smokers (N = 505). There were no differences between groups in blood pressure reactivity. However, the very light occasional smokers (mean = 9.8, SD = 9.58 bpm) tended to exhibit greater HR reactivity than the rest of the smokers (mean = 6.5, SD = 9.15 bpm),  $F(1,589) = 3.33$ ,  $p = .07$ ,  $\eta^2_p = .006$ .

## 4. Discussion

Whereas previous research indicates that the excitatory cardiovascular effects of smoking at the time of exposure add to the haemodynamic impact of acute psychological stress (Davis and Matthews, 1990, MacDougall et al., 1988, Ray et al., 1986), the consequences of being a habitual smoker for reactivity are less clear. Nevertheless, two larger scale studies suggest that, if anything, smokers conventionally exhibit smaller blood pressure and HR reactions to stress than non-smokers (Roy et al., 1994, Sheffield et al., 1997b). The present results from a large community sample provide strong confirmatory evidence. Smokers displayed significantly smaller SBP and DBP reactions to acute stress than ex and non-smokers. Further, current and ex-smokers had lower HR reactivity. The novel inclusion of the ex-smoker category in the present study allows us to determine whether giving up smoking is associated with a similar magnitude of reactivity to that typical of non-smokers; this would certainly appear to be the case for blood pressure, although not HR, reactivity. These associations between smoking status and cardiovascular reactivity remained significant following adjustment for a host of variables likely to be associated with reactivity and/or smoking: baseline cardiovascular level, age cohort, sex, occupational status, body mass index, performance score on the PASAT, whether or not participants were taking antihypertensive medication, and depressive symptoms score. In

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

addition, very light occasional smokers tended to have larger HR reactions to stress than more frequent smokers. Only one previous study, that we know of, considered regularity of smoking; no differences in cardiovascular reactivity were reported between moderate (<15 cigarettes a day) and heavy smokers (Roy et al., 1994). However, it is likely that the present self-categorisation of very light occasional smokers contained individuals who were much closer to non-smokers than the moderate smokers in the previous study.

The direction of the present association finds common ground with the results of other studies reporting generally blunted physiological adjustments in smokers to a range of challenges. For example, in a study using positron emission tomography to measure myocardial perfusion, smokers were reported to show an impaired increase in perfusion to the cold pressor test relative to non-smokers (Meeder et al., 1996). Further, smokers were found to have a smaller flow velocity response than non-smokers in the posterior cerebral arteries, measured by functional transcranial Doppler, to a visual stimulus challenge (Olah et al., 2008). In addition, smokers have been shown to have reduced density and down-regulated function of beta-adrenergic receptors, which might also contribute to reduced cardiovascular reactivity (Laustiola et al., 1988). These effects have been found to be reversible with smoking cessation (Laustiola et al., 1991). Thus, it would appear that habitual smoking may be characterised by depressed rather than exaggerated sympathetic nervous system reactivity. It is also possible that the associations observed in the present study, to an extent, reflect variations in parasympathetic withdrawal, as smokers have been reported to show a blunted response of the low frequency component of the heart rate variability spectrum to orthostatic stress (Lucini et al., 1996).

The direction of the present associations are also in line with a growing body of evidence that smokers exhibit blunted hypothalamic-pituitary-adrenocortical (HPA) axis stress reactivity. Although the act of smoking and acute doses of nicotine activate the HPA axis (Kirschbaum et al., 1992, Pomerleau et al., 1983), habitual smokers have been found to show diminished salivary and plasma cortisol reactions to psychological stress (al'Absi et al., 2003, Kirschbaum et al., 1993, Kirschbaum et al., 1994, Rohleder and Kirschbaum, 2006). In addition, this cortisol hypo-responsiveness has been found to predict relapse among smokers who have recently quit

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

smoking; blunted blood pressure reactivity was also associated with relapse (al'Absi et al., 2005, al'Absi, 2006). Given that haemodynamic and cortisol stress reactivity are strongly correlated (e.g. (Cacioppo, 1994), it is perhaps unsurprising that attenuated reactivity in one system is paralleled by the diminished reactions of the other and that blunted reactivity in the two systems may be similarly implicated in smoking relapse. Indeed, it has been argued recently that blunted reactivity, also found in alcoholics and drug abusers, may be a general marker for risk of dependency and addiction (Lovallo, 2006).

Ex-smokers registered higher resting blood pressure than smokers and non-smokers. In previous epidemiological comparisons of smokers and non-smokers, the former are generally reported to have lower blood pressure (Goldbourt et al., 1997, Green et al., 1986, Sheffield et al., 1997b). However, in the present study, group variations in resting blood pressure were abolished following statistical adjustment for a number of covariates. Accordingly, it is possible that findings of relative low resting blood pressure among smokers are artefactual. In contrast, smokers had higher resting HR levels than the other two groups. This result is also not without precedent (Roy et al., 1994, Sheffield et al., 1997b). Further, the effect withstood adjustment for possible confounders.

The present study is not without limitations. First, we measured only blood pressure and HR. It could have proved instructive to have the sort of comprehensive assessment of haemodynamics afforded by impedance cardiography. However, the decision to test participants in their own homes and the size of the sample precluded the use of impedance cardiography. Second, determining causality is impossible from cross-sectional observational data and confounding is always a potential problem (Christenfeld et al., 2004). However, we did adjust statistically for a broad range of possible confounders. Nevertheless, residual confounding as a consequence of poorly measured or un-measured variables cannot be wholly discounted. For example, one very parsimonious explanation for the direction of relationship observed in the present study is that smokers, as a result of temporary abstinence during stress testing (Roy et al., 1994) found it more difficult to concentrate and engaged less with the stress task. This would be reflected in a poorer stress task performance score and current smokers (mean = 43.0, SD = 9.26)

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

did not perform as well as participants who had never smoked (mean = 44.5, SD = 9.09) with ex-smokers (mean = 43.6, SD = 9.29) occupying the mid position,  $F(2,1643) = 4.21, p = .02, \eta^2_p = .005$ . However, the group differences in performance are small and the associations between smoking status and haemodynamic reactivity remained significant after adjustment for performance score. Third, and relatedly, we had to rely on performance score as our measure of task engagement. Although this seems reasonable, in retrospect it might have proved useful to have included self-report measures of stress task impact. Fourth, it is also possible that nicotine deprivation *per se* as a result of temporary withdrawal may have contributed to blunted reactivity. The present study would undoubtedly have benefited from including a measure of smoking recency. However, blunted cardiac output and heart rate reactivity to acute stressors in female smokers have been observed regardless of whether they were wearing a nicotine replacement patch or placebo patch (Girdler et al., 1997). In addition, cardiovascular reactivity has been compared among non-smokers, smokers who abstained from smoking, and smokers who continued to smoke at their usual rate; smokers, irrespective of their assigned condition, showed attenuated blood pressure reactions to psychological stress (al'Absi et al., 2003). Thus, we think it unlikely that any short-term abstinence in the present study would have affected reactivity dramatically. Finally, our main measure of smoking behaviour was crude to say the least. For example, it has been argued that in terms of the total exposure to the toxic smoke components of tobacco, the way in which cigarettes are smoked may be as important as whether people smoke (Jarvis and Russell, 1980). In the present study no account was taken of the extent of inhalation. However, subjective measures of inhalation have proved unsatisfactory (Stepney, 1982) and most previous studies, including those on smoking and haemodynamic reactivity, have relied on simple taxonomies of the sort used here. Although smoking is sometimes underestimated in self-reports, particularly among ex-smokers (Lewis et al., 2003), there is evidence that smokers' reports can be reasonably reliable and agree with objective measures such as CO exhalation (Mak et al., 2005). Nevertheless, it would have strengthened the study had more detailed information regarding duration and intensity of smoking behaviour been available, as well as biochemical measures to verify smoking behaviour.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

In conclusion, compared to non-smokers and/or ex-smokers, current smokers exhibited blunted haemodynamic reactions to acute psychological stress. The effects, although attenuated, were still significant following adjustment for a wide range of candidate confounders. These findings may have both theoretical and methodological ramifications. First, it would appear that smokers' increased risk for cardiovascular disease is not a function of increased reactivity. Second, the direction of the associations observed is consistent with the notion that blunted stress reactivity in smokers may be common to both the sympathetic nervous system and the HPA axis. Third, it resonates with the idea that blunted reactivity may be characteristic of a range of dependencies. Further studies, however, will be required to confirm this sort of speculation. Fourth, smoking status is rarely if ever mentioned as a variable that needs to be considered when designing stress reactivity studies or in analysing the data from them. As others have noted (Roy et al., 1994), it may be necessary to adjust for smoking status or to treat smokers and non-smokers separately when analysing the effects of other psychosocial or behavioural variables on haemodynamic reactions to stress.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

## References

1980 Classification of Occupations. London, HMSO.

Al'absi, M. 2006 Hypothalamic-pituitary-adrenocortical responses to psychological stress and risk for smoking relapse. *Int J Psychophysiol*, 59, 218-27.

Al'absi, M., Hatsukami, D. & Davis, G. L. 2005 Attenuated adrenocorticotrophic responses to psychological stress are associated with early smoking relapse. *Psychopharmacology (Berl)*, 181, 107-17.

Al'absi, M., Wittmers, L. E., Erickson, J., Hatsukami, D. & Crouse, B. 2003 Attenuated adrenocortical and blood pressure responses to psychological stress in ad libitum and abstinent smokers. *Pharmacol Biochem Behav*, 74, 401-10.

Allen, M. T., Matthews, K. A. & Sherman, F. S. 1997 Cardiovascular reactivity to stress and left ventricular mass in youth. *Hypertension*, 30, 782-7.

Barnett, P. A., Spence, J. D., Manuck, S. B. & Jennings, J. R. 1997 Psychological stress and the progression of carotid artery disease. *J Hypertens*, 15, 49-55.

Cacioppo, J. T. 1994 Social Neuroscience: Autonomic, neuroendocrine, and immune responses to stress. *Psychophysiology*, 31, 113-128.

Carroll, D., Harrison, L. K., Johnston, D. W., Ford, G., Hunt, K., Der, G. & West, P. 2000 Cardiovascular reactions to psychological stress: the influence of demographic variables. *J Epidemiol Community Health*, 54, 876-877.

Carroll, D., Phillips, A. C. & Der, G. 2008 Body mass index, abdominal adiposity, obesity and cardiovascular reactions to psychological stress in a large community sample. *Psychosom Med*, 70, 653-660.

Carroll, D., Phillips, A. C., Hunt, K. & Der, G. 2007 Symptoms of depression and cardiovascular reactions to acute psychological stress: evidence from a population study. *Biological Psychology*, 75, 68-74.

Carroll, D., Ring, C., Hunt, K., Ford, G. & Macintyre, S. 2003 Blood pressure reactions to stress and the prediction of future blood pressure: effects of sex, age, and socioeconomic position. *Psychosom Med*, 65, 1058-1064.

- Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>
- Christenfeld, N. J., Sloan, R. P., Carroll, D. & Greenland, S. 2004 Risk factors, confounding, and the illusion of statistical control. *Psychosom Med*, 66, 868-75.
- Davis, M. C. & Matthews, K. A. 1990 Cigarette smoking and oral contraceptive use influence women's lipid, lipoprotein, and cardiovascular responses during stress. *Health Psychol*, 9, 717-36.
- Dembroski, T. M., Macdougall, J. M., Cardozo, S. R., Ireland, S. K. & Krug-Fite, J. 1985 Selective cardiovascular effects of stress and cigarette smoking in young women. *Health Psychol*, 4, 153-67.
- Ford, G., Ecob, R., Hunt, K., Macintyre, S. & West, P. 1994 Patterns of class inequality in health through the lifespan: class gradients at 15, 35 and 55 years in the west of Scotland. *Soc Sci Med*, 39, 1037-50.
- Girdler, S. S., Jamner, L. D., Jarvik, M., Soles, J. R. & Shapiro, D. 1997 Smoking status and nicotine administration differentially modify hemodynamic stress reactivity in men and women. *Psychosom Med*, 59, 294-306.
- Goldbourt, U., Yaari, S. & Medalie, J. H. 1997 Isolated low HDL cholesterol as a risk factor for coronary heart disease mortality. A 21-year follow-up of 8000 men. *Arterioscler Thromb Vasc Biol*, 17, 107-13.
- Green, M. S., Harari, G. & Schwartz, K. 1991 Cigarette smoking related to ambulatory blood pressure and heart rate. *Am Heart J*, 121, 1569-70.
- Green, M. S., Jucha, E. & Luz, Y. 1986 Blood pressure in smokers and nonsmokers: epidemiologic findings. *Am Heart J*, 111, 932-40.
- Handa, K., Tanaka, H., Shindo, M., Kono, S., Sasaki, J. & Arakawa, K. 1990 Relationship of cigarette smoking to blood pressure and serum lipids. *Atherosclerosis*, 84, 189-93.
- Hasenfratz, M. & Battig, K. 1992 Action profiles of smoking and caffeine: Stroop effect, EEG, and peripheral physiology. *Pharmacol Biochem Behav*, 42, 155-61.
- Heishman, S. J., Snyder, F. R. & Henningfield, J. E. 1993 Performance, subjective, and physiological effects of nicotine in non-smokers. *Drug Alcohol Depend*, 34, 11-8.
- Hori, T., Hayashi, M., Oka, M., Agari, I., Kawabe, K. & Takagi, M. 1994 Re-examination of arousing and de-arousing effects of cigarette smoking. *Percept Mot Skills*, 78, 787-800.



- Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>
- James, J. E. & Richardson, M. 1991 Pressor effects of caffeine and cigarette smoking. *Br J Clin Psychol*, 30 ( Pt 3), 276-8.
- Jarvis, M. J. & Russell, M. A. 1980 Comment on the Hunter Committee's second report. *Br Med J*, 280, 994-5.
- Kamarck, T. W., Everson, S. A., Kaplan, G. A., Manuck, S. B., Jennings, J. R., Salonen, R. & Salonen, J. T. 1997 Exaggerated blood pressure responses during mental stress are associated with enhanced carotid atherosclerosis in middle-aged Finnish men: findings from the Kuopio Ischemic Heart Disease Study. *Circulation*, 96, 3842-8.
- Kirschbaum, C., Scherer, G. & Strasburger, C. J. 1994 Pituitary and adrenal hormone responses to pharmacological, physical, and psychological stimulation in habitual smokers and nonsmokers. *Clin Investig*, 72, 804-10.
- Kirschbaum, C., Strasburger, C. J. & Langkrar, J. 1993 Attenuated cortisol response to psychological stress but not to CRH or ergometry in young habitual smokers. *Pharmacology and Biochemistry of Behavior*, 44, 527-31.
- Kirschbaum, C., Wust, S. & Strasburger, C. J. 1992 'Normal' cigarette smoking increases free cortisol in habitual smokers. *Life Sci*, 50, 435-42.
- Laustiola, K. E., Kotamaki, M., Lassila, R., Kallioniemi, O. P. & Manninen, V. 1991 Cigarette smoking alters sympathoadrenal regulation by decreasing the density of beta 2-adrenoceptors. A study of monitored smoking cessation. *J Cardiovasc Pharmacol*, 17, 923-8.
- Laustiola, K. E., Lassila, R., Kaprio, J. & Koskenvuo, M. 1988 Decreased beta-adrenergic receptor density and catecholamine response in male cigarette smokers. A study of monozygotic twin pairs discordant for smoking. *Circulation*, 78, 1234-40.
- Lewis, S. J., Cherry, N. M., Mc, L. N. R., Barber, P. V., Wilde, K. & Povey, A. C. 2003 Cotinine levels and self-reported smoking status in patients attending a bronchoscopy clinic. *Biomarkers*, 8, 218-28.
- Lovallo, W. R. 2006 Cortisol secretion patterns in addiction and addiction risk. *Int J Psychophysiol*, 59, 195-202.
- Lovallo, W. R. & Gerin, W. 2003 Psychophysiological reactivity: mechanisms and pathways to cardiovascular disease. *Psychosom Med*, 65, 36-45.

- Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>
- Lucini, D., Bertocchi, F., Malliani, A. & Pagani, M. 1996 A controlled study of the autonomic changes produced by habitual cigarette smoking in healthy subjects. *Cardiovasc Res*, 31, 633-9.
- Lynch, J. W., Everson, S. A., Kaplan, G. A., Salonen, R. & Salonen, J. T. 1998 Does low socioeconomic status potentiate the effects of heightened cardiovascular responses to stress on the progression of carotid atherosclerosis? *Am J Public Health*, 88, 389-94.
- Macdougall, J. M., Musante, L., Castillo, S. & Acevedo, M. C. 1988 Smoking, caffeine, and stress: effects on blood pressure and heart rate in male and female college students. *Health Psychol*, 7, 461-78.
- Macintyre, S. 1987 West of Scotland Twenty-07: Health in the community. The survey's background and rationale (Working Paper No. 7). Glasgow, MRC Medical Sociology Unit.
- Mak, Y. W., Loke, A. Y., Lam, T. H. & Abdullah, A. S. 2005 Validity of self-reports and reliability of spousal proxy reports on the smoking behavior of Chinese parents with young children. *Addict Behav*, 30, 841-5.
- Mann, S. J., James, G. D., Wang, R. S. & Pickering, T. G. 1991 Elevation of ambulatory systolic blood pressure in hypertensive smokers. A case-control study. *Jama*, 265, 2226-8.
- Markovitz, J. H., Raczynski, J. M., Wallace, D., Chettur, V. & Chesney, M. A. 1998 Cardiovascular reactivity to video game predicts subsequent blood pressure increases in young men: The CARDIA study. *Psychosom Med*, 60, 186-91.
- Meeder, J. G., Blanksma, P. K., Van Der Wall, E. E., Anthonio, R. L., Willemsen, A. T., Pruijm, J., Vaalburg, W. & Lie, K. I. 1996 Long-term cigarette smoking is associated with increased myocardial perfusion heterogeneity assessed by positron emission tomography. *Eur J Nucl Med*, 23, 1442-7.
- O'brien, E., Waeber, B., Parati, G., Staessen, J. & Myers, M. G. 2001 Blood pressure measuring devices: recommendations of the European Society of Hypertension. *BMJ*, 322, 531-6.
- Olah, L., Raiter, Y., Candale, C., Molnar, S., Rosengarten, B., Bornstein, N. M. & Csiba, L. 2008 Visually evoked cerebral vasomotor response in smoking and nonsmoking young adults, investigated by functional transcranial Doppler. *Nicotine Tob Res*, 10, 353-8.

- Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>
- Parati, G., Trauzzi, S., Ravogli, A., Casadei, R., Omboni, S. & Mancia, G. 1991 Methodological problems in evaluation of cardiovascular effects of stress in humans. *Hypertension*, 17, III50-5.
- Pauli, P., Rau, H., Zhuang, P., Brody, S. & Birbaumer, N. 1993 Effects of smoking on thermal pain threshold in deprived and minimally-deprived habitual smokers. *Psychopharmacology (Berl)*, 111, 472-6.
- Perkins, K. A., Grobe, J. E., Fonte, C. & Breus, M. 1992 "Paradoxical" effects of smoking on subjective stress versus cardiovascular arousal in males and females. *Pharmacol Biochem Behav*, 42, 301-11.
- Perkins, K. A., Sexton, J. E., Reynolds, W. A., Grobe, J. E., Fonte, C. & Stiller, R. L. 1994 Comparison of acute subjective and heart rate effects of nicotine intake via tobacco smoking versus nasal spray. *Pharmacol Biochem Behav*, 47, 295-9.
- Pomerleau, O. F., Fertig, J. B., Seyler, L. E. & Jaffe, J. 1983 Neuroendocrine reactivity to nicotine in smokers. *Psychopharmacology (Berl)*, 81, 61-7.
- Ray, R. L., Nellis, M. J., Brady, J. V. & Foltin, R. W. 1986 Nicotine and caffeine effects on the task-elicited blood pressure response. *Addict Behav*, 11, 31-6.
- Ring, C., Burns, V. E. & Carroll, D. 2002 Shifting hemodynamics of blood pressure control during prolonged mental stress. *Psychophysiology*, 39, 585-90.
- Rohleder, N. & Kirschbaum, C. 2006 The hypothalamic-pituitary-adrenal (HPA) axis in habitual smokers. *Int J Psychophysiol*, 59, 236-43.
- Roy, M. P., Steptoe, A. & Kirschbaum, C. 1994 Association between smoking status and cardiovascular and cortisol stress responsivity in healthy young men. *Int J Behav Med*, 1, 264-83.
- Schwartz, A. R., Gerin, W., Davidson, K. W., Pickering, T. G., Brosschot, J. F., Thayer, J. F., Christenfeld, N. & Linden, W. 2003 Toward a causal model of cardiovascular responses to stress and the development of cardiovascular disease. *Psychosom Med*, 65, 22-35.
- Sheffield, D., Smith, G. D., Carroll, D., Shipley, M. J. & Marmot, M. G. 1997a The effects of blood pressure resting level and lability on cardiovascular reactions to laboratory stress. *Int J Psychophysiol*, 27, 79-86.

- Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>
- Sheffield, D., Smith, G. D., Carroll, D., Shipley, M. J. & Marmot, M. G. 1997b The effects of recent food, alcohol, and tobacco intake and the temporal scheduling of testing on cardiovascular activity at rest and during psychological stress. *Psychophysiology*, 34, 204-12.
- Stepney, R. 1982 Are smokers' self-reports of inhalation a useful measure of smoke exposure. *J Epidemiol Community Health*, 36, 109-12.
- Straneva, P., Hinderliter, A., Wells, E., Lenahan, H. & Girdler, S. 2000 Smoking, oral contraceptives, and cardiovascular reactivity to stress. *Obstet Gynecol*, 95, 78-83.
- Tersman, Z., Collins, A. & Eneroth, P. 1991 Cardiovascular responses to psychological and physiological stressors during the menstrual cycle. *Psychosom Med*, 53, 185-97.
- Treiber, F. A., Kamarck, T., Schneiderman, N., Sheffield, D., Kapuku, G. & Taylor, T. 2003 Cardiovascular reactivity and development of preclinical and clinical disease states. *Psychosom Med*, 65, 46-62.
- Willemsen, G., Ring, C., Carroll, D., Evans, P., Clow, A. & Hucklebridge, F. 1998 Secretory immunoglobulin A and cardiovascular reactions to mental arithmetic and cold pressor. *Psychophysiology*, 35, 252-9.
- Winzer, A., Ring, C., Carroll, D., Willemsen, G., Drayson, M. & Kendall, M. 1999 Secretory immunoglobulin A and cardiovascular reactions to mental arithmetic, cold pressor, and exercise: effects of beta-adrenergic blockade. *Psychophysiology*, 36, 591-601.
- Zigmond, A. S. & Snaith, R. P. 1983 The hospital anxiety and depression scale. *Acta Psychiatr Scand*, 67, 361-70.

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

Table 1. Smoking(%) by age cohort, sex, and household occupational status

		Current	Ex-	Never
<i>Age Cohort:</i>	Youngest	239 (40)	58 (10)	294 (50)
	Middle	218 (35)	140 (22)	266 (43)
	Eldest	136 (32)	140 (32)	155 (36)
<i>Sex:</i>	Male	287 (38)	165 (22)	305 (40)
	Female	306 (34)	173 (20)	410 (46)
<i>Occupational Group:</i>	Manual	343 (45)	155 (20)	273 (35)
	Non-manual	248 (29)	181 (21)	441 (50)

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

Table 2. Mean (SD) SBP, DBP, and HR during baseline and the stress task by age cohort, sex, and occupational status

<i>Group</i>		<i>SBP</i>		<i>DBP</i>		<i>HR</i>	
		<i>Baseline</i>	<i>Task</i>	<i>Baseline</i>	<i>Task</i>	<i>Baseline</i>	<i>Task</i>
<i>Age Cohort:</i>	Youngest	120.0 (15.07)	130.1 (15.93)	73.4 (10.08)	80.2 (10.27)	67.5 (11.00)	77.5 (12.74)
	Middle	127.1 (18.08)	139.4 (18.75)	80.6 (11.13)	87.7 (11.30)	66.7 (11.17)	74.5 (12.17)
	Eldest	144.4 (21.68)	156.7 (22.80)	83.8 (11.17)	90.8 (13.23)	65.7 (9.92)	71.7 (11.15)
<i>Sex:</i>	Male	134.7 (18.25)	147.5 (12.22)	81.2 (11.18)	88.4 (11.73)	64.7 (10.43)	73.4 (12.19)
	Female	124.3 (21.07)	134.7 (21.06)	76.8 (11.56)	85.6 (12.34)	68.4 (10.84)	76.0 (12.32)
<i>Occupational Group:</i>	Manual	130.5 (21.44)	141.7 (22.17)	79.3 (11.93)	85.9 (12.56)	67.0 (11.26)	74.0 (12.19)
	Non-manual	127.8 (19.58)	139.6 (21.16)	78.4 (11.29)	85.7 (12.06)	66.5 (10.40)	75.6 (12.39)

Table 3. Mean (SD) SBP, DBP, and HR baseline and reactivity by smoking status

		Current	Ex	Never
<i>SBP:</i>	Baseline	128.2 (20.30)	133.4 (21.41)	127.8 (19.98)
	Reactivity	9.8 (11.41)	12.4 (11.57)	12.4 (11.97)
<i>DBP:</i>	Baseline	78.3 (11.70)	81.2 (11.74)	78.2 (11.36)
	Reactivity	6.0 (8.09)	7.1 (8.27)	7.6 (9.15)
<i>HR:</i>	Baseline	67.7 (11.18)	66.5 (10.93)	66.0 (10.38)
	Reactivity	6.9 (9.28)	7.1 (8.75)	9.5 (10.43)

Phillips, A.C., Der, G., Hunt, K., & Carroll, D. (2009). Haemodynamic reactions to acute psychological stress and smoking status in a large community sample. *International Journal of Psychophysiology*, 73, 273-278. DOI: 10.1016/j.ijpsycho.2009.04.005. <http://dx.doi.org/10.1016/j.ijpsycho.2009.04.005>

