

## Glucocorticoids modulate human brown adipose tissue thermogenesis in vivo

Scotney, Hannah; Symonds, Michael E.; Law, James; Budge, Helen ; Sharkey, Don; Manolopoulos, Konstantinos

DOI:

[10.1016/j.metabol.2017.01.024](https://doi.org/10.1016/j.metabol.2017.01.024)

License:

Creative Commons: Attribution (CC BY)

### Document Version

Publisher's PDF, also known as Version of record

### Citation for published version (Harvard):

Scotney, H, Symonds, ME, Law, J, Budge, H, Sharkey, D & Manolopoulos, K 2017, 'Glucocorticoids modulate human brown adipose tissue thermogenesis in vivo', *Metabolism*, vol. 70, pp. 125-132.  
<https://doi.org/10.1016/j.metabol.2017.01.024>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

# Metabolism

[www.metabolismjournal.com](http://www.metabolismjournal.com)

## Glucocorticoids modulate human brown adipose tissue thermogenesis *in vivo*



Hannah Scotney<sup>a</sup>, Michael E. Symonds<sup>a,b,\*</sup>, James Law<sup>a</sup>, Helen Budge<sup>a</sup>,  
Don Sharkey<sup>a</sup>, Konstantinos N. Manolopoulos<sup>c,d,\*\*</sup>

<sup>a</sup> Division of Child Health, Obstetrics & Gynaecology, School of Medicine, University of Nottingham, Nottingham, NG7 2UH, UK

<sup>b</sup> Nottingham Digestive Diseases Biomedical Research Unit, Queens Medical Centre Campus, Nottingham, NG7 2UH

<sup>c</sup> Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, B15 2TT, UK

<sup>d</sup> Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, B15 2TT, UK

### ARTICLE INFO

#### Article history:

Received 3 October 2016

Accepted 14 January 2017

#### Keywords:

Brown adipose tissue

Glucocorticoids

Humans

Infrared thermography

Beta adrenoceptor

### ABSTRACT

**Introduction.** Brown adipose tissue (BAT) is a thermogenic organ with substantial metabolic capacity and has important roles in the maintenance of body weight and metabolism. Regulation of BAT is primarily mediated through the  $\beta$ -adrenoceptor ( $\beta$ -AR) pathway. The *in vivo* endocrine regulation of this pathway in humans is unknown. The objective of our study was to assess the *in vivo* BAT temperature responses to acute glucocorticoid administration.

**Methods.** We studied 8 healthy male volunteers, not pre-selected for BAT presence or activity and without prior BAT cold-activation, on two occasions, following an infusion with hydrocortisone ( $0.2 \text{ mg.kg}^{-1}.\text{min}^{-1}$  for 14 h) and saline, respectively. Infusions were given in a randomized double-blind order. They underwent assessment of supraclavicular BAT temperature using infrared thermography following a mixed meal, and during  $\beta$ -AR stimulation with isoprenaline ( $25 \text{ ng.kg fat-free mass}^{-1}.\text{min}^{-1}$  for 60 min) in the fasting state.

**Results.** During hydrocortisone infusion, BAT temperature increased both under fasting basal conditions and during  $\beta$ -AR stimulation. We observed a BAT temperature threshold, which was not exceeded despite maximal  $\beta$ -AR activation. We conclude that BAT thermogenesis is present in humans under near-normal conditions. Glucocorticoids modulate BAT function, representing important physiological endocrine regulation of body temperature at times of acute stress.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

**Abbreviations:** <sup>18</sup>F-DG-PET/CT, <sup>18</sup>F-fluorodeoxyglucose-positron emission tomography/computed tomography; ATP, adenosine triphosphate; AUC, area under the curve; BAT, brown adipose tissue; CRF, clinical research facility; DXA, dual energy x-ray absorptiometry; HC, hydrocortisone; HOMA, homeostatic model assessment; ISO, isoprenaline; IT, infrared thermography; NEFA, non-esterified fatty acids; NIHR, National Institute for Health Research; NRES, National Research Ethics Service; REC, research ethics committee; tAUC, time-averaged area under the curve; T<sub>REF</sub>, reference point temperature; T<sub>SCR</sub>, supraclavicular temperature; UCP1, uncoupling protein 1;  $\beta$ -AR, beta adrenoceptor.

\* Correspondence to: M.E. Symonds, Division of Child Health, Obstetrics & Gynaecology, School of Medicine, University of Nottingham, Nottingham, NG7 2UH, UK. Tel.: +44 115 823 0625.

\*\* Correspondence to: K.N. Manolopoulos, Institute of Metabolism and Systems Research, University of Birmingham, IBR Tower Level 2, Room 228, Birmingham, B15 2TT, UK. Tel.: +44 121 414 7525.

E-mail address: [k.manolopoulos@bham.ac.uk](mailto:k.manolopoulos@bham.ac.uk) (K.N. Manolopoulos).

<http://dx.doi.org/10.1016/j.metabol.2017.01.024>

0026-0495/© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

There is increasing evidence that brown adipose tissue (BAT) has important physiological roles beyond thermoregulation in newborn infants and rodents [1]. Adult humans have significant amounts of BAT [2] and, as a highly metabolic tissue with the capacity to oxidize both glucose and lipid, attention has turned to its involvement in the pathogenesis of obesity and the metabolic syndrome [3]. BAT is characterized by the presence of uncoupling protein (UCP) 1 which uncouples adenosine triphosphate (ATP) production by the mitochondrial respiratory chain, allowing the dissipation of excess chemical energy as heat [4]. The principal factors regulating BAT function in healthy adults have yet to be fully established due, in part, to the technical limitations of assessing BAT function *in vivo*. The majority of studies in humans have used 18F-fluorodeoxyglucose-positron emission tomography/computed tomography ( $^{18}\text{F}$ FDG-PET/CT) as the gold standard to assess BAT activity, but this is constrained by exposure to ionizing radiation, the scanning protocols involved [5] and its unsuitability for live tracking of BAT activation especially after feeding. Systemic  $\beta$ -adrenoceptor ( $\beta$ -AR) activation promotes BAT activity in humans [6], but the role of other endocrine factors remains largely unknown. The pre-partum elevation of cortisol is pivotal in the initiation of nonshivering BAT thermogenesis at birth [7], and glucocorticoids have recently been proposed as regulators of BAT activity in healthy adult females [8] and in individuals pre-selected for the presence of active BAT [9]. BAT has also been considered to contribute to dietary-induced thermogenesis [10,11], although this concept remains controversial [12]. We, therefore, studied whether BAT is activated by feeding, or by an acute increase in cortisol under basal and  $\beta$ -AR stimulated conditions.

## 2. Materials and Methods

### 2.1. Subjects

Eight healthy male volunteers participated in this randomized, double-blind, placebo controlled study, conducted between January and March 2015. Individuals were recruited using print and electronic advertising and none was selected or screened on the basis of presence of any active BAT. All subjects underwent a medical evaluation during a screening visit to ensure they were healthy. No subject had any significant past medical history, smoked tobacco or took any regular medications that could affect the study's outcome measures.

### 2.2. Study Approval

The study was approved by the Edgbaston NRES Committee, UK (REC reference 14/WM/1085). All participants provided written informed consent.

### 2.3. Study Design

#### 2.3.1. Clinical Research Facility

All parts of this study were conducted in a temperature controlled room at the National Institute for Health Research (NIHR)/

Wellcome Trust Clinical Research Facility (CRF) of the University of Birmingham at the Queen Elizabeth Hospital Birmingham, UK. Room temperature was held constant at 23–26 °C and was monitored using an ambient temperature probe.

#### 2.3.2. Anthropometric Measurements


Measurements were taken during the screening visit. Waist circumference was measured midway between the lower margin of the last palpable rib and the top of the iliac crest, and hip circumference at the level of the greater trochanters. Total and regional fat masses were measured by dual-energy x-ray absorptiometry (DXA). Visceral fat mass was estimated by DXA using a proprietary algorithm provided by the manufacturer [13]. Core temperature was measured with a tympanic thermometer.

#### 2.3.3. Study Visits

Study visits were identical, except for the nature of overnight infusion, and were at least 2 weeks apart (Fig. 1A). Subjects were admitted to the CRF in the afternoon, and a cannula for infusion purposes was inserted into a right antecubital fossa vein. At 1800 h, they were served a standardized calorie-controlled meal (vegetable lasagne; total energy 2634 kJ; typical nutritional values per 100 g of product: 1.9 g fat, 12.2 g carbohydrates, 3.3 g protein, 1.5 g fiber), and then fasted until study completion the next day. BAT thermogenesis assessment was performed immediately before and after the meal, which was ingested within 20 min and was accompanied by tap water at room temperature. At 1900 h, a constant infusion of either hydrocortisone (HC, 0.2 mg.kg<sup>-1</sup>.h<sup>-1</sup>) or normal saline (control study visit) was started and given until study completion the following day. Infusions were administered in a double-blind, randomized fashion. At 2200 h, lights were switched off for night rest. In the morning, cannulations for blood sampling purposes were performed and, at 0900 h, the isoprenaline infusion protocol commenced. After baseline measurements for 45 min, a one-step infusion of isoprenaline (ISO, 25 ng.kg fat-free mass<sup>-1</sup>.min<sup>-1</sup>) was given for 60 min. BAT thermogenic activity was measured at baseline and throughout the infusion.

### 2.4. BAT Thermogenesis Assessment

An infrared thermography (IT) camera (FLIR E60 2.3 Megapixel Infrared Camera; FLIR Systems, Danderyd, Sweden) was used to acquire images of the anterior neck and upper chest region, which were sequentially analyzed and processed by an automated analysis program, as described previously [14]. Areas of interest for temperature analysis were the supraclavicular region ( $T_{\text{SCR}}$ ) representing BAT, and a non-adipose tissue reference point ( $T_{\text{REF}}$ ) on the chest, close to the xiphoid. In addition, during the periods of IT, two skin contact temperature sensors (iButton DS1922L, Maxim Integrated, Womersley, UK) recording skin temperature every minute were taped within the supraclavicular fossa (main BAT site) and lateral to the umbilicus (white adipose tissue). For skin contact temperature measurements, data were collected every minute, and analysis was performed using 5-min averages. For meal measurements, the mean of both study days was calculated. Fasting and pre-ISO baseline were defined as the average of time points –15 to 0 min. Postprandial and peak post-ISO infusion periods were defined as time points 0 to 15 min and 40 to 50 min, respectively. For the duration of the study,



**Fig. 1 – Study design and temperature responses to a meal.** Each participant underwent the study twice, whereby BAT thermogenic activity was studied with infrared thermography before and after a standardized meal, followed by either a 14 h overnight constant hydrocortisone or normal saline (control) infusion. Infusions were given in a randomized, double-blind order and continued during and after  $\beta$ -adrenoceptor stimulation with isoprenaline (A). Mean changes in supraclavicular region ( $T_{\text{SCR}}$ , gray circles) and non-adipose tissue reference ( $T_{\text{REF}}$ , open squares) temperatures (dotted line indicates time of meal) (B), individual responses (fasting, open squares; postprandial, black squares) (C), and changes in core temperature (fasting, open circles; postprandial, black squares) (D) following the meal. \* $p < 0.05$  compared to fasting baseline,  $n = 8$ .

participants were wearing a hospital gown, with their torso exposed for the duration of all measurements.

### 2.5. Analytical Methods

Blood samples were drawn into heparinized syringes, and plasma was prepared rapidly at  $4^{\circ}\text{C}$  and immediately frozen at  $-80^{\circ}\text{C}$  before analysis. Plasma glucose and NEFA concentrations were measured enzymatically using commercially available kits on an ILAB600 or ILAB650 clinical analyzer (Instrumentation Laboratory UK, Warrington, UK). Insulin and C-peptide were measured by ELISA (Invitron, Monmouth, UK) at a reference laboratory (Diabetes Research Unit Cymru, Swansea University, UK). Cortisol was measured by a colorimetric assay (R&D Systems, Abingdon, UK).

### 2.6. Calculations and Statistics

Indexes of  $\beta$ -cell function and insulin resistance were calculated according to the homeostatic model assessment (HOMA) method, whereby the mean of three consecutive plasma glucose and insulin postabsorptive measurements was used. Energy expenditure was calculated based on heart rate, age and weight as previously described [15]. Area under the curve (AUC) was calculated using the trapezoid rule and is presented as a time-averaged value (tAUC; AUC divided by the relevant time period). Comparisons between groups were analyzed using t test or non-

parametric tests for data that were not normally distributed. A  $p < 0.05$  was considered statistically significant. Based on previous studies using a similar integrative physiology design [16], the sample size was designed to have 85% power to detect a difference of 0.75 standard deviations at the 5% significance level for metabolic parameters. Data were analyzed using IBM Statistics for Windows v21 and GraphPad Prism for Windows v6.05. All data are presented as mean  $\pm$  SEM, unless otherwise stated.

## 3. Results

### 3.1. Meal Ingestion Leads to Increased BAT Thermogenic Activity

Baseline anthropometric and metabolic characteristics of participants are shown in Table 1 and environmental temperature data for each individual study day are shown in Supplemental Table 1. There was no difference in outside or room temperature between study days. Following the mixed meal, postprandial  $T_{\text{SCR}}$  increased, whereas  $T_{\text{REF}}$  remained stable (Fig. 1B). All participants responded with an increase in BAT thermogenic activity (Fig. 1C), while core temperature did not change (Fig. 1D). Skin contact measurements showed a similar postprandial temperature increase of  $0.39 \pm 0.10^{\circ}\text{C}$  over supraclavicular BAT ( $p < 0.05$  compared to fasting), whereas skin temperature over white adipose tissue remained the same.



**Table 1 – Baseline anthropometric and metabolic characteristics of participants, n = 8.**

Characteristic	
Age (years)	20 (18–34)
Weight (kg)	75.0 (61.5–81.7)
BMI (kg/m <sup>2</sup> )	23.0 (20.6–24.2)
WHR	0.84 (0.78–0.9)
Trunk fat (kg)	7.7 (5.1–9.2)
Leg fat (kg)	5.5 (4.3–7.5)
Visceral fat (kg)	0.2 (0.1–0.4)
Systolic BP (mmHg)	124 (105–145)
Diastolic BP (mmHg)	72 (59–79)
Heart rate (bpm)	58 (53–62)
Fasting glucose (mg/dL)	113 (89–122)
Fasting insulin (pmol/L)	30.6 (14.4–68.5)
Fasting NEFA (μmol/L)	707 (450–873)
TSH (mIU/L)	1.30 (0.93–4.83)
FT4 (pmol/L)	17.7 (15.0–18.8)

Median and range shown. BMI, body mass index; WHR, waist-to-hip ratio; BP, blood pressure; TSH, thyroid stimulating hormone; FT4, free thyroxine.

### 3.2. Acute Hypercortisolemia Induces Peripheral Insulin Resistance and Increases Basal BAT Thermogenic Activity


Overnight HC infusion resulted in significantly increased plasma cortisol concentrations (Fig. 2A). From a metabolic perspective, basal plasma non-esterified fatty acids (NEFAs) were high due to fasting (Fig. 2B). HC increased basal NEFA

and glucose (Fig. 2C), as well as insulin (basal insulin  $30.4 \pm 6.0$  pmol/L vs.  $55.2 \pm 7.4$  pmol/L,  $p = 0.025$  control compared to hypercortisolemia) and C-peptide concentrations (basal C-peptide  $0.25 \pm 0.03$  pmol/mL vs.  $0.38 \pm 0.04$  pmol/mL,  $p = 0.001$  control compared to hypercortisolemia). In line with this, HOMA indices of peripheral insulin resistance increased (HOMA IR index control  $0.62 \pm 0.11$  vs. hypercortisolemia  $1.11 \pm 0.16$ ,  $p = 0.016$ ) (Table 2).

Acute hypercortisolemia increased  $T_{SCR}$  in the basal state (Fig. 2D). This was accompanied by an increase in basal core temperature (Fig. 2E), but we did not observe any effect on blood pressure or heart rate (Fig. 2F).

### 3.3. Acute $\beta$ -AR Stimulation Increases BAT Thermogenic Activity During Control and Hypercortisolemia Conditions

From a metabolic perspective, ISO infusion significantly increased systemic NEFA concentrations, both under control and hypercortisolemia conditions (Fig. 2B). Despite the augmentation of basal systemic lipolysis by HC, the  $\beta$ -AR dependent rise in plasma NEFA was of similar magnitude compared to control conditions ( $\Delta$  AUC  $953 \pm 155$  vs  $979 \pm 175$  μmol/L;  $p = 0.926$  compared to control). Following the initial peak, there was a sharp decline in NEFA concentrations despite continuing ISO infusion. Control plasma glucose concentrations were unaffected by ISO, while the observed increase in concentrations during HC infusion is due to glucocorticoid-induced peripheral tissue insulin resistance (Fig. 2C). This is supported by the concomitant changes in insulin, C-peptide and



**Fig. 2 – Metabolic and cardiovascular responses following saline infusion (S) or hydrocortisone infusion (HC) at baseline and during an isoprenaline infusion (ISO, dotted lines indicate infusion period). Plasma cortisol concentrations (saline, open circles; hydrocortisone, black squares) (A), non-esterified fatty acids (NEFA) (saline, open circles; hydrocortisone, black squares) (B), glucose (saline, open circles; hydrocortisone, black squares) (C), supraclavicular temperature ( $T_{SCR}$ ) (saline, open circles; hydrocortisone, black circles) (D) and core temperature at baseline (saline, open squares; hydrocortisone, black squares) (E). Systolic (sys) and diastolic (dia) blood pressure (BP) and heart rate (HR) at baseline (saline, open bars; hydrocortisone, black bars) (F). \* $p < 0.05$  vs. control, n = 8.**


**Table 2 – Comparison of insulin and C-peptide AUC, and homeostatic model assessment (HOMA) indexes during basal and isoprenaline-stimulated conditions, n = 8.**

	Control		Cortisol		P value control vs cortisol		P value basal vs isoprenaline	
	Basal	ISO	Basal	ISO	Basal	ISO	Control	Cortisol
Insulin (pmol/L)	30.4 ± 6.0	100.4 ± 15.7	55.2 ± 7.4	206.7 ± 30.5	0.025	0.012	0.012	0.012
C-peptide (pmol/mL)	0.25 ± 0.03	0.47 ± 0.05	0.38 ± 0.04	0.90 ± 0.09	0.001	0.001	<0.0001	<0.0001
			Control		Cortisol		P value	
Δ insulin basal to ISO (pmol/L)			70.0 ± 10.1		151.5 ± 28.9		0.012	
Δ C-peptide basal to ISO (pmol/mL)			0.22 ± 0.03		0.52 ± 0.07		0.012	
HOMA %B			42.2 ± 5.9		40.5 ± 4.5		0.731	
HOMA %S			192.2 ± 29.5		105.9 ± 19.2		0.004	
HOMA IR			0.62 ± 0.11		1.11 ± 0.16		0.016	

Mean and SEM shown.

HOMA indexes showing a decrease in glucose sensitivity despite a significant increase in insulin and C-peptide concentrations during ISO infusion (Table 1). Expectedly, non-selective β-AR stimulation with ISO increased heart rate and systolic blood pressure, responses not significantly affected by HC (Fig. 2F and Supplemental Fig. 1). Basal and ISO-induced BAT thermogenic activity measures did not show any significant correlation with BMI or measures of adipose tissue distribution (data not shown).

Adrenergic stimulation resulted in a highly localized increase in temperature within the supraclavicular region, representative of BAT thermogenic activity, both under control and hypercortisolemic conditions (Fig. 3). All study participants responded to ISO with an increase in BAT temperature (Supplemental Fig. 2). Under control conditions, ISO increased T<sub>SCR</sub> by 0.7 °C, plateaued and then returned to baseline after the infusion, implying cessation of β-adrenergic-mediated BAT



**Fig. 3 – BAT thermogenic responses.** Supraclavicular temperature (T<sub>SCR</sub>) and non-adipose tissue reference point (T<sub>REF</sub>) following saline (S) infusion (T<sub>SCR</sub>, gray circles; T<sub>REF</sub> open squares) (A) or hydrocortisone (HC) infusion (T<sub>SCR</sub>, gray circles; T<sub>REF</sub> open squares) (B) during and after isoprenaline stimulation (ISO, dotted lines indicate infusion period). Individual peak BAT temperatures during ISO (S, open circles; HC, black circles) (C). Change in temperature during ISO or following a standardized meal (T<sub>SCR</sub>, gray bars; T<sub>REF</sub> open bars) (D). \*p < 0.001 vs. basal, †p < 0.05 vs. saline, #p < 0.001 vs. saline, ##p < 0.001 vs. T<sub>REF</sub>, n = 8.

thermogenesis (Fig. 3A). These responses to ISO were similar during hypercortisolemia (Fig. 3B), whereby peak  $T_{SCR}$  was slightly higher (Fig. 3C). The ISO-induced  $T_{SCR}$  increase was greater than the physiological stimulus of diet-induced thermogenesis (Fig. 3D). Skin temperature showed similar results (Supplemental Fig. 3). Energy expenditure increased significantly during ISO (Supplemental Fig. 4). During hypercortisolemia, ISO-induced energy expenditure was closely correlated with basal  $T_{SCR}$  during control conditions (Pearson  $r = 0.742$ ,  $p = 0.035$ ) and peak  $T_{SCR}$  during HC ( $r = 0.870$ ,  $p = 0.005$ ). In response to ISO, peak core temperature was similar between control and hypercortisolemia conditions (Supplemental Fig. 5).

#### 4. Discussion

Human supraclavicular BAT is characterized by the presence of thermogenically functional UCP1, with a respiratory capacity that substantially exceeds that of white fat [17]. Understanding the endocrine factors regulating BAT function is an important prerequisite before being able to utilize the metabolic capabilities of this tissue. In this study we sought to study BAT *in vivo* following exposure to a combination of physiological stimuli in order to determine the relative importance of diet and endocrine mediated effects.

BAT glucose uptake has been reported to be increased following a single carbohydrate-rich meal [18], although overfeeding for 24 h did not have any effect [19]. This has led to some controversy regarding the contribution of BAT to dietary-induced thermogenesis in humans. We sought to investigate this using a single standardized mixed meal, serving as a physiological stimulus. While we did not measure whole body energy expenditure, we observed selective temperature changes over the supraclavicular region only, immediately after the meal, suggesting direct BAT activation and not a thermic effect of food. Interestingly, from a mechanistic perspective, postprandial BAT activation would be characterized by both systemic cortisol secretion [20] and sympathetic  $\beta$ -AR stimulation [21], suggesting an acute maximal response following feeding.

Cortisol promotes important physiological maturation effects around the time of birth, including raised UCP1 abundance in adipose tissue [7,22]. However, in adult rodents, glucocorticoids inhibit BAT [23] by interfering with adrenergic signaling [24,25]. Human data are scarce with one study reporting dexamethasone-induced inhibition of UCP1 expression and metabolic rate in human brown adipocytes *in vitro* [26], and another reporting BAT activation following administration of the synthetic glucocorticoid prednisolone *in vivo* [9]. In our study, we chose hydrocortisone to model a physiological acute surge of cortisol, as seen during the perinatal period and at times of acute stress. We observed an increase of basal  $T_{SCR}$  during hypercortisolemia, supporting a physiological role for cortisol in BAT activation, as we did not observe any additive effects on blood pressure or heart rate. The duration of the infusion was chosen to allow for glucocorticoid-mediated genomic effects to take place [27]. While the achieved plasma cortisol concentrations

were in excess of those typical for acute stress [28], it is important to note that tissue-responsiveness can be determined by tissue-specific glucocorticoid metabolism rather than absolute plasma concentrations [29]. Taken together our data indicate the positive relationship between cortisol and BAT temperature as previously indicated from a small study on healthy adult females [8].

$\beta$ -AR stimulation induces BAT thermogenesis in humans [6,30], although findings are inconsistent depending on the  $\beta$ -AR employed [31,32]. We found a localized increase in supraclavicular temperature during ISO infusion both under control and hypercortisolemia conditions. This temperature change was temporally limited for the duration of the infusion, suggesting underlying BAT activation.  $T_{SCR}$  responses for all subjects increased within the first 5 min which is in accordance with acute cold exposure on BAT [14]. The observed  $T_{SCR}$  plateau is suggestive of a limit to BAT thermogenesis *in vivo*. The concomitant sharp decline in NEFA concentrations during the later stages of the infusion is consistent with  $\beta$ -AR desensitization due to maximal receptor stimulation [16]. The finding of a slightly higher peak  $T_{SCR}$  during hypercortisolemia suggests a minor synergistic effect between cortisol and  $\beta$ -AR stimulation. Interestingly, the two pathways are intrinsically connected as catecholamine synthesis is under glucocorticoid control [33].

Previous studies have shown that active BAT decreases with age and obesity, and its activation varies between sexes [14,34]. We studied BAT activity in healthy males using IT to assess temperature changes in the supraclavicular region and a non-adipose tissue reference point. Supraclavicular skin temperature increases upon BAT activation [35,36] and IT has been shown to measure changes in skin temperature overlying the main BAT depot in humans [14]. It has been confirmed as a reliable alternative for *in vivo* BAT activity assessment, correlating with  $^{18}\text{F}$ FDG-PET/CT [9,37], with the additional benefit of enabling real-time tracking of temperature changes. IT-derived BAT temperature measurements might be influenced by subcutaneous adipose tissue thickness [38], however, in our study participants were lean and we monitored dynamic temperature changes over time, as opposed to a single, static measurement. Adrenoceptor-induced vasodilation, both as a result of HC and ISO infusions, could increase skin blood flow and interfere with IT measurements. However, compared to  $T_{SCR}$ , there were clear temporal differences in the change in  $T_{REF}$  which showed a later initial increase and a sustained increase post-infusion. Overall, we demonstrate a BAT-specific thermogenic and vasodilation response to both HC and ISO, clearly differentiated from non-BAT reference areas, confirming that the temperature responses we measured are confined to BAT.

Our findings support previous studies showing  $\beta$ -AR stimulation as a means of activating BAT in humans [6,30], confirming IT as a sensitive, non-invasive method for the *in vivo* assessment of BAT function in humans under near-normal conditions [9,14,37]. This is particularly important when comparing our results with those studies using glucose tracer uptake as an index of BAT activity. Similar ISO doses did not show any significant BAT glucose tracer uptake, likely due to competition between the tracer and fatty acids from ISO-induced lipolysis combined with increased insulin

resistance [32]. Given that BAT primarily utilizes fatty acids for heat generation [1], it is possible that  $^{18}\text{F}$ FDG-PET/CT underestimates the amount of active BAT in humans. This limitation has led to the development of alternative BAT assessment methods, in addition to IT, either using different PET/CT tracers [39], or based on magnetic resonance imaging techniques [40]. We demonstrate that temperature changes in the supraclavicular area upon  $\beta$ -AR stimulation are indicative of localized BAT activity in a cohort of unselected young individuals, maintained at room temperature. This supports the prospect of harnessing BAT activity and the associated increase in energy expenditure as a potential treatment for metabolic diseases. We provide further evidence that in humans, in contrast to rodents, acute hypercortisolemia does not inhibit BAT function, but results in BAT activation [9]. Despite this, there is a threshold of activity that cannot be overcome even during maximal short-term  $\beta$ -AR stimulation.

Our study has some limitations by design, including the small size of our sample, although it is standard for a healthy volunteer study of this type. The acute infusion of hydrocortisone limits the conclusions we can draw in relation to states of chronic glucocorticoid excess that are associated with profound metabolic changes, i.e. Cushing's Syndrome. In addition, the concomitant induction of relative insulin resistance during hydrocortisone infusion might have obscured glucocorticoid-specific effects on BAT function, especially since there is a complex relationship between insulin-mediated glucose uptake and BAT perfusion and activity *in vivo* [41]. The strengths of our study are the randomized double-blind design of the infusion protocol and that all measurements were carried out within a short period of time, reducing the confounding effect of variations seasonal temperature, and thus endogenous BAT activity. In addition, by using IT for the assessment of BAT thermogenic activity, we were able to perform live tracking of BAT function in response to experimental stimuli.

In conclusion, glucocorticoids modulate BAT thermogenesis and may represent an important physiological mechanism for maintaining human body temperature at times of acute stress. Our study suggests that transient stress could act to promote BAT function. This suggests that depending on the type and magnitude of stress BAT could be utilized to improve body weight regulation and metabolic homeostasis.

### Authors' contributions

KNM and MES designed the study; HS and KNM conducted the experiments and analyzed the data with JL, HS, MES, JL, HB, DS and KNM wrote the manuscript.

### Funding

The study was funded by a Starter Grant for Clinical Lecturers of the Academy of Medical Sciences awarded to KNM; HS is supported by a Medical Research Council Doctoral Training Award.

### Acknowledgments

We thank the research nursing staff at the NIHR/Wellcome Trust Clinical Research Facility at the Queen Elizabeth Hospital Birmingham for excellent study support, and the volunteers for participating.

### Conflict of interest

No conflicts of interest, financial or otherwise, are declared by the authors.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.metabol.2017.01.024>.

### REFERENCES

- [1] Ouellet V, Labbe SM, Blondin DP, Phoenix S, Guerin B, Haman F, et al. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *J Clin Invest* 2012;122:545–52.
- [2] van Marken Lichtenbelt WD, Vanhommerig JW, Smulders NM, Drossaerts JM, Kemerink GJ, Bouvy ND, et al. Cold-activated brown adipose tissue in healthy men. *N Engl J Med* 2009;360:1500–8.
- [3] Nedergaard J, Cannon B. The changed metabolic world with human brown adipose tissue: therapeutic visions. *Cell Metab* 2010;11:268–72.
- [4] Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev* 2004;84:277–359.
- [5] Cypess AM, Haft CR, Laughlin MR, Hu HH. Brown fat in humans: consensus points and experimental guidelines. *Cell Metab* 2014;20:408–15.
- [6] Cypess AM, Weiner LS, Roberts-Toler C, Franquet Elia E, Kessler SH, Kahn PA, et al. Activation of human brown adipose tissue by a beta3-adrenergic receptor agonist. *Cell Metab* 2015;21:33–8.
- [7] Mostyn A, Pearce S, Budge H, Elmes M, Forhead AJ, Fowden AL, et al. Influence of cortisol on adipose tissue development in the fetal sheep during late gestation. *J Endocrinol* 2003;176:23–30.
- [8] Robinson LJ, Law JM, Symonds ME, Budge H. Brown adipose tissue activation as measured by infrared thermography by mild anticipatory psychological stress in lean healthy females. *Exp Physiol* 2016;101:549–57.
- [9] Ramage LE, Akyol M, Fletcher AM, Forsythe J, Nixon M, Carter RN, et al. Glucocorticoids acutely increase brown adipose tissue activity in humans, revealing species-specific differences in UCP-1 regulation. *Cell Metab* 2016;24:130–41.
- [10] Symonds ME, Pope M, Budge H. The ontogeny of brown adipose tissue. *Annu Rev Nutr* 2015;35:295–320.
- [11] Hibi M, Oishi S, Matsushita M, Yoneshiro T, Yamaguchi T, Usui C, et al. Brown adipose tissue is involved in diet-induced thermogenesis and whole-body fat utilization in healthy humans. *Int J Obes* 2016.
- [12] Kozak LP. Brown fat and the myth of diet-induced thermogenesis. *Cell Metab* 2010;11:263–7.



- [13] Kaul S, Rothney MP, Peters DM, Wacker WK, Davis CE, Shapiro MD, et al. Dual-energy x-ray absorptiometry for quantification of visceral fat. *Obesity (Silver Spring)* 2012;20:1313–8.
- [14] Symonds ME, Henderson K, Elvidge L, Bosman C, Sharkey D, Perkins AC, et al. Thermal imaging to assess age-related changes of skin temperature within the supraclavicular region co-locating with brown adipose tissue in healthy children. *J Pediatr* 2012;161:892–8.
- [15] Keytel LR, Goedecke JH, Noakes TD, Hiiloskorpi H, Laukkanen R, van der Merwe L, et al. Prediction of energy expenditure from heart rate monitoring during submaximal exercise. *J Sports Sci* 2005;23:289–97.
- [16] Manolopoulos KN, Karpe F, Frayn KN. Marked resistance of femoral adipose tissue blood flow and lipolysis to adrenaline in vivo. *Diabetologia* 2012;55:3029–37.
- [17] Porter C, Herndon DN, Chondronikola M, Chao T, Annamalai P, Bhattarai N, et al. Human and mouse brown adipose tissue mitochondria have comparable UCP1 function. *Cell Metab* 2016;24:246–55.
- [18] Vosselman MJ, Brans B, van der Lans AA, Wierts R, van Baak MA, Mottaghy FM, et al. Brown adipose tissue activity after a high-calorie meal in humans. *Am J Clin Nutr* 2013;98:57–64.
- [19] Schlogl M, Piaggi P, Thiyyagura P, Reiman EM, Chen K, Lutrin C, et al. Overfeeding over 24 hours does not activate brown adipose tissue in humans. *J Clin Endocrinol Metab* 2013;98:E1956–60.
- [20] Korbonits M, Trainer PJ, Nelson ML, Howse I, Kopelman PG, Besser GM, et al. Differential stimulation of cortisol and dehydroepiandrosterone levels by food in obese and normal subjects: relation to body fat distribution. *Clin Endocrinol* 1996;45:699–706.
- [21] van Baak MA. Meal-induced activation of the sympathetic nervous system and its cardiovascular and thermogenic effects in man. *Physiol Behav* 2008;94:178–86.
- [22] Symonds ME, Pope M, Budge H. Adipose tissue development during early life: novel insights into energy balance from small and large mammals. *Proc Nutr Soc* 2012;71:363–70.
- [23] van den Beukel JC, Grefhorst A, Quarta C, Steenbergen J, Mastroberardino PG, Lombes M, et al. Direct activating effects of adrenocorticotrophic hormone (ACTH) on brown adipose tissue are attenuated by corticosterone. *FASEB J* 2014;28:4857–67.
- [24] Soumano K, Desbiens S, Rabelo R, Bakopanos E, Camirand A, Silva JE. Glucocorticoids inhibit the transcriptional response of the uncoupling protein-1 gene to adrenergic stimulation in a brown adipose cell line. *Mol Cell Endocrinol* 2000;165:7–15.
- [25] Martinez-deMena R, Calvo RM, Garcia L, Obregon MJ. Effect of glucocorticoids on the activity, expression and proximal promoter of type II deiodinase in rat brown adipocytes. *Mol Cell Endocrinol* 2016;428:58–67.
- [26] Barclay JL, Agada H, Jang C, Ward M, Wetzig N, Ho KK. Effects of glucocorticoids on human brown adipocytes. *J Endocrinol* 2015;224:139–47.
- [27] Biddie SC, Conway-Campbell BL, Lightman SL. Dynamic regulation of glucocorticoid signalling in health and disease. *Rheumatology (Oxford)* 2012;51:403–12.
- [28] Carmen Wong KY, Wong V, Ho JT, Torpy DJ, McLean M, Cheung NW. High cortisol levels in hyperglycaemic myocardial infarct patients signify stress hyperglycaemia and predict subsequent normalization of glucose tolerance. *Clin Endocrinol* 2010;72:189–95.
- [29] Morgan SA, Hassan-Smith ZK, Lavery GG. Mechanisms in endocrinology: tissue-specific activation of cortisol in Cushing's syndrome. *Eur J Endocrinol* 2016;175:R83–9.
- [30] Carey AL, Formosa MF, Van Every B, Bertovic D, Eikelis N, Lambert GW, et al. Ephedrine activates brown adipose tissue in lean but not obese humans. *Diabetologia* 2013;56:147–55.
- [31] Cypess AM, Chen YC, Sze C, Wang K, English J, Chan O, et al. Cold but not sympathomimetics activates human brown adipose tissue in vivo. *Proc Natl Acad Sci U S A* 2012;109:10001–5.
- [32] Vosselman MJ, van der Lans AA, Brans B, Wierts R, van Baak MA, Schrauwen P, et al. Systemic beta-adrenergic stimulation of thermogenesis is not accompanied by brown adipose tissue activity in humans. *Diabetes* 2012;61:3106–13.
- [33] Rozansky DJ, Wu H, Tang K, Parmer RJ, O'Connor DT. Glucocorticoid activation of chromogranin a gene expression. Identification and characterization of a novel glucocorticoid response element. *J Clin Invest* 1994;94:2357–68.
- [34] Yoneshiro T, Aita S, Matsushita M, Okamatsu-Ogura Y, Kameya T, Kawai Y, et al. Age-related decrease in cold-activated brown adipose tissue and accumulation of body fat in healthy humans. *Obesity (Silver Spring)* 2011;19:1755–60.
- [35] Boon MR, Bakker LE, van der Linden RA, Pereira Arias-Bouda L, Smit F, Verberne HJ, et al. Supraclavicular skin temperature as a measure of 18F-FDG uptake by BAT in human subjects. *PLoS One* 2014;9:e98822.
- [36] van der Lans AA, Vosselman MJ, Hanssen MJ, Brans B, van Marken Lichtenbelt WD. Supraclavicular skin temperature and BAT activity in lean healthy adults. *J Physiol Sci* 2016;66:77–83.
- [37] Salem V, Izzi-Engbeaya C, Coello C, Thomas DB, Chambers ES, Comninos AN, et al. Glucagon increases energy expenditure independently of brown adipose tissue activation in humans. *Diabetes Obes Metab* 2016;18:72–81.
- [38] Gatidis S, Schmidt H, Pfannenbergl CA, Nikolaou K, Schick F, Schwenzler NF. Is it possible to detect activated brown adipose tissue in humans using single-time-point infrared thermography under thermoneutral conditions? Impact of BMI and subcutaneous adipose tissue thickness. *PLoS One* 2016;11:e0151152.
- [39] Hwang JJ, Yeckel CW, Gallezot J-D, Aguiar RB-D, Ersahin D, Gao H, et al. Imaging human brown adipose tissue under room temperature conditions with 11C-MRB, a selective norepinephrine transporter PET ligand. *Metab Clin Exp* 2015;64:747–55.
- [40] Holstila M, Virtanen KA, Grönroos TJ, Laine J, Lepomäki V, Saunavaara J, et al. Measurement of brown adipose tissue mass using a novel dual-echo magnetic resonance imaging approach: a validation study. *Metab Clin Exp* 2013;62:1189–98.
- [41] Orava J, Nuutila P, Lidell Martin E, Oikonen V, Noponen T, Viljanen T, et al. Different metabolic responses of human brown adipose tissue to activation by cold and insulin. *Cell Metab* 2011;14:272–9.