

Adipogenic differentiation of MSC alters their immunomodulatory properties in a tissue-specific manner

Munir, Hafsa; Clarke, Lewis; Sheriff, Lozan; Kemble, Samuel; Nayar, Saba; Barone, Francesca; Nash, Gerard; McGettrick, Helen

DOI:

[10.1002/stem.2622](https://doi.org/10.1002/stem.2622)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Munir, H, Clarke, L, Sheriff, L, Kemble, S, Nayar, S, Barone, F, Nash, G & McGettrick, H 2017, 'Adipogenic differentiation of MSC alters their immunomodulatory properties in a tissue-specific manner', *Stem Cells*, vol. 35, no. 6, pp. 1636–1646. <https://doi.org/10.1002/stem.2622>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Adipogenic Differentiation of Mesenchymal Stem Cells Alters Their Immunomodulatory Properties in a Tissue-Specific Manner

HAFSA MUNIR,^a LEWIS S. C. WARD,^b LOZAN SHERIFF,^a SAMUEL KEMBLE,^b SABA NAYAR,^b FRANCESCA BARONE,^b GERARD B. NASH,^a HELEN M. MCGETTRICK ^{id b}

Key Words. Mesenchymal stem cells • Adipocytes • Endothelial cells • Neutrophils • Inflammation • Tissue-specific

^aInstitute for Cardiovascular Sciences, College of Medical and Dental Sciences and ^bInstitute of Inflammation and Ageing, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom

Correspondence: Helen M. McGettrick, PhD, Institute of Inflammation and Ageing, University of Birmingham, Birmingham, B15 2WB, U.K. Telephone: 44-121-371-3228; e-mail: h.m.mcgettrick@bham.ac.uk

Received October 6, 2016; accepted for publication March 10, 2017; first published online in *STEM CELLS EXPRESS* April 4, 2017.

© AlphaMed Press
1066-5099/2017/\$30.00/0

<http://dx.doi.org/10.1002/stem.2622>

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Chronic inflammation is associated with formation of ectopic fat deposits that might represent damage-induced aberrant mesenchymal stem cell (MSC) differentiation. Such deposits are associated with increased levels of inflammatory infiltrate and poor prognosis. Here we tested the hypothesis that differentiation from MSC to adipocytes in inflamed tissue might contribute to chronicity through loss of immunomodulatory function. We assessed the effects of adipogenic differentiation of MSC isolated from bone marrow or adipose tissue on their capacity to regulate neutrophil recruitment by endothelial cells and compared the differentiated cells to primary adipocytes from adipose tissue. Bone marrow derived MSC were immunosuppressive, inhibiting neutrophil recruitment to TNF α -treated endothelial cells (EC), but MSC-derived adipocytes were no longer able to suppress neutrophil adhesion. Changes in IL-6 and TGF β 1 signalling appeared critical for the loss of the immunosuppressive phenotype. In contrast, native stromal cells, adipocytes derived from them, and mature adipocytes from adipose tissue were all immunoprotective. Thus disruption of normal tissue stroma homeostasis, as occurs in chronic inflammatory diseases, might drive “abnormal” adipogenesis which adversely influences the behavior of MSC and contributes to pathogenic recruitment of leukocytes. Interestingly, stromal cells programmed in native fat tissue retain an immunoprotective phenotype. *STEM CELLS* 2017; 00:000–000

SIGNIFICANCE STATEMENT

Mesenchymal stem cells (MSC) can act to control inflammation, by regulating the ability of neutrophils to enter into inflamed tissues. Our data indicates that disruption of normal tissue homeostasis might drive “abnormal” MSC adipogenesis, causing the cells to lose their regulatory function. Thus, adipocytes may exist in at least two functional states: immunoprotective in healthy adipose tissue and stimulatory in sites of ectopic (e.g., chronic inflammation) fat deposition. Importantly, changes in the phenotype of MSC at sites of chronic inflammation may contribute to uncontrolled leukocyte infiltration and pathogenesis.

INTRODUCTION

Mesenchymal stem cells (MSC) are tissue-resident stromal precursors that undergo lineage-specific differentiation to repair damaged tissue and modulate a variety of immune responses [1, 2]. Indeed exploiting these properties therapeutically is the principle underpinning clinical trials using MSC in chronic inflammatory diseases [1]. Our recent studies revealed that MSC communicate with neighboring blood vascular endothelial cells (EC) to limit leukocyte recruitment during inflammation [3, 4]. This homeostatic role is a shared characteristic of MSC in a range of tissues [5]. Thus, MSC act as tissue-resident regulators of

leukocyte trafficking into inflamed peripheral tissue or might be delivered therapeutically to limit acute inflammatory infiltrates or to resolve chronic inflammatory disease.

Studies of the effects of lineage-specific differentiation on the immunoprotective properties of MSC have been rare and have yielded conflicting results [reviewed in ref. 6]. For example, human bone marrow (BM)-derived MSC have been reported to maintain their ability to suppress the proliferation of T cells from healthy individuals or rheumatoid arthritis (RA) patients following differentiation into osteoblasts [7–9] or chondrocytes [10]. Moreover, BMMSC-derived chondrocytes reduced T-cell activation and their ability to release proinflammatory

mediators *in vitro* [9, 10]. On the other hand, chondrogenically differentiated rat BMMSC were unable to suppress T-cell proliferation and activation [11] or dendritic cell maturation and crosstalk with T-cells [12]. Comparable studies following adipogenic differentiation have not been reported. Thus it remains unclear whether differentiation of MSC adversely affects their immunoprotective functions, in particular their ability to regulate EC and leukocyte trafficking during inflammation.

Adipogenic differentiation of MSC is particularly interesting because ectopic fat deposits and/or alterations in local adipose tissue are associated with a number of inflammatory disorders including Duchenne muscular dystrophy [13], myocardial infarction [14], and type II diabetes [15]. These deposits could be the result of inappropriate differentiation of tissue-resident MSC, possibly induced by inflammatory mediators in the affected tissue. Indeed platelet-derived growth factor (PDGFR) receptor-alpha positive (PDGFR α^+) skeletal muscle MSC were identified as the source of fat deposits in a murine model of glycerol-induced muscle fibre degeneration [13]. Interestingly, cardiotoxin-induced fibre degeneration did not lead to ectopic fat deposition in this muscle [13]. These data suggest that aberrant adipogenic differentiation in peripheral tissues may be stimulus-specific.

Chronic inflammation has been reported to alter the phenotype of BMMSC distal to the affected site. For instance, in RA and systemic lupus erythematosus, the proliferation and senescence of BMMSC were accelerated (e.g., [7, 8, 16–18]), and their osteogenic capacity was reduced [19], possibly worsening disease. What changes occur to MSC locally resident in chronically inflamed peripheral tissue is uncertain, but we have shown that tissue-resident stromal cells from chronically inflamed sites acquire a pathogenic proinflammatory phenotype, modifying EC to inappropriately recruit leukocytes [20–23]. The transformed stromal cells mediated their effects by altering the bioactivity of interleukin 6 (IL-6) or transforming growth factor beta 1 (TGF β 1), switching the function of these cytokines from an immunoprotective to pro-inflammatory state [20, 21, 23]. Whether adipogenic differentiation of MSC could drive a similar proinflammatory transformation is unknown.

The foregoing led us to test the hypothesis that aberrant differentiation of MSC to adipocytes might contribute to chronicity in inflamed tissue. We assessed the effects of adipogenic differentiation of BMMSC on their capacity to regulate neutrophil recruitment during inflammation *in vitro* and compared the differentiated cells to primary adipocytes from adipose tissue. Unlike MSC, MSC-derived adipocytes were not able to suppress neutrophil adhesion to inflamed EC. Changes in IL-6 and TGF β 1 signalling appeared critical for the loss of immunosuppressive phenotype. In contrast, native stromal cells, adipocytes derived from them and mature adipocytes (mAD) from adipose tissue were immunoprotective. Thus disruption of normal tissue stroma homeostasis, as occurs in chronic inflammatory diseases, might drive “abnormal” adipogenesis which adversely influences the behavior of MSC and contributes to pathogenic recruitment of leukocytes.

MATERIALS AND METHODS

Isolation and Culture of Human MSC or Adipocytes

Commercially available primary human BMMSC (Lonza Ltd., Basel, Switzerland) from healthy donors were obtained at

passage 2 and expanded three times in culture (i.e., to passage 5) in Mesenchymal Stem Cell Growth Medium (MSCGM) Bulletkit (Lonza Ltd., Basel, Switzerland, <http://www.lonza.com/>) [3–5]. Based on manufacturer's information, cells had undergone 11–12 doublings at passage 5 and underwent approximately 2.25 population doublings per passage [5].

Adipose derived mesenchymal stromal cells (ADSC; preadipocytes) or mature adipocytes (mAD) were isolated from subcutaneous adipose tissue collected from healthy patients undergoing abdominal surgery. Tissue pieces were suspended in Dulbecco's modified Eagle's medium (DMEM)/F-12 medium (Biosera, ZI du Bousquet, France, <http://www.biosera.com/>) supplemented with 25 mM HEPES (Sigma-Aldrich, Paisley, U.K., <http://www.sigmaaldrich.com>) containing 2 mg/ml collagenase type II (Sigma) at 37°C for 1 hour on a rotator. The cell suspension was filtered through a 70 μ m pore filter and centrifuged at 400g for 5 minutes. Mature adipocytes floated and were harvested at the air–liquid interface, resuspended in DMEM/F-12 medium and cultured for 24 hours prior to use. Pelleted ADSC were resuspended in DMEM/F-12 media supplemented with 10% fetal calf serum (FCS), 100 U/ml penicillin, and 100 μ g/ml streptomycin (AD basal media; all from Sigma) and cultured as previously described [24]. ADSC were passaged once before use.

Alternatively commercially available primary human ADSC from healthy donors were obtained at passage 2 and expanded three times in culture (i.e., to passage 5) in MSC growth medium 2 (C-28009; all from PromoCell GmbH, Heidelberg, Germany, <http://www.promocell.com/>).

Differentiation of MSC into Adipocytes

BMMSC were cultured for 21 days in adipogenic induction medium (Lonza) according to the manufacturer's instructions and as previously described [5]. To assess differentiation into MSC-derived adipocytes, samples were fixed in 10% neutral buffered formalin (Sigma) for 30 minutes, treated with 60% isopropanol (Sigma), and stained for 30 minutes with 0.3% oil red O (Sigma) dissolved in isopropanol. Samples were washed in distilled water and counterstained with hematoxylin solution (Sigma). Cells were imaged, and digitized images were acquired using an EVOS FL Imaging System (Thermo Scientific, Loughborough, U.K., <https://www.thermofisher.com/uk/en/home.html>) (Supporting Information Fig. 1A).

In addition, differentiation was confirmed by the upregulation of expression of the adipocyte genes: PPAR γ , C/EBP α , and FABP4. Briefly, mRNA was isolated from BMMSC and BMMSC-derived adipocytes using the RNeasy Mini Kit (Qiagen, Crawley, U.K., <http://www1.qiagen.com>) [25], converted to cDNA, and analyzed by quantitative polymerase chain reaction (qPCR) using Taqman Universal PCR Mastermix according to manufacturer's instructions (Applied Biosystems, Warrington, U.K., <http://www.appliedbiosystems.com>) [26]. Primers for PPAR γ , C/EBP α , and FABP4 were bought as Assay on Demand kits from Applied Biosystems and amplified using the 7900HT Real-Time PCR machine, analyzed using SDS 2.2 (Applied Biosystems) and expressed as $2^{-\Delta C_T}$, where C_T represents the difference in the cycle number between the gene of interest and 18S (Supporting Information Fig 1B–1D).

Differentiation of Adipose Derived Preadipocytes into Adipocytes

ADSC were (5×10^5) were seeded onto inverted 0.4 μm pore Transwell filter inserts (BD Biosciences, Cowley, U.K., <http://www.bdbiosciences.com/eu/home>) as previously described [4, 5] and cultured in AD basal media for 24 hours. Adipogenic differentiation was induced by culturing cells in DMEM/F-12 supplemented with 166 nM recombinant insulin, 0.2 nM triiodothyronine (T3), 33 μM Biotin, 17 μM pantothenic acid, 0.01 mg/ml transferrin, and 100 nM cortisol (all from Sigma) for 21–28 days as previously described [24]. For the first 5 days, media was also supplemented with 50 $\mu\text{g}/\text{ml}$ 3-isobutyl-1-methylxanthine and 2 μM Rosiglitazone (PPAR agonist) (all from Sigma).

Isolation and Culture of EC

Human umbilical vein endothelial cells (HUVEC) were isolated from umbilical cords as previously described [4, 27] and cultured in Medium 199 (Life Technologies, Paisley, U.K., <http://www.lifetech.com>) supplemented with 20% FCS, 35 $\mu\text{g}/\text{ml}$ gentamicin, 10 ng/ml epidermal growth factor, 1 $\mu\text{g}/\text{ml}$ hydrocortisone (all from Sigma), and 2.5 $\mu\text{g}/\text{ml}$ Amphotericin B (Life Technologies).

Isolation of Neutrophils and Lymphocytes

Venous blood was collected from healthy donors into EDTA tubes (Sarstedt, Leicester, U.K., <https://www.sarstedt.com/en/home/>). Neutrophils (PMN) or peripheral blood mononuclear cells (PBMC) were isolated by centrifugation on two-step histopaque density gradients as previously described [5, 21]. PBMC were panned on culture plastic for 30 minutes to remove monocytes and purify peripheral blood lymphocytes (PBL). Purified leukocytes were washed twice in PBS containing 1 mM Ca^{2+} , 0.5 mM Mg^{2+} , and 0.15% bovine serum albumin (PBSA; all from Sigma) at 250g for 5 minutes, counted and resuspended to 2×10^6 cells per ml in PBSA.

Establishing Cocultures of EC with MSC or Adipocytes

Cocultures were established by seeding EC and stromal cells (either MSC or MSC-derived adipocytes) on opposite sides of 0.4 μm pore Transwell filter inserts as previously described [4, 5]. Briefly, MSC, MSC-derived adipocytes, or ADSC (5×10^5) were seeded onto inverted filters and cultured for 24 hours. HUVEC were then seeded on the inner surface of inserts. Cells were cocultured for 24 hours prior to treatment with or without 100 U/ml tumour necrosis factor- α (TNF α ; R&D Systems, Abingdon, U.K., <http://www.rndsystems.com>) for a further 4 hours (neutrophils) or TNF α in combination with 10 ng/ml interferon- γ (IFN γ ; Peprotech Inc., London, U.K., <http://www.peprotech.com>) for a further 24 hours (PBL) [3, 5, 25]. Parallel endothelial monocultures were set up as controls.

In a separate series of experiments, nonadherent mAD (5×10^5 cells) were suspended in a well, on top of which a filter was placed (such that the mAD came into contact with the basal surface of the filter). Cells were cultured 24 hours prior to seeding EC into the filter as above. Cocultures were established for 24 hours and cytokine treated as above.

In some experiments, a neutralizing antibody against IL-6 (5 $\mu\text{g}/\text{ml}$; clone 6708; R&D Systems) was added when

cocultures were established and was present throughout the coculture and cytokine stimulation.

To investigate the bioactivity of conditioned media, supernatants were obtained from MSC and MSC-derived adipocyte monoculture and cocultures at 24 hours. Fresh EC monocultures were treated with conditioned media for 24 hours prior to stimulation with TNF α in the same conditioned media for a further 4 hours [5].

Cocultures were also formed with collagen gels containing MSC, and EC cultured on top. Rat-tail collagen type 1 (2.15 mg/ml; First Link Ltd., West Midlands, U.K., <http://www.firstlinkuk.co.uk/>) was mixed with $10 \times$ M199 and then neutralized by addition of 1 N NaOH on ice, as described [21, 28, 29]. MSC (2.5×10^4 cells per well in a 12-well plate) were added to 500 μl , and the gel was allowed to set for 15 minutes at 37°C, and then equilibrated for 24 hours. HUVEC were seeded on the surface of the gel and cocultured with MSC for 24 hours, prior to cytokine treatment as above.

Flow-Based Adhesion Assay

Flow-based adhesion assays were performed for filters incorporated into a custom-made parallel-plate flow chamber, using phase-contrast digital microscopy as previously described [3, 4]. Purified leukocytes were perfused over EC for 4 minutes followed by washout with cell-free PBSA, all at a wall shear stress of 0.1 Pa. Digitized recordings of 5–10 random fields were made 2 and 9 minutes after the end of the leukocyte bolus to assess leukocyte adhesion and transmigration, respectively. Images were analyzed offline using Image-Pro Plus software (Media Cybernetics, Marlow, U.K., <http://www.mediacy.com/>). Total leukocytes bound to the endothelium were counted and classified as either: (a) rolling adherent; (b) stationary, or (c) transmigrated under the endothelial monolayer. The total number of bound cells at 2 minutes was averaged per field and expressed as cells/ mm^2 per 10^6 perfused; at 9 minutes the proportion of the bound cells that had transmigrated was evaluated.

Collagen Gel-Based Static Adhesion Assay

Neutrophils (5×10^5 cells per ml) were added to the EC for 20 minutes, and nonadherent cells were removed by washing three times in PBSA. Neutrophil adhesion and migration was assessed using phase contrast digital microscopy by acquiring digitised z-stack images from five random fields on the surface of the endothelium throughout the depth of the gel at 2 hours, as described [29]. Images were analyzed offline using Image-Pro Plus software. The number of neutrophils, adherent on the surface of the endothelium and throughout the gel, was averaged per field. From the known areas of the field and well, these counts were converted to totals per gel and expressed as a percentage of the number of cells added. The proportion of the adherent neutrophils migrated into the gel was also calculated.

Gene Expression Analysis

Gene expression was analyzed using mRNA isolated from EC, MSC, and MSC-derived adipocyte monocultures, or cocultures as described previously [5]. Data were calculated as $2^{-\Delta\text{CT}}$ relative to expression of 18S, as described above.

Quantification of Soluble Mediators

Culture supernatants were obtained from unstimulated EC and MSC cultured alone or in coculture for 24 hours. IL-6 and soluble IL-6 receptor (sIL-6R) were quantified using IL-6 Duo-Set ELISA and sIL-6R Quantikine ELISA Kit, respectively (R&D Systems) according to manufacturer's instructions.

Ethics

The study was conducted in compliance with the Declaration of Helsinki. All human samples were obtained with written, informed consent and approval from the Human Biomaterial Resource Centre (Birmingham, U.K.), West Midlands and Black Country Research Ethics Committee, North East—Tyne and West South Research Ethics Committee, or University of Birmingham Local Ethical Review Committee.

Statistical Analysis

Data are expressed as mean \pm SEM, where different EC and leukocyte donors were used for each experiment. Three to ten different MSC and three primary ADSC donors were incorporated in each experiment for each independent experiment. Multivariate data were analyzed using analysis of variance (ANOVA), followed by Tukey or Dunnett post hoc tests for comparisons between treatments or to controls respectively. Univariate data were analyzed using paired *t* test. $p \leq 0.05$ was considered statistically significant.

RESULTS

Effect of Adipogenic Differentiation on the Ability of MSC to Suppress Leukocyte Recruitment by EC

MSC differentiation into adipocytes was confirmed by the presence of prominent lipid droplets and the significant upregulation of the adipocyte-related genes (PPAR γ , C/EBP α , and FABP4) compared to undifferentiated MSC (Supporting Information Fig. 1). In coculture with EC, BMMSC suppressed neutrophil adhesion to TNF α -stimulated EC under flow conditions (Fig. 1A) and in a static 3D tissue model (Fig. 1B). This effect was lost when BMMSC-derived adipocytes were incorporated into coculture, such that they no longer inhibited neutrophil recruitment from flow (Fig. 1A) or adhesion in the 3D model (Fig. 1B). Similar effects were observed when trabecular bone or umbilical cord MSC, or MSC-derived adipocytes were cocultured with EC (Supporting Information Fig. 2; also see Supporting information for derivation of MSC). Lymphocyte recruitment to TNF α and IFN γ -stimulated EC was suppressed by BMMSC, but in this case, MSC-derived adipocytes continued to suppress recruitment when incorporated into the cocultures (Fig. 1C). Adipogenic differentiation of MSC had no significant effect on the proportion of adherent leukocytes migrating through the endothelium when compared to the undifferentiated MSC cocultures or the EC cultured alone (data not shown). Thus, MSC lost their immunosuppressive effects on neutrophil, but not lymphocyte, recruitment upon differentiation into adipocytes. These responses were shared by MSC from several tissues. All subsequent experiments were performed using MSC isolated from BM.




Figure 1. Effects of differentiation of MSC into adipocytes on their immunomodulation of neutrophil or lymphocyte recruitment. Cocultures formed by seeding EC with bone marrow MSC (BMMSC) or BMMSC-derived adipocytes (**A, C**) on opposite sides of 0.4 μ m porous filters or (**B**) seeding EC onto a collagen gel containing BMMSC or BMMSC-derived adipocytes for 24 hours. Cultures were stimulated with (**A, B**) 100 U/ml tumor necrosis factor α (TNF α) for 4 hours or (**C**) 100 U/ml TNF α and 10 ng/ml interferon γ for 24 hours. In (**A**) and (**C**), a 4 minute bolus of purified leukocytes was perfused over the endothelium. (**A**) Neutrophil or (**C**) PBL adhesion was assessed and expressed as the number of cells adherent/mm² per 10⁶ cells perfused. In (**B**), neutrophils were allowed to adhere to the EC cultured on collagen gel containing MSC or adipocytes for 20 minutes before adhesion was assessed at 2 hours and expressed as a percentage of cells added. Data are mean \pm SEM from $n = 3$ experiments incorporating a different EC and leukocyte donor in each experiment and two different BMMSC donors. ANOVA showed a significant effect of culture conditions on neutrophil ($p < 0.01$ in **A**, $p < 0.05$ in **B**) and PBL adhesion ($p < 0.01$ in **C**). *, $p < 0.05$; **, $p < 0.01$ by Tukey post-test. Abbreviations: EC, endothelial cells; MSC, mesenchymal stem cells; PBL, peripheral blood lymphocytes; PMN, neutrophil.




Figure 2. Secretion and role in immunosuppression of IL-6 in cocultures of MSC or MSC-derived adipocytes. **(A)** IL-6 and **(B)** sIL-6R release into supernatants from EC, bone marrow MSC (BMMSC), and BMMSC-derived adipocyte monoculture and cocultures was assessed after 24 hours. **(C)** BMMSC or **(D)** BMMSC-derived adipocyte cocultures were treated with neutralizing antibodies against IL-6 for the duration of the coculture and cytokine treatment. Neutrophil adhesion was assessed and expressed as the number of cells adherent/mm² per 10⁶ cells perfused. **(E)** SOCS3 gene expression analyzed by quantitative polymerase chain reaction in EC cultured alone, with MSC or with MSC-derived adipocytes. Data are expressed as 2^{-ΔCt} relative to SOCS3 expression by EC cultured alone. Data are mean ± SEM (A) *n* = 16–19 and (B, C) *n* = 4, (D) *n* = 3, or (E) *n* = 3 independent experiments using a different EC and (C, D) neutrophil donor in each experiment. Three different BMMSC donors were used in all experiments, except (A) where 10 different donors were used. ANOVA showed significant effects of treatment in (A) (*p* < 0.01), (C and D) (each *p* < 0.05) *, *p* < 0.05 and **, *p* < 0.01 by Tukey post-test. ¥¥ = *p* < 0.01 compared to the sum of the EC and respective MSC monoculture supernatant by paired *t* test. Abbreviations: AD, adipocytes; EC, endothelial cells; IL-6, interleukin-6; MSC, mesenchymal stem cells; sIL-6R, soluble IL-6 receptor; SOCS3, suppressor of cytokine signaling 3.

Role of IL-6 in Effects of MSC and MSC-Derived Adipocytes in Coculture

We have previously identified IL-6 as the bioactive agent required for the inhibitory effects of MSC in coculture with EC [3, 5]. Therefore, we compared the presence and activity of IL-6 released in different cocultures, hypothesizing that they might account for the differences in effects of MSC and MSC-derived adipocytes on neutrophil adhesion. MSC-derived adipocyte monocultures released significantly more IL-6 compared to undifferentiated MSC monocultures (Fig. 2A). MSC cocultures released a higher concentration of IL-6 than MSC monocultures, but production of IL-6 did not increase when

MSC-derived adipocytes were cocultured with EC (Fig. 2A). As a result, the two types of coculture generated similar levels of IL-6. sIL-6R levels were also similar between all culture conditions tested (Fig. 2B). Thus, loss of MSC-mediated suppression upon adipogenic differentiation was not due to a decrease in the release of IL-6 or its receptor in MSC-derived adipocyte cocultures. This raised the question of whether IL-6 present in the coculture secretome remained bioactive.

In previous studies comparing stromal cells from healthy and diseased tissues, we have reported that similar concentrations of IL-6 produced during coculture had opposing effects (immunosuppression versus stimulation) on leukocyte

recruitment [21, 30]. Here we hypothesized that the differentiation of MSC into adipocytes may alter the bioactivity of IL-6 during coculture. Neutralization of IL-6 significantly reduced the inhibitory effects of MSC in coculture (Fig. 2C). In contrast, neutralization of IL-6 in MSC-derived adipocyte cocultures significantly reduced neutrophil adhesion (Fig. 2D), restoring immunoprotective function. Thus IL-6 had essentially opposite effects in cocultures with either MSC or MSC-derived adipocytes; supporting the concept that conversion of MSC to adipocytes resembled a pathological process.

Suppressor of cytokine signaling 3 (SOCS3) is an IL-6 inducible gene known to regulate TNF α responses [reviewed in ref. 31]. Here, expression of SOCS3 in endothelial cells was upregulated by coculture with MSC, but not MSC-derived adipocytes, when compared to the EC monoculture controls (Fig. 2E). Thus, downstream signaling from IL-6/sIL-6R differed between the two forms of coculture.

Role of TGF β 1 Signalling in Variations in Immunosuppression

We have previously reported a role for TGF β ₁ in mediating the immunosuppressive actions of MSC in coculture [3]. We wondered whether MSC differentiation might alter TGF β ₁ responses in our cocultures. Following coculture with MSC, EC expressed significantly higher levels of TGF β -R₁ and TGF β -R₃ at the transcript level compared to the EC monoculture controls (Fig. 3). In contrast, MSC-derived adipocytes had no effect on the expression of these TGF β receptors by EC (Fig. 3). Thus loss of TGF β ₁ downstream signaling may contribute to the changes in IL-6 signaling and loss of immunosuppression in MSC-derived adipocyte cocultures.

Profile and Activity of Secretome Released by MSC and MSC-Derived Adipocytes in Coculture

Next we examined whether differentiation into adipocytes altered the secretome generated during coculture, such that it was no longer immunosuppressive. In an unbiased analysis using a cytokine expression array (see Supporting Information for methods), we detected higher levels of some analytes (e.g., angiogenin, CD105, and platelet derived growth factor (PDGF)-AA) and lower levels of others (e.g., Dickkopf-related protein 1, CXCL12, vascular endothelial growth factor (VEGF)) in the MSC-derived adipocyte coculture supernatants when compared to the MSC cocultures (Fig. 4A; Supporting Information Fig 3). In some cases, the expression of analytes, such as adiponectin, resistin, or IL-6, was comparable between both coculture conditions (Supporting Information Fig 3). We thus tested whether the MSC-derived adipocyte coculture secretome had different bioactivity compared to MSC in our recruitment assays. MSC coculture conditioned media mimicked the effect of MSC-EC coculture, inhibiting neutrophil adhesion to TNF α -treated EC monocultures *in vitro* (Fig. 4B), as previously described [5]. In contrast, conditioned media from MSC-derived adipocyte monoculture or coculture had no effect on neutrophil adhesion (Fig. 4C). This indicates that differentiation of MSC into adipocytes alters both the composition and bioactivity of the secretome generated during coculture. Thus raises the possibility that other bioactive soluble agents generated by MSC-derived adipocyte cocultures are capable of modulating IL-6 and TGF β ₁ responses, such that they are no longer immunoprotective.




Figure 3. Expression and function of TGF β receptors in endothelial cells cocultured with MSC or MSC-derived adipocytes. Cocultures formed by seeding EC with MSC or MSC-derived adipocytes on opposite sides of 0.4 μ m porous filters for 24 hours. Gene expression for (A) TGF β R1, (B) TGF β R3, and (C) decorin was analyzed in endothelial cells cultured alone or with either MSC or MSC-derived adipocytes by qPCR. Data are expressed as 2^{- Δ CT} relative to 18S. Data are mean \pm SEM from (A) $n = 5-6$, (B) $n = 3$ or (C) $n = 4$ independent experiments using a different donor for each cell type in each experiment. ANOVA showed significant effects of treatment in (A) and (B) (both $p < 0.01$). *, $p < 0.05$ and **, $p < 0.01$ by Tukey post-test. Abbreviations: EC, endothelial cells; MSC, mesenchymal stem cells; TGF β , transforming growth factor beta.

Immunosuppressive Effects of Adipocytes Derived from Adipose Tissue

Finally we analyzed whether stromal cells from "true" adipose tissue (ADSC), adipocytes derived from these cells (ADSC-derived adipocytes), and mature adipocytes (mAD) extracted




Figure 4. Content and immunosuppressive effects of conditioned media from MSC or adipocyte cocultures. **(A)** Conditioned media from EC alone or EC in coculture were analyzed by a cytokine expression array. Representative immunoblots from a single experiment. EC monocultures were treated with conditioned media from **(B)** BMMSC or **(C)** BMMSC-derived adipocytes either cultured alone (monocultures) or cocultured with EC for 24 hours. Neutrophil adhesion was assessed at 2 minutes post perfusion and expressed as the number of cells adherent/mm² per 10⁶ cells perfused. In **(B)**, ANOVA showed a significant effect of treatment on neutrophil adhesion, $p < 0.05$. Data are mean \pm SEM, **(B)** $n = 3$ –5 and **(C)** $n = 5$ independent experiments using a different EC and neutrophil donor in each experiment. **(B)** 2 or **(C)** 3 different BMMSC donors were used. *, $p < 0.05$ by Dunnett post-test. Abbreviations: AD, adipocytes; EC, endothelial cells; MSC, mesenchymal stem cells.

from the tissue exerted the same effects as BMMSC-derived adipocytes on leukocyte recruitment to EC. Coculture with ADSC significantly reduced neutrophil recruitment to inflamed EC (Fig. 5A), sharing the immunosuppressive capabilities seen in MSC from other tissue sources [3, 5]. Unlike BMMSC-derived adipocytes, both ADSC-derived adipocytes and mAD

were immunosuppressive in coculture, significantly inhibiting neutrophil adhesion to TNF α -stimulated EC (Fig. 5B, 5C). Suppression of adhesion was also evident when lymphocytes were perfused over these cocultures (Fig. 6). We also detected high levels of IL-6 in ADSC and mAD monocultures, which did not increase when these cells were cocultured with EC (Supporting Information Fig. 4). However, in contrast to BMMSC-derived adipocytes, neutralization of IL-6 significantly inhibited the immunosuppressive effects of ADSC in coculture (Fig. 5D). Thus adipocytes derived from adipose tissue stromal cells and “native” adipocytes maintained immunoprotective effects, while adipose cells derived from BMMSC lost this capability. It appears, therefore, that ectopic MSC-derived adipose cells would lack the ability of ‘true’ adipose tissue derived stromal cells to regulate leukocyte recruitment.

DISCUSSION

Surprisingly little is known about how differentiation of MSC alters their immunomodulatory properties, with much of the work focusing on osteogenesis and chondrogenesis producing conflicting findings [reviewed in ref. 6]. We examined for the first time the effects of adipogenesis on MSC crosstalk with EC and regulation of the inflammatory infiltrate. We also compared this form of immunosuppression between adipocytes generated from adipose tissue, and adipocytes derived from MSC from other tissues. Interestingly, they were not the same. Upon coculture with EC, BMMSC-derived adipocytes had lost the ability of BMMSC to suppress neutrophil adhesion. The ability to suppress lymphocyte adhesion was retained. This phenotype was shared with adipocytes differentiated from MSC derived from other non-adipose tissue. In contrast, adipocytes differentiated from adipose-derived stromal cells and mature adipocytes retained immunoprotective capacity, suppressing both neutrophil and lymphocyte recruitment. Thus the effect of adipogenesis on immunoregulation appears to be tissue specific, with stromal cells from adipose and nonadipose tissues generating cells with different capabilities.

A key attribute of MSC is their capacity to differentiate into various cell lineages to aid the tissue repair processes. However, we understand little about how this process affects MSC behavior, or if the context in which differentiation is triggered can alter the outcome. Here, MSC isolated from non-adipose tissue lost some of their immunoprotective effects following adipogenesis, such that they were no longer able to inhibit neutrophil adhesion. This characteristic was not shared by ADSC, which retained their immunomodulatory capacity even after differentiation. Of note, the latter observation argues against the loss of suppression upon adipogenesis by non-adipose-derived MSC being an artefact of *in vitro* differentiation, as *in vitro* differentiation of ADSC had no effect on their suppressive potential.

IL-6 is known to have different functions, eliciting immunoprotective responses or proinflammatory effects, depending on the inflammatory and stromal context [reviewed in ref. 32]. Indeed here and in our previous studies [3, 5], IL-6 was identified as a bioactive agent required for the inhibitory effects of MSC in coculture. However, neutralization of IL-6 reversed the effects of MSC-derived adipocytes in coculture




Figure 5. Immunosuppressive effects by ADSC and adipocytes on neutrophil recruitment. EC were cocultured with (A) ADSC, (B) ADSC-derived adipocytes (ADSC adipo) or (C) mature adipocytes (Mature adipo) for 24 hours prior to stimulation with tumor necrosis factor α (TNF α) for 4 hours. (D) ADSC cocultures were treated with neutralizing antibodies against IL-6 for the duration of the coculture and cytokine treatment. Neutrophil adhesion was assessed at 2 minutes postperfusion and expressed as the number of cells adherent/mm² per 10⁶ cells perfused. In (D), ANOVA showed a significant effect of treatment on neutrophil adhesion, $p < 0.05$. Data are mean \pm SEM, (A) $n = 6$, (B) $n = 3$, and (C, D) $n = 4$ independent experiments using a different EC and neutrophil donor in each experiment. Three different stromal cell donors were used in all experiments, except (D) where one donor was used. *, $p < 0.05$ compared to EC monoculture by paired t test. ¥, $p < 0.05$ and ¥¥, $p < 0.01$ by Tukey post-test. Abbreviations: ADSC, adipose-derived stromal cells; EC, endothelial cells; IL, interleukin.

to be inhibitory, suggesting that under these conditions, IL-6 had not just lost efficacy but triggered stimulatory rather than inhibitory signals. These findings are akin to those reported for synovial fibroblasts from patients with rheumatoid arthritis in coculture with EC, where IL-6 signaling stimulated recruitment of leukocytes [20, 21]. Collectively these data support the concept that conversion of MSC to adipocytes resembles a pathological process. Here, we found that in MSC cocultures, IL-6 induced the expression of SOCS3 in EC, a typically IL-6-regulated gene known to downregulate cellular responses to cytokines [31], and which we previously found to play a role in modification of neutrophil recruitment by viral infection [33]. Interestingly, we detected no changes in EC SOCS3 expression following coculture with MSC-derived adipocytes, despite the presence of IL-6 at similar concentrations as those detected in MSC cocultures. Collectively these data reveal two distinct IL-6 signalling pathways, which are induced in a context-specific manner and elicit different functional consequences in EC.

To the best of our knowledge only two studies have assessed MSC following adipogenesis, and neither observed any alterations in their ability to inhibit T-cell proliferation [9] or dendritic cell maturation [12]. Similarly, suppression of T-cell proliferation remained unaffected by osteogenic differentiation of human BMMSC [7–9] or ADSC [8]. On the other hand, MSC-derived chondrocytes have been reported to

inhibit [9, 10] or have no effect on [11, 12] on T-cell responses. In the latter studies, reduction in bioactive soluble mediators, such as nitric oxide or prostaglandin E₂, were linked to the defect in suppressive activity of MSC after differentiation [11]. Here we also observed a different profile of soluble mediators released by MSC and differentiated MSC in coculture. Moreover, conditioned medium from MSC-derived adipocyte cocultures was unable to suppress neutrophil recruitment in the same manner as medium generated from MSC cocultures. This clearly shows that the secretomes generated by crosstalk between EC and MSC or MSC-derived adipocytes have different bioactivity. Thus we propose that the changes in the bioactivity of IL-6 in MSC-derived adipocyte cocultures occurs due to the presence of additional soluble mediators (exclusive to these cocultures) that cause a reduction in TGF β receptors and thus response to TGF β ₁, along with a failure to induce the suppressor genes SOCS3, usually associated with IL-6 stimulation. Further work is required to identify the exact soluble mediator(s) responsible for these changes, which have the potential to be novel therapeutic targets for diseases involving ectopic fat deposition.

Healthy adipose tissue is considered to be intrinsically immunoprotective, largely through the abundant production of the immunosuppressive adipokine, adiponectin. Indeed our data would agree with this, with stromal cells at all stages of lineage commitment (preadipocytes, differentiated adipocytes,




Figure 6. Immunosuppressive effects of ADSC and adipocytes on lymphocyte recruitment. EC were cocultured with (A) ADSC, (B) ADSC-derived adipocytes, or (C) mature adipocytes (mature adipo) for 24 hours prior to stimulation with TNF α and IFN γ for 24 hours. PBL adhesion was assessed at 2 minutes postperfusion and expressed as the number of cells adherent/mm² per 10⁶ cells perfused. Data are mean \pm SEM from $n = 3-4$ independent experiments using a different donor for each cell type in each experiment. *, $p < 0.05$; **, $p < 0.01$ compared to EC monoculture by paired t test. Abbreviations: ADSC, adipose-derived stromal cells; EC, endothelial cells; PBL, peripheral blood lymphocytes.

mAD) exhibiting immunosuppressive effects in our assays. Moreover, adipose-derived MSC are known to inhibit T-cell proliferation [reviewed in ref. 34] and appear to retain this capacity after differentiation into osteoblasts [7, 8] or adipocytes [35]. Thus adipocytes, much like stromal cells from other healthy tissues [5, 21, 36], have a homeostatic role in limiting

inflammatory leukocyte infiltration. Indeed, IL-6 is the shared common bioactive immunoprotective mediator of healthy stromal cells [5, 21, 36] and was responsible for the immunoprotective capacity of adipose-derived stromal cells in coculture. Here, MSC-derived adipocytes, ADSC, and mAD all secreted high levels of IL-6 in culture, which were not altered by coculture with EC. Given the different effects of these cocultures, clearly it is not simply the concentration of IL-6 but also the downstream signalling pathway(s) it elicits which are critical for its anti-inflammatory effect. Further work is required to determine the mechanisms by which adipocytes from adipose tissue mediate their immunomodulatory effects and if these effects become modified in obesity and other metabolic disorders.

Adipogenesis in non-adipose tissue is unlikely to be a homeostatic response, but rather the inappropriate response of the tissue to repeated episodes of damage and insult associated with chronic inflammatory diseases. Therefore, it may not be surprising that the endogenous immunoprotective effect of the tissue stroma may become distorted. Supporting this concept, neutrophil infiltration into intra-abdominal adipose tissue was increased in mice on a high-fat diet compared to those on a normal chow diet [37]. Thus adipocytes may have the potential to exist in at least two functional states: (a) immunosuppressive as found in healthy adipose tissue and (b) stimulatory in sites of abnormal (obesity) or ectopic (chronic inflammation) fat deposition. Whether the adipocytes from the latter two sites are phenotypically and functionally the same remains to be determined. Our work also suggests an unexplored scenario; that changes in the phenotype of MSC at sites of chronic inflammation may contribute to uncontrolled leukocyte infiltration and pathogenesis.

CONCLUSION

In their natural environment MSC act to endogenously dampen EC responses to cytokines, thereby limiting leukocyte recruitment in an IL-6 dependent manner. However, aberrant differentiation of non-adipose-derived MSC causes the cells to lose this intrinsic immunoprotective capacity. Alterations in the phenotype and function of MSC may take the brake off and contribute to pathogenic inflammatory responses by altering the way in which IL-6 and TGF β_1 exerts their effects, switching from anti-inflammatory to pro-inflammatory. This does not appear to be a characteristic of adipose tissue per se but may occur when MSC differentiate in an inappropriate location or circumstance.

ACKNOWLEDGMENTS

H.M. was supported by BBSRC Ph.D. studentship, while L.S.C.W. was supported by MRC Ph.D. studentship, all awarded to University of Birmingham. H.M.M. and F.B. were supported by an Arthritis Research UK Career Development Fellowship (19899) and an Arthritis Research UK Senior Fellowship (21236), respectively. Studies were supported by grants from the British Heart Foundation (PG/14/28/30774) and Pfizer (PO:8500669699). Arthritis Research UK Rheumatoid Arthritis Pathogenesis Centre of Excellence (RACE) was part-funded by Arthritis Research UK (20298); this Centre is a collaboration between the Universities of Glasgow, Newcastle, and

Birmingham. Umbilical cords were collected with the assistance of the research midwives at Birmingham Women's Health Care NHS Trust and Sandwell and West Birmingham Hospitals NHS Trust.

AUTHOR CONTRIBUTIONS

H.M., L.W., L.S., and S.K.: collection and/or assembly of data, data analysis and interpretation, manuscript writing, and final approval of manuscript; S.N. and F.B.: provision of study material or patients, data analysis and interpretation,

manuscript writing, and final approval of manuscript; G.N. and H.M.: conception and design, financial support, collection and/or assembly of data, data analysis and interpretation, manuscript writing, and final approval of manuscript.

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

H.M.M. has received research funding from Pfizer. All other authors declare that they have no conflicts of interest.

REFERENCES

- Munir H, McGettrick HM. Mesenchymal stem cells therapy for autoimmune disease: Risks and rewards. *Stem Cells Dev* 2015;24:2091–3100.
- Le Blanc K, Mougiakakos D. Multipotent mesenchymal stromal cells and the innate immune system. *Nat Rev Immunol* 2012;12:383–396.
- Luu N-T, McGettrick HM, Buckley CD et al. Crosstalk between mesenchymal stem cells and endothelial cells leads to downregulation of cytokine-induced leukocyte recruitment. *Stem Cells* 2013;31:2690–2702.
- Munir H, Rainger GE, Nash GB et al. Analyzing the effects of stromal cells on the recruitment of leukocytes from flow. *J Vis Exp* 2015;95:e52480.
- Munir H, Luu NT, Clarke LSC et al. Comparative ability of mesenchymal stromal cells from different tissues to limit neutrophil recruitment to inflamed endothelium. *Plos One* 2016;11:e0155161.
- Lohan P, Coleman CM, Murphy JM et al. Changes in immunological profile of allogeneic mesenchymal stem cells after differentiation: Should we be concerned?. *Stem Cell Res Ther* 2014;5:1–11.
- Montespan F, Deschaseaux F, Sensebe L et al. Osteodifferentiated mesenchymal stem cells from bone marrow and adipose tissue express HLA-G and display immunomodulatory properties in HLA-mismatched settings: Implications in bone repair therapy. *J Immunol Res* 2014;2014:230346.
- Niemeyer P, Kornacker M, Mehlhorn A et al. Comparison of immunological properties of bone marrow stromal cells and adipose tissue-derived stem cells before and after osteogenic differentiation in vitro. *Tissue Eng* 2007;13:111–121.
- Le Blanc K, Tammik C, Rosendahl K et al. HLA expression and immunologic properties of differentiated and undifferentiated mesenchymal stem cells. *Exp Hematol* 2003;31:890–896.
- Zheng ZH, Li XY, Ding J et al. Allogeneic mesenchymal stem cell and mesenchymal stem cell-differentiated chondrocyte suppress the responses of type II collagen-reactive T cells in rheumatoid arthritis. *Rheumatology* 2008;47:22–30.
- Ryan AE, Lohan P, Flynn L et al. Chondrogenic differentiation increases antidonor immune response to allogeneic mesenchymal stem cell transplantation. *Mol Ther* 2014;22:655–667.
- Chen X, McClurg A, Zhou GQ et al. Chondrogenic differentiation alters the immunosuppressive property of bone marrow-derived mesenchymal stem cells, and the effect is partially due to the upregulated expression of B7 molecules. *Stem Cells* 2007;25:364–370.
- Uezumi A, Fukada S, Yamamoto N et al. Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. *Nat Cell Biol* 2010;12:143–152.
- Goldfarb JW, Roth M, Han J. Myocardial fat deposition after left ventricular myocardial infarction: Assessment by using MR water-fat separation imaging. *Radiology* 2009;253:65–73.
- Goodpaster BH, Wolf D. Skeletal muscle lipid accumulation in obesity, insulin resistance, and type 2 diabetes. *Pediatric Diabetes* 2004;5:219–226.
- Kastrinaki MC, Sidiropoulos P, Roche S et al. Functional, molecular and proteomic characterisation of bone marrow mesenchymal stem cells in rheumatoid arthritis. *Ann Rheum Dis* 2008;67:741–749.
- Nie Y, Lau CS, Lie AKW et al. Defective phenotype of mesenchymal stem cells in patients with systemic lupus erythematosus. *Lupus* 2010;19:850–859.
- Sun L, Wang D, Liang J et al. Umbilical cord mesenchymal stem cell transplantation in severe and refractory systemic lupus erythematosus. *Arthritis & Rheumatism* 2010;62:2467–2475.
- Tang Y, Xie H, Chen J et al. Activated NF-KB in bone marrow mesenchymal stem cells from systemic lupus erythematosus patients inhibits osteogenic differentiation through downregulating smad signaling. *Stem Cells Dev* 2012;22:668–678.
- Lally F, Smith E, Filer A et al. A novel mechanism of neutrophil recruitment in a coculture model of the rheumatoid synovium. *Arthritis Rheum* 2005;52:3460–3649.
- McGettrick HM, Smith E, Filer A et al. Fibroblasts from different sites may promote or inhibit recruitment of flowing lymphocytes by endothelial cells. *Eur J Immunol* 2009;39:98–107.
- Rainger GE, Nash GB. Cellular pathology of atherosclerosis: Smooth muscle cells prime cocultured endothelial cells for enhanced leukocyte adhesion. *Circ Res* 2001;88:615–622.
- Tull SP, Anderson SI, Hughan SC et al. Cellular pathology of atherosclerosis: Smooth muscle cells promote adhesion of platelets to cocultured endothelial cells. *Circ Res* 2006;98:98–104.
- McNelis JC, Manolopoulos KN, Gathercole LL et al. Dehydroepiandrosterone exerts antiglycocorticoid action on human preadipocyte proliferation, differentiation, and glucose uptake. *Am J Physiol-Endocrinol Metab* 2013;305:E1134.
- McGettrick HM, Buckley CD, Filer A et al. Stromal cells differentially regulate neutrophil and lymphocyte recruitment through the endothelium. *Immunology* 2010;131:357–370.
- Chimen M, McGettrick HM, Apta B et al. Homeostatic regulation of T cell trafficking by a B cell-derived peptide is impaired in autoimmune and chronic inflammatory disease. *Nat Med* 2015;21:467–475.
- Cooke BM, Usami S, Perry I et al. A simplified method for culture of endothelial cells and analysis of adhesion of blood cells under conditions of flow. *Microvasc Res* 1993;45:33–45.
- Jeffery HC, Buckley CD, Moss P et al. Analysis of the effects of stromal cells on the migration of lymphocytes into and through inflamed tissue using 3-D culture models. *J Immunol Methods* 2013;400-401:45–57.
- Luo D, McGettrick HM, Stone PC et al. The roles of integrins in function of human neutrophils after their migration through endothelium into interstitial matrix. *Plos One* 2015;10:e0118593.
- McGettrick HM, Filer A, Raza K et al. Fibroblasts lose their immunosuppressive ability early in the development of rheumatoid arthritis: Effects on lymphocyte recruitment. *Ann Rheum Dis* 2014;73:A19.
- Alexander WS, Hilton DJ. The role of suppressors of cytokine signaling (SOCS) proteins in regulation of the immune response. *Ann Rev Immunol* 2004;22:503–529.
- McGettrick HM, Butler LM, Buckley CD et al. Tissue stroma as a regulator of leukocyte recruitment in inflammation. *J Leukocyte Biol* 2012;91:385–400.
- Butler LM, Jeffery HC, Wheat RL et al. Kaposi's sarcoma-associated Herpesvirus inhibits expression and function of endothelial cell major histocompatibility complex class II via suppressor of cytokine signaling 3. *J Virol* 2012;86:7158–7166.
- Zhao Q, Ren H, Han Z. Mesenchymal stem cells: Immunomodulatory capability and clinical potential in immune diseases. *J Cell Immunother* 2016;2:3–20.
- Hashemi SM, Hassan ZM, Pourfathollah AA et al. In vitro immunomodulatory

properties of osteogenic and adipogenic differentiated mesenchymal stem cells isolated from three inbred mouse strains. *Biotechnol Lett* 2013;35:135–142.

36 Kuravi SJ, McGettrick HM, Satchell SC et al. Podocytes regulate neutrophil recruitment by glomerular endothelial cells via IL-6 mediated crosstalk. *J Immunol* 2014;193:234–243.

37 Elgazar-Carmon V, Rudich A, Hadad N et al. Neutrophils transiently infiltrate intra-abdominal fat early in the course of high-fat feeding. *J Lipid Res* 2008;49:1894–1903.



See www.StemCellsTM.com for supporting information available online.