

Concentrations of polychlorinated biphenyls in soil and indoor dust associated with electricity generation facilities in Lagos, Nigeria

Folarin, Bilikis Temitope; Abdallah, Mohamed Abou-Elwafa; Oluseyi, Temilola; Olayinka, Kehinde; Harrad, Stuart

DOI:

[10.1016/j.chemosphere.2018.05.110](https://doi.org/10.1016/j.chemosphere.2018.05.110)

License:

Other (please provide link to licence statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Folarin, BT, Abdallah, MA-E, Oluseyi, T, Olayinka, K & Harrad, S 2018, 'Concentrations of polychlorinated biphenyls in soil and indoor dust associated with electricity generation facilities in Lagos, Nigeria', *Chemosphere*, vol. 207, pp. 620-625. <https://doi.org/10.1016/j.chemosphere.2018.05.110>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Published in *Chemosphere* on 19/05/2018

DOI: 10.1016/j.chemosphere.2018.05.110

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

**CONCENTRATIONS OF POLYCHLORINATED BIPHENYLS IN SOIL
AND INDOOR DUST ASSOCIATED WITH ELECTRICITY
GENERATION FACILITIES IN LAGOS, NIGERIA**

Bilikis Temitope Folarin^{1, 2}, Mohamed Abou-Elwafa Abdallah², Temilola Oluseyi¹, Kehinde
Olayinka¹, Stuart Harrad^{2*}

¹Department of Chemistry,
University of Lagos,
Lagos,
Nigeria

²School of Geography, Earth and Environmental Sciences,
University of Birmingham,
Birmingham,
United Kingdom

*Corresponding author

S.J.Harrad@bham.ac.uk

Abstract

Concentrations of 7 indicator polychlorinated biphenyls (PCBs) were measured in dust and soil samples from 12 power stations collected over the two major seasons of the Nigerian climate. Median Σ PCB₇ concentrations in soil ranged from 2 ng/g for power station A to 220 ng/g for power station I; while those in dust ranged from 21 ng/g for power station L to 2,200 ng/g for power station I. For individual congeners, median PCB concentrations ranged from 3.8 ng/g for PCB 101 to 52 ng/g for PCB 180 in dust, and <0.07 ng/g for PCB 28 to 5.9 ng/g for PCB 153 in soil. The type of power station activity exerted a significant influence on concentrations of Σ PCB₇ in dust and soil (generation > transmission > distribution). Congener patterns in dust and soil samples were compared using principal component analysis (PCA) with those in transformer oil samples from 3 of the power stations studied and with common PCB mixtures (Aroclors). This revealed congener patterns in soil were more closely related to that in the transformer oil than dust. Congener patterns in most samples were similar to Aroclor 1260. Concentrations of PCBs in soil samples close to the transformers significantly exceeded those in soil sampled further away.

Keywords

PCBs; power generation plants; Africa; transformers

1. Introduction

Polychlorinated biphenyls (PCBs) are a class of industrial chemicals listed as persistent organic pollutants (POPs) under the Stockholm convention (UNEP, 2009). Prior to their restriction, PCB mixtures had been used extensively as dielectric fluids in electrical appliances and power generating plants as well as additives in many other non-electrical products in everyday use (Xing *et al.*, 2011; Breivik *et al.*, 2002).

The large-scale production and usage of PCBs, has resulted in widespread contamination of the environment (Harrad *et al.*, 1994). Efforts to gradually eliminate the presence of these toxic, ubiquitous chemicals, led to ban on their production (Robinson, 2009; Koppe and Keys, 2002). However, over four decades post-restriction, PCBs remain in circulation within the global environment, due to their persistent nature (Gouin *et al.*, 2005). Moreover, PCBs continue to be released into the environment, for example through spillage from transformers during maintenance and day-to-day operations in power stations (Tue *et al.*, 2016; Xing *et al.*, 2011).

The soil is a primary recipient of many outdoor contaminants (Domotorova *et al.*, 2012). PCBs have been detected in soil around power plants (generating, transmitting and distribution transformers) and other e-waste sites in China (Tang *et al.*, 2010; Shen *et al.*, 2008). Meanwhile, reported sources of PCBs in indoor environments include paints, building sealants and adhesives. PCB fumes can be released from hot electrical gadgets, wires and cables, since PCBs have been used as components of these products because of their excellent insulating properties (Xing *et al.*, 2011; Kohler *et al.*, 2005).

The high K_{OW} values of PCBs increases their tendency to bind to dust and soil (Kuusisto *et al.*, 2007). Indoor dust has been reported as a major sink of PCBs and other pollutants in indoor environments (Wang *et al.*, 2015; Whitehead *et al.*, 2011), leading to exposure via inhalation, ingestion and/or dermal uptake (Harrad *et al.*, 2006; Jones-Otazo *et al.*, 2005; Freels *et al.*, 2007).

To the best of our knowledge, there are only two reports of PCB concentrations in indoor dust in Africa. Firstly, Abafe *et al.*, 2015 measured concentrations of three PCB congeners (PCB 28, 153, and 180) in indoor dust samples from ten homes, eleven offices, and thirteen computer laboratories in Durban, South Africa. The second report measured six indicator PCBs in dust samples from three microenvironment categories (homes, cars and offices) in Nigeria (Harrad *et al.*, 2016). The present study measures concentrations of PCBs in dust and soil from the vicinity of power stations in Nigeria to test our hypothesis that such stations are a source of PCBs to the surrounding environment. This work reports for the first time, the concentrations of PCBs in dust samples from power stations in Nigeria and provides only the second report on PCBs in Nigerian dust.

Concentrations of PCBs in soil from non-African countries have been extensively reported. Vane *et al.*, 2014 (UK); and Jiang *et al.*, 2011 (China), reported concentrations of PCBs in soil from urban areas, while Melynk *et al.*, 2015 (Poland) reported PCBs in landfill soil. Concentrations of PCBs in industrial and agricultural soils were also compared in Pakistan by Syed *et al.*, 2013. The few reports on PCBs in African soil are from South Africa (Batterman *et al.*, 2009); Kenya (Sun *et al.*, 2016); Ghana (Tue *et al.*, 2016) and Nigeria (Alani *et al.*, 2013). All these African reports are centred around PCBs in soil from urban areas, except for Tue *et al.*, 2016 that measured dioxin-like PCBs and other dioxin related compounds released from open burning of e-waste in Ghana. The current study is the first investigation of PCBs in soil in the vicinity of power stations in Nigeria.

Against this backdrop, this paper reports the concentrations of PCBs in 48 office dust and 48 soil samples collected within and in the vicinity of 12 power stations in Lagos, Nigeria (6 power distribution stations, 5 power transmission stations, and 1 former generation station within Lagos, Nigeria). These data are used to: (i) test the hypothesis that power stations in Nigeria constitute a source of PCBs to the surrounding environment; (ii) evaluate the effect of

seasonal variation (wet and dry seasons) on concentrations of PCBs in soil and dust samples; and (iii) compare the pattern of PCBs in transformer oil samples with dust and soil samples.

2. Materials and Methods

2.1 Sample collection

A total of 48 dust samples and 48 soil samples were collected from twelve power stations in Lagos, Nigeria in the months of June 2015 (24 dust samples and 24 soil samples) and December 2015 (24 dust samples and 24 soil samples). Surface dusts were collected from two offices within each power station while soil samples were collected from two points for each power station. The first point was 1-3 m and the second sampling point 50-100 m away from the transformer plants. Collection of top soil (0 - 5 cm depth) was done using a hand trowel, which was washed and pre-cleaned with hexane before each sampling. Following collection, each sample was passed through a 2 mm sieve, wrapped in aluminium foil and stored at – 20 °C in individual zip lock bags.

Dust samples were collected from elevated surfaces (e.g. shelves and air conditioning panels, electrical control panels etc) using a paintbrush which had been washed and pre-cleaned with hexane before each sampling, and wrapped in aluminium foil. Similar to soils, dust samples were kept in individual zip lock bags and stored at – 20 °C. Both soil and dust samples were transported to the University of Birmingham for analysis. At Birmingham, the dust samples were sieved with a 500 µm sieve, using procedures similar to Harrad et al. (2016). In Birmingham, both soil and dust samples were stored at 4 °C prior to analysis.

Transformer oil samples were collected from three of the studied power stations. Two were used transformer oil while the third one was a fresh unused oil. Oil sample 1 was collected from Site I and J, sites that are linked - one being a former generation station (I) and the other

a transmission station (J). The other used oil was from site H, while the unused oil was from site K.

2.2 Chemicals and materials

Individual indicator PCB standards (#s 28, 52, 101, 118, 138, 153, 180), and 5 native internal standards (ISs) (PCB 34, 62, 119, 131 and 173) were purchased from LGC standards, UK. These native PCB congeners are absent in environmental samples and have been previously reported to be suitable for use as internal standards (Ayris *et al.*, 1997). Recovery determination standard (RDS), PCB 129 was from QMX laboratories, UK; HPLC grade solvents (DCM, n-Hexane) were from Fisher Scientific, UK. Analytical grade reagents purchased from Sigma Aldrich, UK were: anhydrous sodium sulfate, which was baked at 450 °C for 4 hours before use and florisil (60-100 mesh) activated at 120 °C for 2 hours.

2.3 Sample extraction and clean up

Accurately weighed dust (approx. 200 mg) and soil (between 1-3 g) samples were carefully introduced into glass extraction tubes that had prior to use been washed, oven dried for several hours, and thoroughly rinsed with extraction solvent (1:1 (v/v) hexane: DCM). A known amount of internal standard was added prior to addition of 2 mL and 5 mL extraction solvent (1:1 (v/v) hexane: DCM) for dust and soil respectively. The mixture was vortexed for about 1 minute before ultrasonic extraction for 15 minutes and 30 minutes for dust and soil samples respectively. The supernatant was separated from the solute by centrifugation at 3500 rpm for 5 mins. The extraction was repeated twice for each sample using fresh solvent for each repeat extraction. Extracts were combined and blown down to dryness under a gentle stream of nitrogen.

Concentrated crude extracts were reconstituted in 2 mL hexane and shaken with 2 mL concentrated H₂SO₄, to eliminate lipids. The organic layer was removed and concentrated to about 1 mL before passing through a bed of florisil, pre-conditioned with 6 mL hexane, and the PCBs on the florisil column were eluted with 20 mL hexane. The eluate was reduced under N₂ to incipient dryness and reconstituted in 150 µL recovery determination standard in iso-octane. While for the majority of samples, this relatively simple clean-up was sufficient; for some samples, DMSO clean up was introduced before florisil clean up to eliminate aliphatic hydrocarbons which were major interferences by introducing 2 mL DMSO to 1 mL acid cleaned extract (in hexane). In such instances, the mixture was vortexed for 1 minute and then centrifuged at 3500 rpm for 2 minutes to facilitate removal of the top hexane aliphatic-containing layer. To the DMSO aromatic-containing layer, 1 mL water was added, and the mixture vortexed to break down the DMSO charge transfer complex. The aqueous DMSO mixture was then back extracted with 4 × 1 mL hexane. These hexane back extracts were combined and blown down ready for florisil clean up as described above.

For transformer oil samples, 50 µL (approx. 40 mg) of oil was diluted with 1 mL hexane and cleaned up using a sequential combination of sulfuric acid, DMSO, and florisil clean up methods as described above.

2.4 Instrumental Analysis

Target PCBs were quantified in calibration standard and sample extracts on an Agilent 5975C GC-MSD, fitted with a RestekRXi-PAH capillary column (40 m × 180 µm × 0.01 µm) using the GC and MS parameters stated in the supplementary material (Table S1). The GC inlet temperature was set to 280 °C, injection was performed in splitless mode. A carrier gas (Helium) flow of 0.5 mL/min was employed. The initial oven temperature was set at 140 °C and held for 2 mins, ramped at 5 °C/min to 215 °C, held for 5 mins and thereafter ramped at 2

°C/min to 245 °C. The temperatures of the quadrupole and ion source were 150 °C and 230 °C respectively.

The target PCBs monitored belong to 5 PCB homologues. To quantify target PCBs (PCB # - 28, 52, 101, 118, 138, 153, and 180), internal standard quantification was employed. PCB # - 34, 62, 119, 131, and 173 were used as internal standards for the tri CB, tetra-CB, penta-CB, hexa-CB, and hepta-CB respectively. The MS was programmed in the selected ion monitoring (SIM) mode, with the two most abundant ions monitored for each PCB homologue. The masses of the 2 ions monitored and their percentage abundance are presented in Table S2 (supplementary material).

2.5 QA/QC

Two standard reference materials (SRM 2585, NIST organics in indoor dust and CRM 481, PCBs in industrial soil) were analysed for method validation. As a measure of precision, 6 replicate analysis of PCBs in SRM 2585 and CRM 481, extracted and cleaned-up as described above, gave a RSD ranging from 4 % to 13 % and 5 % to 10 % for dust and soil reference materials respectively. Concentrations of PCBs in these reference materials obtained using our methods were deemed acceptably similar to the certified values (Tables S3 and S4). Mean recoveries of internal standards in SRM2585 ranged from 80 % to 94 % and are provided in table S5. Analyses of method blanks were carried out once after every 12 samples. The method blanks were treated in a similar way as the samples but sodium sulfate was extracted in place of dust/soil under study. Results obtained were not corrected for blank concentrations since almost all the target PCBs were absent from blanks, and even where present in blanks, concentrations were less than 5 % of the concentration found in the corresponding sample. The limit of detection was obtained as the concentration equivalent to a signal to noise ratio of 3:1. LOD values obtained for dust were: 0.09 ng/g, 0.08 ng/g, 0.11 ng/g, 0.11 ng/g, 0.11 ng/g, and 0.08 ng/g for PCB 28, 52, 101, 118, 138 and 180

respectively. The corresponding LOD values for soil were: 0.07 ng/g, 0.06 ng/g, 0.07 ng/g, 0.06 ng/g, 0.07 ng/g, 0.07 ng/g, and 0.05 ng/g respectively.

2.6 Data treatment

The results were analysed statistically using Minitab 18. Given the distribution of our data revealed by visual inspection, non-parametric methods were used to test for significant differences in concentrations of PCBs in different sample groups. Specifically, a Wilcoxon test was performed to test: (a) differences in concentrations of PCBs in soil samples taken at two distances from each power station, and (b) the influence of season on concentrations of PCBs in our soil samples. In addition, a Kruskal-Wallis test was used to test if different categories of power station activity exerted a significant influence on the concentration of PCBs in our soil samples. Finally, principal component analysis (PCA) was used to compare PCB congener patterns in samples with those in transformer oil and Aroclor mixtures.

3. Results and Discussion

3.1 Differences in concentration of PCBs in soil samples from point 1 and 2.

We hypothesised that soil samples taken closer to the power station facility would have higher concentrations of PCBs compared to samples acquired further away. Table 1 shows the ΣPCB_7 concentrations in samples taken from both points at each power station studied. A Wilcoxon test revealed that concentrations of ΣPCB_7 closer to the power station (point 1 samples) were significantly higher ($p=0.013$) than those in samples taken further away (point 2 samples). Similar significantly ($p<0.05$) elevated concentrations in point 1 samples were observed for all individual PCB congeners. This increment in PCB contamination of soil

close to power generation facilities, suggests strongly that such activities have constituted a substantial source of PCBs to their local environments.

Power station I (PS I), the former generation station, was the only site that had high concentrations of target PCBs in both point 1 as well as point 2 samples (Table 1). The high concentrations of PCBs in the point 2 soil from PS I could be due to emissions arising from the conveying of transformers in and out of the power station since the power station is now used for transformer repair purposes.

3.2 Influence of power station activity category on PCB concentrations in dust and soil

The 12 power stations studied comprised 6 distribution sub-stations, 5 transmission stations and 1 former generation station, now used for repair of transformers. A Kruskal Wallis test was performed to test whether the type of activity (transmission, distribution, or transformer dismantling (station I)) impacted on the concentrations of PCBs in soil and dust. On the basis of power station activity, results revealed a statistically significant difference ($P < 0.05$) in concentrations of total PCBs in dust and soil-between the three categories. For dust, the mean ΣPCB_7 concentration for generation station samples was 2241.8 ng/g; in transmission station samples ($n=5$) the mean ΣPCB_7 concentration was 314.5 ng/g, while concentrations in the distribution station samples ($n=6$; mean $\Sigma\text{PCB}_7 = 122.8$ ng/g) were the lowest. For soil, the mean ΣPCB_7 concentration for generation station samples was 197.7 ng/g; in transmission station samples ($n=5$) the mean ΣPCB_7 concentration was 92.6 ng/g, while distribution station samples ($n=6$; mean $\Sigma\text{PCB}_7 = 38.3$ ng/g) had the lowest concentrations. The persistent nature of PCBs was confirmed from the results obtained for PCBs in soil samples around power generation transformers from PS I that has been non-functional for over a decade, with residual concentrations very close to being the highest in soil samples from power stations in this study. These high levels could be residual concentration from day to day operations (before the ban of PCBs) when the power station was fully functional.

3.3 Congener pattern in dust, soil and transformer oil samples relative to PCB commercial mixtures

PCBs were mostly used as mixtures, manufactured under various trade names by different producers. One of the most popular of these PCB mixtures were known as Aroclors. Principal component analysis (PCA) was conducted to identify which of the Aroclors have a congener pattern similar to those in samples from this study. It was observed that many of the soil samples, some of the dust samples, as well as transformer oil samples have positive component scores in PC 1, similar to Aroclor 1260. PC 1 could explain 39 % of the congener variations within the data set and was driven positively by PCB 153 > 101 > 118 > 138, with eigenvectors of 0.490, 0.446, 0.435, 0.279 respectively. PC 1 was driven negatively by PCB 52 > 28 > 180 with eigenvectors of -0.360, -0.310, -0.262 respectively.

Aroclor 1260 had positive component scores in PC 2, together with some of the dust and soil samples. The oil from PS I and J also displayed positive PC 2 scores. PC 2, which explained an additional 24.8 % of the variation in congener patterns in the samples, was driven towards the positive direction substantially by PCB 180, with an eigenvector of 0.512, and also by PCB 138 and 153, with eigenvectors of 0.399 and 0.142 respectively. PCB 28 made the greatest negative contribution to PC 2, with an eigenvector of -0.427; while PCB 101, 52 and 118 also contributed negatively to PC 2, with eigenvectors of -0.375, -0.36, and -0.326 respectively.

Worthy of note are the soil (L-SOIL) and dust (L-DUST) samples from PS L (Figure 1), these samples from power station L both had the highest negative scores in PC 2, of all the soil and dust samples. Also, of all the soil and dust samples, L-SOIL contributed most negatively to scores of PC 1 while L-DUST contributed most positively to the scores of PC 1. It was observed that Aroclor 1242 displayed a very similar PC1 score to L-SOIL. Moreover,

Aroclors 1242 and 1016 displayed PC2 scores similar to those for both soil and dust from PS L (Figure 1).

3.4 Effect of seasonal variation on concentrations of target PCBs in dust and soil

Wilcoxon test employed to test the effect of seasonal variation revealed statistically indistinguishable concentrations of individual target PCBs in dust samples for the month of June and December. A similar absence of seasonal variation was observed when comparing PCB concentrations in soil in June with those from the same locations in December.

3.5 Comparison of concentrations obtained in soil and dust with relevant reports

Concentrations of PCBs in dust from our Nigerian power station offices exceed those reported for PCBs in dust from Nigerian cars, homes and office (Harrad et al, 2016). They also exceed those in other previous studies, (Tue *et al.*, 2013; Tan *et al.*, 2007; Ali *et al.*, 2012) reported PCBs in Vietnamese homes around e-waste sites; Singaporean, Kuwaiti, and Pakistani homes respectively; but are lower than those reported by Abafe and Martincigh, 2015 and Zheng *et al.*, 2015 for South African offices and homes around e-waste handling sites respectively. Worthy of note is the similar congener profile in our report to the other two reports on PCBs in dust from African countries (Harrad *et al.*, 2016; Abafe and Martincigh, 2015) (Table 2). This may imply that the PCB formulations used in Nigeria and South Africa are similarly dominated by higher chlorinated congeners (Harrad *et al.*, 2009) – such as Aroclor 1260.

With respect to soil, concentrations of both individual congeners and ΣPCB_7 in our study exceed those in other studies from locations not directly impacted by power station activities (Table 3). This is further evidence of the potential importance of power stations as sources of local contamination with PCBs, and further study appears warranted.

4. Conclusion

The concentrations of PCBs in samples of soil taken 1-3 m distant from Nigerian power stations exceed significantly those detected in soil samples taken 50-100 m from the same facilities. The type of activity conducted at each power station exerted a significant effect on the concentrations of PCBs in soil and dust samples with the distribution stations having the lowest concentrations.

Irrespective of the power station activity, the PCB congener pattern in soil was observed to be more similar to that in transformer oil taken from some of our power stations than that in dust samples. We thus infer that concentrations of PCBs in soils in this study were substantially influenced by emissions from transformer oil, while our dust samples were influenced by alternative sources such as control panels present inside power stations. The evidence presented here supports the hypothesis that activities at power stations have contributed significantly to PCB contamination of the surrounding environment.

Acknowledgements

The authors acknowledge gratefully funding received from the Analytical Chemistry Trust Fund in the guise of a Developing World Scholarship to Bilikis Folarin (Ref. 15/600504/14).

References

Abafe, O. A. & Martincigh, B. 2015. An assessment of polybrominated diphenyl ethers and polychlorinated biphenyls in the indoor dust of e-waste recycling facilities in South Africa: implications for occupational exposure. *Environmental Science and Pollution Research*, 22, 14078-14086.

- Abafe, O. A. & Martincigh, B. S. 2015. Polybrominated diphenyl ethers and polychlorinated biphenyls in indoor dust in Durban, South Africa. *Indoor Air*, 25, 547-556.
- Abdallah, M. A., Drage, D. & Harrad, S. 2013. A one-step extraction/clean-up method for determination of PCBs, PBDEs and HBCDs in environmental solid matrices. *Environmental Science Processes & Impacts*, 15, 2279-2287.
- Alani R., Olayinka K. & Alo B. 2013. The level of persistent, bioaccumulative, and toxic (PBT) organic micropollutants contamination of Lagos soils. *Journal of Environmental Chemistry and Ecotoxicology Vol. 5(2)*, 26-38.
- Ali, N., Ali, L., Mehdi, T., Dirtu, A. C., Al-Shammari, F., Neels, H. & Covaci, A. 2013. Levels and profiles of organochlorines and flame retardants in car and house dust from Kuwait and Pakistan: Implications for human exposure via dust ingestion. *Environment International*, 55, 62-70.
- Ayris, S., Currado, G. M., Smith, D. & Harrad, S. 1997. GC/MS procedures for the determination of PCBs in environmental matrices, *Chemosphere*, 35, 905-917.
- ATSDR, 2000. Toxicological profile for Polychlorinated biphenyls: Chemical and physical information, chapter 4, 450-463. <https://www.atsdr.cdc.gov/toxprofiles/tp17-c4.pdf>
- ATSDR, 2014. Case studies in environmental medicine: Polychlorinated biphenyls (PCBs) Toxicity. 33-34. <http://www.atsdr.cdc.gov/csem/csem.asp?csem=30&po=8>
- Batterman, S., Chernyak, S., Gouden, Y., Hayes, J., Robins, T. & Chetty, S. 2009. PCBs in air, soil and milk in industrialized and urban areas of KwaZulu-Natal, South Africa. *Environ Pollut*, 157, 654-63.
- Breivik, K., Sweetman, A., Pacyna, J. M. & Jones, K. C. 2002a. Towards a global historical emission inventory for selected PCB congeners — a mass balance approach: 1. Global production and consumption. *Science of The Total Environment*, 290, 181-198.


- Breivik, K., Sweetman, A., Pacyna, J. M. & Jones, K. C. 2002b. Towards a global historical emission inventory for selected PCB congeners — a mass balance approach: 2. Emissions. *Science of The Total Environment*, 290, 199-224.
- Diamond, M., Melymuk, L., Csiszar, S.A. & Robson, M. 2010. Estimation of PCB stocks, emissions, and urban fate: will our policies reduce concentrations and exposure? *Environ. Sci. Technol.* 44, 2777-2783.
- Domotorova, M., Sejakova, Z., Kocan, A., Conka, K., Chovancova, J. & Fabisikova, A. 2012. PCDDs, PCDFs, dioxin-like PCBs and indicator PCBs in soil from five selected areas in Slovakia. *Chemosphere*, 89, 480-485.
- Freels, S., Chary, L. K., Turyk, M., Piorkowski, J., Mallin, K., Dimos, J., Anderson, H., McCann, K., Burse, V. & Persk, Y. V. 2007. Congener profiles of occupational PCB exposure versus PCB exposure from fish consumption. *Chemosphere*, 69, 435-443.
- Gouin, T., Harner, T., Daly, G. L., Wania, F., Mackay, D. & Jones, K. C. 2005. Variability of concentrations of polybrominated diphenyl ethers and polychlorinated biphenyls in air: implications for monitoring, modeling and control. *Atmospheric Environment*, 39, 151-166.
- Harrad, S., Abdallah, M. A. E. & Oluseyi, T. 2016. Polybrominated diphenyl ethers and polychlorinated biphenyls in dust from cars, homes, and offices in Lagos, Nigeria. *Chemosphere*, 146, 346-353.
- Harrad, S. & Diamond, M. 2006. New Directions: Exposure to polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs): Current and future scenarios. *Atmospheric Environment*, 40, 1187-1188.
- Harrad, S., Ibarra, C., Robson, M., Melymuk, L., Zhang, X. M., Diamond, M. & Douwes, J. 2009. Polychlorinated biphenyls in domestic dust from Canada, New Zealand, United Kingdom and United States: Implications for human exposure. *Chemosphere*, 76, 232-238.

- Harrad, S., Ibarra, C., Diamond, M., Melymuk, L., Robson, M., Douwes, J., Roosens, L., Dirtu, A. C. & Covaci, A. 2008. Polybrominated diphenyl ethers in domestic indoor dust from Canada, New Zealand, United Kingdom and United States. *Environment International*, 34, 232-238.
- Jiang, Y., Wang, X., Zhu, K., Wu, M., Sheng, G. & Fu, J. 2011. Polychlorinated biphenyls contamination in urban soil of Shanghai: level, compositional profiles and source identification. *Chemosphere*, 83, 767-773.
- Jones-Otazo, H.A., Clarke, J.P., Diamond, M.L., Archbold, J.A., Ferguson, J.A., Harner, T., Richardson, G.M., Ryan, J.J., Wilford, B. 2005. Is house dust the missing exposure pathway to PBDEs? An analysis of the urban fate and human exposure to PBDEs. *Environ. Sci. Technol*, 39, 5121-5130.
- Kohler, M., Tremp, J., Zennegg, M., Siler, C., Minder-Kohler, S., Beck, M., Lienemann, P., Wegmann, L. & Schmid, P. 2002. Joint sealants: an overlooked diffuse source of polychlorinated biphenyls in buildings. *Environ. Sci. Technol*, 39, 1967-1973.
- Koppe, J.G. & Keys, J. 2002. PCBs and the precautionary principle. European Environmental Agency, Luxembourg: *Environment issue report*, Report No. 22.
- Kuusisto, S., Lindroos, O., Rantio, T., Priha, E. & Tuhkanen, T. 2007. PCB contaminated dust on indoor surfaces - Health risks and acceptable surface concentrations in residential and occupational settings. *Chemosphere*, 67, 1194-1201.
- Leung, A., Cai, Z. W. & Wong, M. H. 2006. Environmental contamination from electronic waste recycling at Guiyu, southeast China. *Journal of Material Cycles and Waste Management*, 8, 21-33.
- Liu, M., Huang, B., Bi, X., Ren, Z., Sheng, G., & Fu, J. 2013. Heavy metals and organic compounds contamination in soil from e-waste region in South China. *Environ. Sci. Process. Impacts*. 15, 919-29.

- Melnyk, A., Dettlaff, A., Kuklinska, K., Namiesnik, J. & Wolska, L. 2015. Concentration and sources of polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) in surface soil near a municipal solid waste (MSW) landfill. *Sci Total Environ*, 530-531, 18-27.
- Robinson, B. H. 2009. E-waste: An assessment of global production and environmental impacts. *Science of The Total Environment*, 408, 183-191.
- Shen, C., Huang, S., Wang, Z., Qiao, M., Tang, X., Yu, C., Shi, D., Zhu, Y., Shi, J., Chen, X., Setty, K. & Chen, Y. 2008. Identification of Ah receptor agonists in soil of e- waste recycling sites from Taizhou area in China. *Environ.Sci. Technol.* 42: 49-55.
- Sun, H., Qi, Y., Zhang, D., Li, Q. X. & Wang, J. 2016. Concentrations, distribution, sources and risk assessment of organohalogenated contaminants in soils from Kenya, Eastern Africa. *Environ Pollut*, 209, 177-85.
- Syed, J. H., Malik, R. N., Li, J., Zhang, G. & Jones, K. C. 2013. Levels, distribution and air-soil exchange fluxes of polychlorinated biphenyls (PCBs) in the environment of Punjab Province, Pakistan. *Ecotoxicol Environ Saf*, 97, 189-95.
- Tan, J., Cheng, S. M., Loganath, A., Chong, Y. S. & Obbard, J. P. 2007. Selected organochlorine pesticide and polychlorinated biphenyl residues in house dust in Singapore. *Chemosphere*, 68, 1675-1682.
- Tang, X., Shen, C., Shi, D., Cheema, S. A., Khan, M. I., Zhang, C. & Chen, Y. 2010. Heavy metal and persistent organic compound contamination in soil from Wenling: An emerging e-waste recycling city in Taizhou area, China. *Journal of Hazardous Materials*, 173, 653-660.
- Tue, N. M., Goto, A., Takahashi, S., Itai, T., Asante, K. A., Kunisue, T. & Tanabe, S. 2016. Release of chlorinated, brominated and mixed halogenated dioxin-related compounds to soils from open burning of e-waste in Agbogbloshie (Accra, Ghana). *Journal of Hazardous Materials*, 302, 151-157.

- Tue, N. M., Takahashi, S., Suzuki, G., Isobe, T., Viet, P. H., Kobara, Y., Seike, N., Zhang, G., Sudaryanto, A. & Tanabe, S. 2013. Contamination of indoor dust and air by polychlorinated biphenyls and brominated flame retardants and relevance of non-dietary exposure in Vietnamese informal e-waste recycling sites. *Environment International*, 51, 160-167.
- UNEP, 2009. Stockholm convention on persistent organic pollutants.<http://chm.pops.int>.
- Vane, C. H., Kim, A. W., Beriro, D. J., Cave, M. R., Knights, K., Moss-Hayes, V. & Nathanail, P. C. 2014. Polycyclic aromatic hydrocarbons (PAH) and polychlorinated biphenyls (PCB) in urban soils of Greater London, UK. *Applied Geochemistry*, 51, 303-314.
- Wang, B. L., Pang, S. T., Sun, J. P., Zhang, X. L., Li, X. L., Sun, Y. G., Lu, X. M. & Zhang, Q. 2015. Levels of polychlorinated biphenyls in settled house dust from urban dwellings in China and their neurodevelopmental effects on preschool-aged children. *Science of the Total Environment*, 505, 402-408.
- Wang, Y. L., Hu, J. X., Lin, W., Wang, N., Li, C., Luo, P., Hashmi, M. Z., Wang, W. B., Su, X. M., Chen, C., Liu, Y. D., Huang, R. L. & Shen, C. F. 2016. Health risk assessment of migrant workers' exposure to polychlorinated biphenyls in air and dust in an e-waste recycling area in China: Indication for a new wealth gap in environmental rights. *Environment International*, 87, 33-41.
- Whitehead, T., Metayer, C., Buffler, P. & Rappaport, S. M. 2011. Estimating exposures to indoor contaminants using residential dust. *Journal of Exposure Science and Environmental Epidemiology*, 21, 549-564.
- Whitehead, T. P., Brown, F. R., Metayer, C., Park, J. S., Does, M., Dhaliwal, J., Petreas, M. X., Buffler, P. A. & Rappaport, S. M. 2014. Polychlorinated Biphenyls in Residential Dust: Sources of Variability. *Environmental Science and Technology*, 48, 157-164.

- Xing, G. H., Liang, Y., Chen, L. X., Wu, S. C. & Wong, M. H. 2011. Exposure to PCBs, through inhalation, dermal contact and dust ingestion at Taizhou, China - A major site for recycling transformers. *Chemosphere*, 83, 605-611.
- Zhang, X. M., Diamond, M. L., Robson, M. & Harrad, S. 2011. Sources, Emissions, and Fate of Polybrominated Diphenyl Ethers and Polychlorinated Biphenyls Indoors in Toronto, Canada. *Environmental Science and Technology*, 45, 3268-3274.
- Zheng, X. B., Xu, F. C., Chen, K. H., Zeng, Y. H., Luo, X. J., Chen, S. J., Mai, B. X. & Covaci, A. 2015. Flame retardants and organochlorines in indoor dust from several e-waste recycling sites in South China: Composition variations and implications for human exposure. *Environment International*, 78, 1-7.



Soil samples (yellow triangles); dust samples (purple rhombuses); oil from power station I&J (red circle); oil from power station H (blue triangle); oil from power station K (green square)

Figure 1: Score plot for principal component 1 versus principal component 2 for soil, indoor dust, transformer oil and Aroclors

Table 1: Concentrations of Σ PCB₇ (ng/g) in soil from both sampling points at each site

Power station (type)	June		December		Mean Point 1	Mean Point 2	Mean Σ PCB ₇
	point 1	point 2	point 1	point 2			
A (distribution)	3.5	2.4	0.4	1.2	2	1.8	1.9
B (transmission)	40.9	1.6	16.8	0.5	29	1.1	15
C (distribution)	27.4	17.4	10.8	2.8	19	10	14.6
D (distribution)	50	26.6	310	4.6	180	16	96.6
E (transmission)	320	10.7	71.3	5	200	7.9	101.7
F (distribution)	148.4	51.4	43.3	26.3	96	39	67.3
G (distribution)	74.6	10.5	15.7	16.3	45	13	29.3
H (transmission)	162.2	18	254.9	12.9	210	16	112
I (generation)	220.1	211.1	125.7	233.9	170	220	197.7
J (transmission)	317.6	57.3	422.4	33.7	370	46	207.8
K (distribution)	33	17.9	27	3.1	30	11	20.3
L (transmission)	83.2	10.8	7.4	4.3	45	7.6	26.4

Table 2: Comparison of median (mean) concentrations (ng/g) of 7 indicator PCBs in dust from this study with those from selected other studies

COUNTRY (REF.)	MICROENVIRONMENT	N	PCB 28	PCB 52	PCB 101	PCB 118	PCB 138	PCB 153	PCB 180
NIGERIA (This study)	POWER STATION OFFICES	4	16.3 (22.4)	12.5 (23.4)	3.8 (38.3)	4 (38.3)	24.9 (73.3)	8.9 (91.3)	51.7 (92.4)
NIGERIA (Harrad et al., 2016)	OFFICES	1	5.2 (4.8)	4.4 (4.6)	6.2 (8.7)	NA	5.1 (7.3)	6.5 (10)	14 (14)
S. AFRICA (Abafe et al., 2015)	OFFICES	1	28.3 (64)	NA	NA	NA	NA	136 (170)	812 (689)
S. AFRICA (Abafe et al., 2015)	E-WASTE&ICT WORKSHOP	5	9 (7.8)	NA	NA	NA	NA	47 (93)	109 (134)
UK (Harrad et al., 2009)	HOME	2	3.4 (6.3)	1.8 (5.6)	1.2 (6.1)	0.92 (4.3)	1.1 (4.1)	1.2 (3.3)	0.89 (1.8)
UK (Abdallah et al., 2013)	HOME	5	8.2 (10.1)	7.4 (7.5)	6.9 (5.3)	1.3 (1.8)	0.8 (1.8)	0.9 (1.6)	0.3 (0.8)
VIETNAM (Tue et al., 2013)	HOMES AROUND E-WASTE SITES	1	1.3 (NS)	0.65 (NS)	0.91 (NS)	1.5 (NS)	2 (NS)	1 (NS)	0.59 (NS)
SINGAPORE (Tan et al., 2007)	HOME	3	0.2 (0.3)	NS	0.5 (0.6)	0.3 (0.7)	NS	0.5 (0.8)	0.2 (0.3)
CHINA (Zheng et al., 2015)	E-WASTE SITES	1	NA	NA	634 (NS)	21 (NS)	462 (NS)	415 (NS)	120 (NS)
KUWAIT (Ali et al., 2013)	HOME	5	NA	NA	<0.2 (63)	<0.2 (35)	<0.1 (28)	<0.2 (20)	0.4 (7)
PAKISTAN (Ali et al., 2013)	HOME	1	NA	NA	<0.2 (<0.2)	<0.2 (<0.2)	<0.1 (0.5)	0.4 (0.8)	0.3 (1.2)
US (Harrad et al., 2009)	HOME	2	5.1 (9.5)	6.7 (7.9)	8.7 (10)	5.5 (10)	6.5 (8.6)	7.1 (8.4)	2.6 (4.5)
CANADA (Harrad et al., 2009)	HOME	1	7.3 (10)	7.2 (12)	8.8 (15)	8.7 (13)	9.5 (12)	9.9(11)	6.8 (8.5)
NEW ZEALAND (Harrad et al., 2009)	HOME	2	2.3 (3.3)	1.4 (2.6)	1.6 (3.1)	0.95 (1.9)	1.8 (2.8)	1.4(2.7)	1.3 (2.3)

NA- NOT APPLICABLE (Congeners were not analysed) ; NS-NOT SPECIFIED (Congeners were analysed but concentrations were not reported)

Table 3: Comparison of median (mean) concentrations (ng/g) of PCBs in soil from this study with those from selected other studies

COUNTRY (REF.)	SAMPLING LOCATIONS	N	PCB 28	PCB 52	PCB 101	PCB 118	PCB 138	PCB 153	PCB 180
NIGERIA (This study)	VICINITY OF POWER STATIONS	48	<0.07 (6.9)	0.3 (2.9)	1.6 (7.9)	1.5 (5.6)	4.3 (23.6)	5.9 (15.4)	4 (11.9)
NIGERIA (Alani et al.,2013)	URBAN	5	ND (0.8)	ND (0.4)	ND (0.3)	NS	0.24 (0.5)	0.48 (0.5)	ND (0.5)
S. AFRICA (Batterman et al., 2009)	URBAN	3	NS (2.4)	NS (2.1)	NS (4.2)	NS (4.4)	NS (8.4)	NS (9.9)	NS (4.9)
KENYA (Sun et al., 2016)	RURAL	6	NS (0.2)	NS (2.4)	NS (6.7)	NS (3.8)	NS (1.9)	NS (10.3)	NS (1.9)
UK (Vane et al., 2014)	URBAN	76	0.2 (0.8)	0.5 (3)	0.4 (3.3)	<0.2 (2.1)	1.6 (5.9)	1.6 (5.1)	0.1 (1.4)
POLAND (Melnyket al., 2015)	AROUND LANDFILL	12	0.8 (1.13)	0.5 (0.55)	0.4 (0.49)	0.3 (0.57)	0.5 (0.62)	0.4 (0.55)	0.5 (0.61)
CHINA (Jiang et al., 2011)	URBAN	55	NS (0.6)	NS (0.03)	NA	NS (0.06)	NS (0.14)	NS (0.16)	NA
PAKISTAN (Syed et al., 2013)	INDUSTRIAL/AGRICULTURAL	50	NS (1.2)	NS (0.3)	NS (0.5)	NS (0.3)	NS (1.3)	NS (0.9)	NS (0.9)
AUSTRALIA (Abdallah et al., 2013)	URBAN	5	<LOD	0.05 (0.06)	<LOD (0.38)	NA	0.4 (0.54)	0.4 (0.45)	0.1 (0.14)

NA- NOT APPLICABLE (Congeners were not analysed) ; NS-NOT SPECIFIED (Congeners were analysed but concentrations were not reported).

Supplementary Material

[Click here to download Supplementary Material: Folarin et al 2018 Supplementary Material.docx](#)