

General group exercise in low back pain management in a military population, a comparison with specific spine group exercise

Surtees, Joanna; Heneghan, Nicola

DOI:

[10.1136/jramc-2018-001011](https://doi.org/10.1136/jramc-2018-001011)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Surtees, J & Heneghan, N 2018, 'General group exercise in low back pain management in a military population, a comparison with specific spine group exercise: a service evaluation', *Journal of the Royal Army Medical Corps*. <https://doi.org/10.1136/jramc-2018-001011>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This article has been accepted for publication in *Journal of the Royal Army Medical Corps* following peer review. The definitive copyedited, typeset version is available online at:

Surtees JE, Heneghan NR

General group exercise in low back pain management in a military population, a comparison with specific spine group exercise: a service evaluation

Journal of the Royal Army Medical Corps Published Online First: 30 October 2018. doi: 10.1136/jramc-2018-001011

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

General group exercise in low back pain management in a military population: a service evaluation

Journal:	<i>Journal of the Royal Army Medical Corps</i>
Manuscript ID	jramc-2018-001011.R1
Article Type:	Original research
Keywords:	chronic low back pain, group rehabilitation, exercise, service evaluation
Abstract:	<p>Objective: To investigate whether general group exercise (GGE) offers the same outcomes when compared with a specific spinal group exercise (SSGE) for chronic low back pain (CLBP) in a military population.</p> <p>Design: Retrospective service evaluation using routine service activity data.</p> <p>Setting: A UK military rehabilitation centre.</p> <p>Participants: A total of 106 CLBP patients.</p> <p>Interventions: Three-week intensive (five days per week, 15-day intervention) rehabilitation course for patients with CLBP. Six SSGE groups (n= 64); CLBP only. Six GGE groups (n=42); CLBP patients grouped with chronic lower limb injuries.</p> <p>Outcome Measures: Oswestry Disability Index (ODI), Numerical Pain-Rating Scores (NPRS) and the Modified Multi-Stage Fitness Test (Mod-MSFT). Long term effects were measured by Medical Employment Standard (MES) status and physiotherapy follow-up at three and 12 months.</p> <p>Results: A between-group analysis showed no significant difference in GGE compared to SSGE. Mean changes (standard deviation) in pain were -2.71 ± 2.35 and -1.20 ± 1.99 ($p=0.018$), ODI were -3.6 ± 5.7 and -4 ± 8.5 respectively ($p = 0.649$) and Mod-MSFT 28.4 ± 30.8 and 29.7 ± 31.7 respectively ($p = 0.792$). At three months, a greater proportion of the GGE were having on-going physiotherapy; GGE = 50%, SSGE = 30.2%, ($p = 0.016$) although, some differences were evident across MES with 32.5% of GGE compared to 20.6% of SSGE being medically fit with no restrictions. . At 12 months, groups were largely comparable for follow up physiotherapy and MES; 22.5% of GGE and 20.6% of SSGE continued to have physiotherapy input; 47.5% of GGE and 50.8% of SSGE medically fit with no restrictions.</p> <p>Conclusion: Patients with CLBP who completed a 3-week rehabilitation programme had comparable outcomes when grouped with LL patients, although only improvements in pain in the GGE group achieved a meaningful change. Further evaluation of potential costs and savings to service costs are now required.</p>

TITLE**General group exercise in low back pain management in a military population, a comparison with specific spine group exercise: a service evaluation**

Surtees JE, Heneghan NR

Joanna Surtees: Primary Care Rehabilitation Facility, RAF Waddington, Lincoln, LN5 9NB

Nicola Heneghan: Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

Correspondence:

Nicola R Heneghan

Centre of Precision Rehabilitation for Spinal Pain (CPR Spine),

School of Sport, Exercise & Rehabilitation Sciences

University of Birmingham, Edgbaston,

Birmingham. B15 2TT. UK

Email: n.heneghan@bham.ac.uk

Tel: +44 121 415 8367

Key words: chronic low back pain, group therapy, exercise, service evaluation

Word count: 3034

ABSTRACT

Objective: To investigate whether general group exercise (GGE) offers the same outcomes when compared with a specific spinal group exercise (SSGE) for chronic low back pain (CLBP) in a military population.

Design: Retrospective service evaluation using routine service activity data.

Setting: A UK military rehabilitation centre.

Participants: A total of 106 CLBP patients.

Interventions: Three-week intensive (five days per week, 15-day intervention) rehabilitation course for patients with CLBP. Six SSGE groups (n= 64); CLBP only. Six GGE groups (n=42); CLBP patients grouped with chronic lower limb injuries.

Outcome Measures: Oswestry Disability Index (ODI), Numerical Pain-Rating Scores (NPRS) and the Modified Multi-Stage Fitness Test (Mod-MSFT). Long term effects were measured by Medical Employment Standard (MES) status and physiotherapy follow-up at three and 12 months.

Results: A between-group analysis showed no significant difference in GGE compared to SSGE. Mean changes (standard deviation) in pain were -2.71 ± 2.35 and -1.20 ± 1.99 ($p=0.018$), ODI were -3.6 ± 5.7 and -4 ± 8.5 respectively ($p = 0.649$) and Mod-MSFT 28.4 ± 30.8 and 29.7 ± 31.7 respectively ($p = 0.792$). At three months, a greater proportion of the GGE were having on-going physiotherapy; GGE = 50%, SSGE = 30.2%, ($p = 0.016$) although, some differences were evident across MES with 32.5% of GGE compared to 20.6% of SSGE being medically fit with no restrictions. . At 12 months, groups were largely comparable for follow up physiotherapy and MES; 22.5% of GGE and 20.6% of SSGE continued to have physiotherapy input; 47.5% of GGE and 50.8% of SSGE medically fit with no restrictions.

Conclusion: Patients with CLBP who completed a 3-week rehabilitation programme had comparable outcomes when grouped with LL patients, although only improvements in pain in the GGE group achieved a meaningful change. Further evaluation of potential costs and savings to service costs are now required.

INTRODUCTION

As well as the leading cause of disability in the general working age population globally, chronic low back pain (CLBP) is also the most common musculoskeletal presentation in the US and UK armed forces.^{1,2} In 2009 7% of UK armed forces medical discharges were a consequence of low back pain.³ Where exercise is recommended for management of LBP⁴ in the general population, exercise rehabilitation has been identified as a research priority by the UK Defence Directive of Rehabilitation (DDR).⁵

Recent National Institute for Health and Clinical Excellence guidelines recommend group exercise as a management option for CLBP.⁶ Recent systematic reviews (2015 and 2017) comparing group exercise programmes to one-to-one physiotherapy for management of chronic musculoskeletal conditions, including CLBP, reported similar clinical outcomes for improvements in pain intensity and functional disability.^{7,8} Whilst both reviews endorse the use of physiotherapist led group exercise as a cost effective management approach for CLBP, another Cochrane review concluded that no one form of exercise afforded superior outcomes⁹; mirroring earlier review findings from Searle et al.,¹⁰ with evidence of support for a wide variety of exercise interventions, including yoga, Pilates, and strength/resistance training and stability/coordination exercises.¹⁰

1
2
3 Group exercise therapy is a core component in CLBP management within the Defence Medical
4 Rehabilitation Programme (DMRP).^{11,12} Regional Rehabilitation Units (RRU) provide 3-week intensive
5 rehabilitation courses (five days per week, 15-day intervention) for patients with chronic
6 musculoskeletal conditions, including CLBP, with 15 patients enrolled on each course. This long
7 established mode of delivery has historically differentiated upper limb, lower limb (LL) and specific
8 spinal group exercise (SSGE) courses. Patients are referred to the RRU from physiotherapists working
9 at Primary Healthcare Rehabilitation Facilities (PCRf), (Figure 1).
10
11
12
13
14

15 In June 2015, the investigating RRU was unable to fill the SSGE course quota; moreover, LL courses
16 were routinely full. Consequently, LL patients had to wait longer for a LL course, exceeding Key
17 Performance Indicators (KPIs) outlined in the DMRP. Based on the available research evidence at
18 that time, amalgamating the LL and SSGE into a General Group Exercise (GGE) course was a
19 justifiable course of action to afford positive outcomes for patients with CLBP. Additionally, the
20 ability to offer courses more regularly could reduce the socioeconomic burden of CLBP in a military
21 population, hastening return to duties and reducing healthcare usage.
22
23
24
25
26
27

28 From a detailed literature search the authors identified one study that investigated the effectiveness
29 of exclusively LL exercise in the treatment of CLBP. Cai et al.¹³ demonstrated that LL strengthening
30 was equally effective to lumbar extensor or lumbar stabilisation exercises for improving lumbar
31 multifidus muscle activation and superior for running functional outcomes in a recreational running
32 population with CLBP (n=84). No study was identified specifically investigating outcomes in exercise
33 groups comprising both CLBP and LL conditions or in a more representative population. In view of
34 these findings, and in line with the identified DDR research priorities, the aim of this service
35 evaluation (SE) was to evaluate the outcomes of the GGE for CLBP patients compared to existing
36 data for SSGE.
37
38
39
40
41
42
43

44 **METHODS**

45 **Design**

46
47 A retrospective SE was designed using routine service activity data. In the absence of reporting
48 guidelines¹⁴ the Standards for Reporting Implementation Studies (StaRI) document was used to
49 inform the methods of the SE.¹⁵ A priori protocol was developed with expertise from the University
50 of Birmingham and approved by the Academic Department of Military Rehabilitation. Ethical
51 approved was granted by the School of Sport and Exercise Sciences Ethics Committee University of
52 Birmingham, and the Ministry of Defence Research Ethics Committee.
53
54
55
56
57
58
59
60

Inclusion Criteria

All CLBP patients accepted for residential rehabilitation from December 2014 to June 2015 were admitted to six established SSGE courses; patients accepted from December 2015 to June 2016 were admitted to six GGE courses. Inclusion criteria: Army, Royal Air Force (RAF) or Royal Navy (RN) personnel, aged 17 - 55. All patients were seen by a GP and physiotherapist at PCRf and, in 69% of cases, an exercise therapist. All patients had CLBP (>3months)¹⁶ and a diagnosis confirmed by a Sports and Exercise Medicine Physician. For the purpose of this SE, clinical presentations were categorised as: non-specific LBP, radiculopathy, sacroiliac joint, trauma, and post-operative spinal surgery. All potential participants were screened to confirm eligibility for participation in active exercise in a group setting with no medical contraindications e.g. cardiovascular, respiratory, neurological or mental health conditions.

Course Design

There is no published evidence supporting the RRU 3-week residential rehabilitation course as opposed to different models or timescale. Duration of 3-weeks for the programmes is pragmatic, with a need to balance time to deliver a clinical intervention, manage waiting lists and allowing individuals' time away from their unit and primary duties to focus on rehabilitation. The course elements include: *strength training and functional conditioning, sensorimotor training, motor control and dynamic stability training, range of motion, flexibility and general movement, cardiovascular conditioning, hydrotherapy, education (pain, goal setting, anatomy and physiology, diet and nutrition, principles of training and relaxation)*¹⁷

Each course is led by a designated senior physiotherapist and exercise therapist. Exercise prescription, progression and intensity is controlled and monitored by the physiotherapist and exercise therapist, and always conducted in a group environment. In line with normal service delivery one-to-one treatment was available, if required. The main difference between the SSGE and GGE was that unlike the GGE, the SSGE had daily, mat-based, spinal mobility sessions.

Outcome Measures

Pain: Numerical Pain-Rating Score (NPRS). A valid and responsive self-report measure of pain intensity, (0-10) where 0 = no pain and 10 = worst possible pain with values recorded at pre and post rehabilitation course.¹⁸

Disability: Oswestry Disability Index (ODI). The ODI is a back specific patient reported questionnaire, consisting of 10 questions that assess the level of pain interference with physical activities of daily living.¹⁹ Test retest reliability is reported to be excellent ICC 0.88 (95%CI 0.77-0.94) and ICC 0.94 (95%CI 0.89-0.97).^{20 21}

Long term outcomes of the intervention were assessed using individual's **Medical Employment Standard (MES)** status at three and 12 months. MES categories are listed below²² although this SE had a specific focus on medical fitness with no restrictions (P2):

P0	Medically unfit for duty and under medical care
P2	Medically fit for unrestricted service worldwide
P3	Medically fit for duty with minor employment limitations
P4	Medically fit for duty within the limitations of pregnancy
P7	Medically fit for duty with major employment limitations

Fitness: Modified multi-stage fitness test (Mod-MSFT). A measure physical function, the Mod-MSFT is a modification of the established Multi Stage Fitness Test (MSFT).²³ It was first used with traumatic brain injuries demonstrating excellent reliability and validity.^{24 25} Markers are placed at 0, 10 and 20metres where the test involves walking, and then running, the 20 metre distance in time to a shortening frequency of beeps, played out on an audio device. The test is terminated by the patient due to pain or fatigue and has been used in a CLBP military population.²⁶ The MSFT is used by the RAF and RN as measurement of physical fitness; achieving an age and sex appropriate pass mark is essential to achieve medical fitness with no restrictions (P2) MES.^{22 27}

Healthcare use including ongoing physiotherapy interventions was also evaluated at three and 12 months.²⁸

Procedure

ODI and Mod-MSFT were recorded by the individual course physiotherapist at the start and end of each course. The researcher extracted baseline demographic characteristics and all outcome measure data from a manual search of electronic defence medical records of all individuals participating in the six SSGE and six GGE courses.

Data analysis

Data was analysed using primarily descriptive methods, utilising the statistical analysis software SPSS version 21 (SPSS Inc., Chicago, Illinois). The alpha level was set at 0.05. Prior to statistical analysis, the Shapiro-Wilks test for normality was utilised due to the small sample size. As a result, the Mann-Whitney U test was selected as an appropriate non-parametric test.²⁹

RESULTS

A total of 106 CLBP patients met the inclusion criteria and were included in the evaluation. The personal characteristics of participants are presented in Table 1.

Table 1 shows there was no between-group statistical significant difference for the following characteristics: age ($p = 0.864$), waiting time ($p = 0.864$) or male/female ratio ($p = 0.170$). The most common clinical diagnosis was non-specific CLBP; 73 of the 106 sample; SSGE 70.3%, GGE 65.9%. There was no statistically significant difference between groups in the clinical presentation ($p = 0.413$). The frequency of non-specific LBP was lower than the commonly reported 90% of all presentations of LBP.³⁰ Waiting time was measured from first presentation at PCRF to the first day of the course. The most common LL presentations within the GGE were post-op anterior cruciate ligament reconstructions, anterior knee pain and hip pain.

		SSGE (n=64)	GGE (n=42)	p value
Age (years)		33.3 ± 7.19	33.9 ± 7.9	0.864*
Waiting Time (days)		277.8 ± 220.6	272.6 ± 241.6	0.864*
Male / Female Ratio		76.6% / 23.4%	64.3% / 35.7%	0.170**
No. of Spine / LL patients on each course (% spinal)	Course 1	12	7 / 6 (53.8%)	n/a
	Course 2	13	5 / 10 (33.3%)	
	Course 3	9	10 / 5 (66.7%)	
	Course 4	7	4 / 9 (30.8%)	
	Course 5	13	9 / 6 (60%)	
	Course 6	10	7 / 8 (46.7%)	

Data presented as mean ± SD *Mann-Whitney **Chi-squared test

Table 1. Characteristics of participants in SSGE and GGE

A between-group analysis, summarised in Table 2, showed no significant difference in the GGE outcomes when compared with the SSGE group. Pain mean change was -2.71 and -1.20 ($p=0.018$), ODI mean change -3.6±5.7 and -4±8.5 respectively ($p=0.649$) and mod-MSFT mean change 28.4±30.8 and 29.7±31.7 respectively, ($p=0.792$).

	SSGE (n=63)	GGE (n=40)	p value
Pain			

Mean NPRS pre course	4.63 ± 2.31	5.17 ± 2.00	0.800
Mean NPRS post course	3.33 ± 2.32	1.81 ± 2.14	0.907
Mean NPRS change	-1.20 ± 1.99	-2.71 ± 2.35	0.018
Disability			
Mean ODI pre course	21.9 ± 11.4	22.7 ± 9.8	0.761
Mean ODI post course	17.6 ± 11.1	19.4 ± 10	0.322
Mean ODI change	-4 ± 8.5	-3.6 ± 5.7	0.649
	SSGE (n=60)	GGE (n=40)	
Physical Function			
Mean Mod-MSFT pre course*	141.8 ± 51.7	138.4 ± 63.2	0.867
Mean Mod-MSFT post course*	171.5 ± 54.4	166.7 ± 58.1	0.725
Mean Mod-MSFT change*	29.7 ± 31.7	28.4 ± 30.8	0.792

Data presented as mean ± SD; SD: Standard Deviation; ODI: Oswestry disability Index; Mod-MSFT: Modified Multistage fitness test; *no of individual shuttles of 10metres

Table 2. Pre and post intervention measures for pain, disability and physical function

Physiotherapy and functional status at 3 and 12 month follow up

At 3 months a greater proportion of the GGE group were still having on-going physiotherapy care (50%) compared with 30.2% in SSGE, although more of the GGE were medically fit with no employment restrictions (32.5%) compared to 20.6% of the SSGE group. Notwithstanding some differences across categories of MES, at 12 months, groups were largely comparable with 22.5% of the GGE and 20.6% of the SSGE continuing to have physiotherapy and 47.5% of GGE and 50.8% of SSGE deemed medically fit with no employment restrictions (see Table 3).

		SSGE (n=63)	GGE (n=40)
Percentage of physio follow up at 3 and 12 months			
3 month post course	Ongoing physio	30.2%	50%
	Discharged Care Complete	41.3%	15%
	Admin Discharge	28.5%	35%
12 months post course	Ongoing physio	20.6%	22.5%
	Discharged Care Complete	54%	40%
	Admin Discharge	25.4%	37.5%
Percentage of post intervention employment standard at 3 and 12 months			

3 month post course	P2	20.6%	32.5%
	P3	28.6%	10%
	P7	47.6%	55%
	P4	0%	0%
	P0	3.2%	2.5%
12 months post course	P2	50.8%	47.5%
	P3	22.2%	20%
	P7	23.8%	32.5%
	P4	1.6%	0%
	P0	1.6%	0%

Table 3. Physiotherapy and functional status at 3 and 12 month follow up

Missing data

Full data sets were available for ODI evaluation, although 3 values were missing for post course Mod-MSFT SSGE, and 11 for NPRS pre and post SSGE, 13 pre and 19 post GGE.

DISCUSSION

The aim of this SE, the first of its kind was to examine outcomes in CLBP patients completing a 3-week course when grouped with LL patients, compared to a group of CLBP patients only. Given the inherent difficulties of conducting clinical trials in a military setting, where participants may be deployed or posted at short notice, use of SE offers an alternative approach to evaluate practice and implement changes in a timely manner. Additionally, this offers a means to systematically assess activities and outcomes to examine efficiency and effectiveness of a service.³¹ Whilst cost effectiveness is a key driver a new multi-criteria decision analysis model incorporates a more comprehensive evaluation inclusive of access, equity, effectiveness of treatment and impact on future services.³² This SE therefore provides a robust evaluation of the impact of changes in patient outcomes in a military setting. In summary this SE found more than 2 point difference in pain scores in favour of GGE, although no between-group difference was found with respect to disability or physical function. A greater proportion of the GGE were still having on-going physiotherapy care at 3 months although 32.5% of the GGE were medically fit with no restrictions compared to only 20.6%

of the SSGE. At 12 months, groups were largely comparable with respect to follow up physiotherapy and those who were deemed medically fit with no or minor employment restrictions.

Measures

Selection, administration and interpretation of outcome measures are important facets in evaluation³³. In this SE outcome measures were informed DDR policy, with 4 of the 5 well established areas for measuring outcomes in LBP included; disability, back specific function, generic health status pain, and work.³⁴

Pain

Notwithstanding the extent of the missing data for NPRS change scores achieved those reported in the wider literature for MCID.¹⁸ It is interesting that this sizeable change was observed during the 3-week course which would suggest that the non-physical factors such as beliefs, knowledge *etc.* had a role in pain perception; reflective of the multidimensional nature of LBP.³⁵ Caution should be taken when interpreting these findings given the extent of the missing data for pain.

Disability

Based on the reported requirements for the general population neither group achieved a meaningful change in disability scores. However, where both groups were largely of minimal disability based on ODI and the absence of a population-specific measure of disability, where a sensitive and specific has not yet been established, groups did meet the values required at the lower end of the range for the reported MCID from other populations, ranging 2.92 to 15.36³⁶ to a 10 point change combined with a 30% improvement from baseline³⁷

Fitness

The only MCID documented in the literature for a shuttle-based test is the Shuttle Walking Test.³⁸ The authors found a change of 76m would be required to represent a 95% confidence interval. However, this was in a population of 29 patients with a diagnosis of spinal stenosis, with a mean age of 69; notably higher than the mean age reported here. Moreover, ceiling effects of this test were found³⁹; out of a total of 90 patients, 31 had achieved 11 of the 12 levels at baseline assessment. Furthermore, it has a different format to the Mod-MSFT being conducted over a 10m not 20m distance.

The only comparable study using the Mod-MSFT in a CLBP military population involved 56 subjects completing an equivalent 3-week course at the Defence Medical Rehabilitation Centre (DMRC).

1
2
3 Roberts et al.²⁶ reported a mean change of 120m, considerably less than distances in this SE; 284m
4 (GGE) and 297m (SSGE). Given their lower mean pre-course and post-course distances of 1040m and
5 1160m respectively this suggests a lower functioning group compared to this SE (GGE = 1715m;
6 SSGE= 1667m). As a tertiary tier of the care pathway (Figure 1) these findings for DMRC are not
7 unexpected. In the absence of other published data the mean post-course scores achieved by both
8 groups is 500m below the pass mark required for males of a comparable age in the RAF and RN
9 MSFT,²⁷ suggesting that full MES was still not achieved by the end of the course.

10
11
12
13
14
15 MES was used to evaluate work disability rather than the Functional Assessment Tool (FAA) reported
16 elsewhere in the literature which limits the ability to draw comparison with this current service
17 evaluation.⁴⁰ MES selection was chosen primarily due to the known problem of clinicians assessing
18 FAA in practice, rather than the patient.⁴¹ That said the MES correlates well to the FAA and critically
19 has been linked to military operational effectiveness.

20 21 22 23 24 **Implications for practice and policy**

25
26 It has been documented that prompt recovery (in non-specific LBP) is most likely to occur during a
27 three months post-onset, with only gradual improvements thereafter.^{42 43} Moreover, studies have
28 found that 62% of all patients continued to complain of pain at 12 months.⁴⁴ This raises the question
29 as to what we can realistically expect given patients commenced the course, on average, 8+ months
30 from initial presentation. Only 69% of patients saw an Exercise Therapist at PCRf pre-course; one of
31 their primary functions is group exercise. There may be greater potential for improvements if
32 rehabilitation courses are offered earlier in the care pathway, and which may in turn ameliorate
33 some of the cost burden of managing more established chronic pain presentations.

34
35
36
37
38
39
40 With a lack of comparable data for those individuals who did not attend an RRU course; this study
41 raises the question of whether the right patients are being selected for course participation as part
42 of the DMRP tier approach. Despite DMRP referral guidelines and timescales, referral patterns to the
43 RRU are patient- and therapist- dependant, informed by therapist expertise, patient operational
44 demands and clinical presentation. This may also explain the variability in waiting times seen in this
45 SE. With a review citing 1501 potential prognostic factors associated with poor recovery from LBP⁴³,
46 decision making for patient referral is complex. With the recent introduction of the STarTBack tool
47 into the DMRP this may now better differentiate different presentations of LBP and inform targeted
48 management.^{11 45}

49
50
51
52
53
54 Maher⁴⁶ summaries the challenges clinicians face where no single treatment cures CLBP, and the
55 abundant unregulated, non-evidence based management options that bombard patients confuse
56
57

1
2
3 the issue further. Whilst there is no specific evidence for the 3-week model, elements found within
4 the course are well evidenced. This SE goes some way to justify the need for more research into the
5 modes of rehabilitation delivery in the UK military setting to assist with clinical decision making.
6
7

8 9 **Strengths and limitations**

10 One of the main limitations is that pain, ODI and Mod-MSFT data was not available at the three or
11 12-month follow up points. The habitual use of outcome measures in clinical practice has challenges,
12 and is widely reported in the literature.⁴⁷ The lack of routine outcome measures recorded across the
13 DMRP limits the impact of the findings of this SE and warrants further investigation. Moreover,
14 population specific measures with established measurement properties are required to further
15 inform practice decisions.⁴⁸ This SE has highlighted the inconsistent recording of the numerical pain
16 rating scale¹⁸ pre- and post-course, despite being an outcome measure documented in DDR policy.
17 Finally evaluation of the impact of a GGE course on LL patients was beyond the scope of this SE,
18 although could be useful to strengthen proposed service changes.
19
20
21
22
23
24
25

26 **CONCLUSION**

27
28 Patients with CLBP who completed a 3-week rehabilitation programme had comparable outcomes
29 when grouped with LL patients, although only improvements in pain in the GGE group achieved
30 more than the MCID on completion of the course. At 12 month follow up both groups were largely
31 comparable with respect to achieving medical fitness with no or minor employment restrictions. This
32 service evaluation supports the need to further consider timing for rehabilitation in the care
33 pathway, comprehensive use of patient reported outcomes and further evaluation of potential costs
34 and savings to service costs.
35
36
37
38
39
40
41

42 **Legend: Figure 1.** Summary of Defence Medical Rehabilitation Centre model
43
44
45
46

47 **Clinical messages**

- 48 • Comparable outcomes are achieved if CLBP patients complete a 3-week exercise programme
49 when grouped with LL patients, compared to a group comprised of CLBP patients only.
- 50 • Efficiency savings could be made by grouping mixed military musculoskeletal presentations
51 together for group exercise therapy earlier in the care pathway.
52
53
54
55
56
57

Conflict of interest

The author declares that there are no known conflicts of interests.

Funding

The author received no specific grant from any funding agency in the commercial, public or not-for-profit sectors.

Words: 2953 (excluding tables and figures)

REFERENCES

1. Knox J, Orchowski J, Scher DL, et al. The incidence of low back pain in active duty United States military service members. *Spine* 2011;**36**(18):1492-500.
2. Strowbridge NF, Burgess KR. Sports and training injuries in British soldiers: the Colchester Garrison Sports Injury and Rehabilitation Centre. *J R Army Med Corps* 2002;**148**(3):236-43.
3. Defence Analytical Services Agency. Secondary Defence Analytical Services Agency. <http://webarchive.nationalarchives.gov.uk/20130403160018/http://www.dasa.mod.uk/modintranet/UKDS/UKDS2010/ukds.php>.
4. National Institute for Health and Care Excellence (NICE) guidelines. Neck pain - non-specific. Secondary Neck pain - non-specific 2015. <https://cks.nice.org.uk/neck-pain-non-specific>.
5. Department AR. Emerging Themes and Research Questions Academic Department of Military Research. RCG Research Workshop DMRC Headley Court, 2014.
6. National Institute for Health and Care Excellence (NICE) guidelines. Low back pain and sciatica in over 16s: assessment and management. Secondary Low back pain and sciatica in over 16s: assessment and management 2016. <https://www.nice.org.uk/guidance/ng59>.
7. O'Keefe M, Hayes A, McCreesh K, et al. Are group-based and individual physiotherapy exercise programmes equally effective for musculoskeletal conditions? A systematic review and meta-analysis. *British Journal of Sports Medicine* 2017;**51**:126-32.
8. Toomey E, Currie-Murphy L, Matthews J, et al. The effectiveness of physiotherapist-delivered group education and exercise interventions to promote self-management for people with osteoarthritis and chronic low back pain: A rapid review Part I. *Man Ther* 2015;**20**:265-86.
9. Saragiotto BT, Maher CG, Yamato TP, et al. Motor control exercise for chronic non-specific low-back pain (Review) *Cochrane Db Syst Rev*, 2016.
10. Searle A, Spink M, Ho A, et al. Exercise interventions for the treatment of chronic low back pain: a systematic review and meta-analysis of randomised controlled trials. *Clinical rehabilitation* 2015;**29**(12):1155-67.
11. Defence Mo. JSP 950: Chapter 22: Rheumatology and Rehabilitation. Leaflet 2-22-1 Defence Medical Rehabilitation Programme (DMRP). Available from: MOD Intranet, 2010.
12. Roberts AJ, Dew A, Bridger R, et al. Predicting low back pain outcome following rehabilitation for low back pain *J Back Musculoskelet Rehabil* 2015;**28**(1):119-29.
13. Comparison of 3 exercise therapies for recreational runners with chronic low back pain. 34 International Conference of Biomechanics in Sport 2016; Tsukuba, Japan.
14. Altman DG, Simera I, Hoey J, et al. EQUATOR: reporting guidelines for health research. *Lancet* 2008;**371**(9619):1149-50.
15. Pinnock H, Barwick M, Carpenter CR, et al. Standards for reporting implementation studies (StaRI): explanation and elaboration document. *BMJ Open* 2017;**356**(i6795).
16. Charlton JE. *Core Curriculum for Professional Education in Pain*. 3rd ed. Seattle, USA, 2005.
17. Lawson E. Regional Rehabilitation Unit - Spines rehabilitation course standardisation. Tidworth Garrison 2017.
18. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine* 2005;**30**(11):1331-34.
19. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine* 2000;**25**(22):2940-52.
20. Grotle M, Garratt AM, Jenssen HK, et al. Reliability and construct validity of self-report questionnaires for patients with pelvic girdle pain. *Phys Ther* 2012;**92**(1):111-23.
21. Dawson AP, Steele EJ, Hodges PW, et al. Utility of the Oswestry Disability Index for studies of back pain related disability in nurses: evaluation of psychometric and measurement properties. *Int J Nurs Stud* 2010;**47**(5):604-07.
22. Defence Mo. JSP 950: Chapter 6: Joint service manual of medical fitness Leaflet 6-7-7 MOD Intranet, 2016.

23. Lamb KL, Rogers L. A re-appraisal of the reliability of the 20 m multi-stage shuttle run test. *Eur J Appl Physiol* 2001;**100**(3):287-92.
24. Vitale AE, Jankowski LW, Sullivan SJ. Reliability for a walk/run test to estimate aerobic capacity in a brain-injured population. *Brain Injury* 1997;**11**(1):67-76.
25. Hassett LM, Harmer AR, Moseley AM, et al. Validity of the modified 20-metre shuttle test: assessment of cardiorespiratory fitness in people who have sustained a traumatic brain injury. *Brain Injury* 2007;**21**:1069-77.
26. Roberts A, Seah R, Dickens JCW, et al. A comparison of pain levels after the Biering-Sorensen and the modified 20-metre shuttle test in patients with chronic low back pain. *J Back Musculoskelet Rehabil* 2014;**27**(2):173-79.
27. Defence Mo. Ministry of Defence. AP3342: RAF Fitness Test, Section 4, Leaflet 402 MOD Intranet.
28. Department for International Development FCO. National Security Strategy and Strategic Defence and Security Review 2015.
29. Field A. *Discovering Statistics with IBM SPSS*. Newbury Park, CA: Sage, 2013.
30. Koes BW, van Tulder MW, Thomas S. Diagnosis and treatment of low back pain. *BMJ* 2006;**332**(7555):1430-4.
31. Authority HR. Defining research. 2016 Secondary Defining research. 2016 2016. <http://www.hra.nhs.uk/research-community/before-you-apply/determine-whether-your-study-is-research/>
32. Dionne F, Mitton C, MacDonald T, et al. The challenge of obtaining information necessary for multi-criteria decision analysis implementation: the case of physiotherapy services in Canada. *Cost Effectiveness and Resource Allocation* 2013;**11**(11).
33. Walton DM. Making (common) sense of outcome measures. *Man Ther* 2015;**20**:723-26.
34. Bombardier C. Outcome assessments in the evaluation of treatment of spinal disorders: summary and general recommendations. *Spine* 2000;**25**:3100-3.
35. Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *The Lancet*.
36. Johnsen LG, Hellum C, Nygaard OP, et al. Comparison of the SF6D, the EQ5D, and the Oswestry disability index in patients with chronic low back pain and degenerative disc disease. *BMC Musculoskelet Disord* 2013;**14**:148-57.
37. Ostelo RWJG, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine* 2008;**33**(1):90-4.
38. Pratt RK, Fairbank JCT, Virr A. The Reliability of the Shuttle Walking Test, the Swiss Spinal Stenosis Questionnaire, the Oxford Spinal Stenosis Score, and the Oswestry Disability Index in the Assessment of Patients With Lumbar Spinal Stenosis. *Spine* 2002;**27**(1):84-91.
39. Chown M, Whittamore L, Rush M, et al. A prospective study of patients with chronic back pain randomised to group exercise, physiotherapy or osteopathy. *Physiotherapy* 2008;**94**(1):21-28.
40. Roberts AJ, Franklyn-Miller AD, Etherington J. A new functional outcome assessment tool for military musculoskeletal rehabilitation: a pilot validation study. *PM R* 2011;**3**(6):527-32.
41. Roberts AJ, Etherington J. The functional activity assessment: a validated PROM, unreliable in the hands of clinicians. *J R Army Med Corps* 2013;**159**(4):287-90.
42. Pengel LH. Acute low back pain: systematic review of its prognosis. *BMJ* 2003;**327**(7410):1-5.
43. Kent PM, Keating JL. Can we predict poor recovery from recent-onset nonspecific low back pain? A systematic review. *Man Ther* 2008;**13**(1):12-28.
44. Hestbaek L L-YC, Manniche C. Low back pain: what is the long-term course? A review of studies of general patient populations. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society* 2003;**12**(2):149-65.

- 1
- 2
- 3 45. Kongsted A, Andersen CH, Hansen MM, et al. Prediction of outcome in patients with low back
- 4 pain – A prospective cohort study comparing clinicians' predictions with those of the Start
- 5 Back Tool. *Man Ther* 2016;**21**:120-27.
- 6 46. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet* 2017;**389**(10070):736
- 7 - 47.
- 8 47. Colquhoun HL, Lamontagne ME, Duncan EAS, et al. A systematic review of interventions to
- 9 increase the use of standardized outcome measures by rehabilitation professionals. *Clinical*
- 10 *rehabilitation* 2017;**31**(3):299-309.
- 11 48. Kyte DG CM, van der Wees PJ, ten Hove R, Tolan S, Hill JC, . An introduction to patient-reported
- 12 outcome measures (PROMs) in physiotherapy. *Physiotherapy* 2015;**101**(2):119-25.
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



30x22mm (300 x 300 DPI)

review only