

Association between periodontal health status and patient-reported outcomes in patients managed in a non-specialist, general dental practice

Sharma, Praveen; Yonel, Zehra; Chapple, Iain; Dietrich, Thomas

DOI:

[10.1111/jcpe.13022](https://doi.org/10.1111/jcpe.13022)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Sharma, P, Yonel, Z, Chapple, I & Dietrich, T 2018, 'Association between periodontal health status and patient-reported outcomes in patients managed in a non-specialist, general dental practice', *Journal of Clinical Periodontology*. <https://doi.org/10.1111/jcpe.13022>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Sharma P, Yonel Z, Busby M, Chapple IL, Dietrich T. Association between periodontal health status and patientreported outcomes in patients managed in a nonspecialist, general dental practice. *J Clin Periodontol*. 2018, which has been published in final form at: <https://doi.org/10.1111/jcpe.13022>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Association between periodontal health status and patient-reported outcomes in patients managed in a non-specialist, general dental practice

Sharma, Praveen; Yonel, Zehra; Chapple, Iain; Dietrich, Thomas

DOI:

[10.1111/jcpe.13022](https://doi.org/10.1111/jcpe.13022)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Sharma, P, Yonel, Z, Chapple, I & Dietrich, T 2018, 'Association between periodontal health status and patient-reported outcomes in patients managed in a non-specialist, general dental practice' *Journal of Clinical Periodontology*. DOI: 10.1111/jcpe.13022

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Association between periodontal health status and patient reported outcomes in patients managed in a non-specialist, general dental practice

Journal:	<i>Journal of Clinical Periodontology</i>
Manuscript ID	CPE-08-18-7877.R1
Manuscript Type:	Epidemiology (Cohort study or case-control study)
Date Submitted by the Author:	02-Oct-2018
Complete List of Authors:	Sharma, Praveen; University of Birmingham, School of Dentistry Yonel, Zehra; University of Birmingham, School of Dentistry Busby, Mike; Simply Health, Dental Division Chapple, Iain; The University of Birmingham, Periodontal Research Group; Dietrich, Thomas; The School of Dentistry, University of Birmingham, Oral Surgery
Topic:	Treatment
Keywords:	Periodontal disease, Patient reported outcomes, Dental Practice
Main Methodology:	Epidemiology

SCHOLARONE™
Manuscripts

1
2
3 Title: Association between periodontal health status and patient reported outcomes in patients
4 managed in a non-specialist, general dental practice
5
6

7 Key words: Periodontitis, patient reported outcomes, dental practice
8
9

10
11 Authors

12 Praveen Sharma (1), Zehra Yonel (1), Michael Busby (2), Iain L Chapple (1), Thomas Dietrich (1)
13
14

15 (1) School of Dentistry, College of Medical & Dental Sciences, University of Birmingham

16 (2) Dental Advisor Denplan, The Stables, Heritage Ct, Clifton Reynes, Olney, MK46 5FW
17
18

19
20 Corresponding Author

21 Mr Praveen Sharma BDS, MJDF (RCS Eng.), FHEA

22 NIHR Doctoral Research Fellow

23 Clinical Lecturer in Restorative Dentistry

24 University of Birmingham School of Dentistry

25 Birmingham, UK.

26 0121 466 5128
27
28

29 praveen.sharma@nhs.net
30
31

32 Conflict of Interest

33
34 Mr Michael Busby acts as a Denplan advisor and Prof Iain Chapple acts as an advisor to Oral Health
35 Innovations, the license holder for DEPPA in the UK and Ireland.
36

37 Praveen Sharma is funded by a doctoral research fellowship grant by the National Institute of Health
38 Research (NIHR), UK (grant reference: DRF-2014-07-109). The views expressed in this article are
39 those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract:

Aims: To explore the associations between periodontal status and patient reported outcomes (PROs) in a large cohort of patients based in non-specialist general dental practice.

Materials and methods: Analysis was conducted using data from 14,620 patients, in 233 non-specialist dental practices across the UK. As part of routine clinical care, data on periodontal probing depths (PPD), alveolar bone loss (ABL), bleeding on probing (BoP) as well as PROs (oral pain/discomfort, dietary restrictions and dental appearance) were recorded using an online database. The associations between periodontal status and PROs were investigated using logistic regression analysis, adjusting for confounders.

Results: We found a positive association between worse periodontal health and the prevalence of PROs. After adjustment for confounders, 13.8% of patients in the healthiest category (PPD <5mm, ABL <2mm, no BoP) reported pain/discomfort, compared to 20.7% of patients in the worst category (PPD >7mm, ABL >4mm). A similar trend was seen with reporting a restricted diet and unhappiness with appearance.

Conclusion: This study provides novel insights into the associations between periodontal status and PROs in a non-specialist, general dental practice, highlighting the benefits of prevention and management of periodontitis.

Clinical Relevance

Scientific rationale for study

To date, no studies have sought to investigate the relationship between patient reported outcomes in a non-specialist dental setting.

Principal findings

The probability of reporting pain or restrictions in diet or unhappiness with dental appearance increases with worsening periodontal status in a dose-dependent manner. Patients with alveolar bone loss but no deep periodontal pockets are less likely to report dental pain or restrictions in diet compared to patients with deep periodontal pockets.

Practical implications

This study highlights the impact of periodontitis on the well-being of patients and hints at the potential improvements in well-being that may be obtained by periodontal intervention.

Acknowledgements:

The authors would like to thank the patients and dentists contributing to the DEPPA database.

Introduction:

Patient reported outcomes (PROs) are defined as “any report of the status of a patient’s health condition that comes directly from the patient, without interpretation of the patient’s response by a clinician or anyone else” (U. S. Department of Health and Human Services, 2009, U.S. Department of Health and Human Services). PROs are a way for the patient’s perception of their disease or health to be incorporated into their clinical assessment and by doing so, ensuring that all dimensions of health are incorporated in the diagnosis and care for the patient (WHO, 2018).

The WHO defines oral health as a state of being free from diseases that “limit an individual’s capacity in biting, chewing, smiling, speaking, and psychosocial wellbeing” (WHO, 2012). However, often the classification of oral health or oral disease is based on assessments carried out by the dental practitioner with little input from the patient. In the field of periodontology, disease status and treatment outcomes are measured using clinical measures such as periodontal probing depths (PPDs), clinical attachment loss (CAL), bleeding on probing (BOP) and radiographic bone loss. Clearly, these parameters may not be directly relevant to patients, who are thought not to be affected by their periodontal condition for many years before symptoms such as pain or tooth mobility present. Hence, periodontitis has widely been regarded as a “silent disease”. However, this notion has recently been called into question, as several studies have suggested that periodontitis patients report worse oral health related quality of life (Buset et al., 2016).

The aim of the present study was to explore the associations between clinical and radiographic periodontal parameters and patient reported experience of oral pain, dietary restrictions and dental appearance in a large, non-specialist dental practice patient cohort.

Methods:

Patient population

This study used data of 14,620 dentate patients, collected as part of routine clinical care by 355 dentists in 233 non-specialist dental practices across the UK. The dentists were part of a dental payment plan scheme (Denplan), which enables them to charge their patients a fixed monthly fee that covers regular examinations and treatments (Busby et al., 2014a). Participating dental practices comply with a range of key performance/quality outcome measures, deemed consistent with “optimal” dental service provision (Busby et al., 2013). Beginning in 2013, dentists enrolled in the scheme have used an online tool, the Denplan PreViser Patient Assessment (DEPPA). The use of DEPPA is free to dentists who are Denplan Excel Certified, which is a voluntary quality assurance programme. Hence, the vast majority of DEPPA users are Denplan Excel Certified. The DEPPA system is used by participating dentists to record data on a patient’s demographics, risk factors for oral diseases and clinical and radiographic findings. In addition, DEPPA also contains a short patient questionnaire that captures data on the patient’s perceptions of their oral health and behaviours. For the present analyses, data of the first data entry for each patient were used, which may represent either an initial consultation for a patient new to the practice or a check-up appointment for a previously seen patient. These two types of appointment cannot be distinguished from the data available. While the DEPPA system uses these data to derive a number of scores as previously reported (Busby et al., 2013, Busby et al., 2014a., Busby et al., 2014b, Newton and Asimakopoulou, 2017, Sharma et al., 2016), only the raw data entered by clinicians were used for the analyses described in this paper.

All personal identifiers are anonymised and collected in an encrypted format and the system is used as part of routine clinical care. The UK Data Commissioner has confirmed that collected data is non-

1
2
3 personalised. Ethical review was therefore not required for this analysis. As data were not recorded
4 for research purposes, no formal calibration and no standardisation of clinical or radiographic
5 procedures was performed.
6
7

8 *Periodontal/exposure variables*

9
10 As part of the clinical examination, dentists recorded the deepest periodontal probing depth (PPD)
11 per sextant (<5mm, 5-7mm or >7mm), any bleeding on probing (BoP) in each sextant (yes/no) and
12 the maximum radiographic alveolar bone loss (ABL) per sextant (<2mm, 2-4mm, >4mm) as measured
13 from the cemento-enamel junction to the alveolar crest, from available radiographs. Third molars
14 were excluded from the periodontal assessment.
15
16
17

18 *Patient reported outcome variables*

19
20 PROs were assessed on a 3-point Likert scale using questions on pain ("Are you experiencing any
21 pain or discomfort in your mouth?" [Yes/Some/No]), dietary restriction ("Do your teeth allow you to
22 eat an unrestricted diet?" [Yes/Mainly/No]) and appearance ("How do you feel about the
23 appearance of your teeth?" [Happy/Some concerns/Unhappy]).
24
25
26

27 *Other data*

28
29 Data were collected on the following variables based on patients' self-report: age in years and sex
30 (male/female); smoking status (ever/never smoker); diabetes status (yes/no); tooth grinding habits
31 (yes/no) and frequency of sugar intake (less than 4 times/ 4 or more times in a typical day). Data on
32 numbers of teeth with restorations and number of teeth needing restorations; patient's oral hygiene
33 (adequate/inadequate); patient's dental attendance not as regularly as advised by the dentist
34 (yes/no); presence of cervical tooth-wear as a measure of abrasion (yes/no) and salivary flow
35 (adequate/inadequate) were collected, as judged by the clinician.
36
37

38 *Statistical analysis*

39
40 The following statistical analysis plan was defined *a priori*. To avoid sparse strata and facilitate
41 interpretability, binary PRO variables were generated for pain (yes/some vs. no), dietary restriction
42 (yes vs. mainly/no) and appearance (happy vs. some concerns/unhappy). In addition, we created
43 binary outcome variables based on the number of positive PRO responses (at least one vs. none and
44 more than one vs. one or none).
45

46 Based on the worst sextant score of PPD, BoP and ABL, patients were categorised into 8 different
47 periodontal status (exposure) categories (Table 1).
48

49 Logistic regression analysis was used to investigate the associations between each PRO as the
50 outcome variable and the various categories of periodontal parameters as independent variables. In
51 multivariable analyses, adjustments were made for variables which might confound these
52 associations. These included age, sex, smoking and diabetes status, number of teeth with
53 restorations, number of teeth needing restorations, oral hygiene, dental attendance, abrasion,
54 grinding habits and frequency of sugar intake. In addition, salivary flow was included as a covariate
55 in the association between periodontal health status and dental pain and diet and but not for the
56 association between periodontal health and appearance.
57
58
59
60

1
2
3 The adjusted probabilities and two-sided 95% CI of reporting various outcomes were calculated by
4 periodontal status. We carried out a sensitivity analysis by investigating these associations in a sub-
5 set of patients with no outstanding restorative needs.

6
7 To assess whether the reporting of oral pain, dietary restriction and unhappiness with appearance
8 were clustered in a sub-group of patients, we investigated the correlation between reporting pain
9 and restricted diet, reporting restricted diet and unhappiness with appearance and reporting
10 unhappiness with appearance and pain using tetrachoric correlation coefficients.

11
12 Additional sensitivity analyses were conducted including comparisons of findings of the above
13 associations between patients requiring no restorations and patients requiring at least one
14 restoration. In addition, a sensitivity analysis was conducted omitting the variables "oral hygiene",
15 "dental attendance" and "salivary flow".

16
17 Patients with missing data were not included in the analysis.

18
19
20
21 Results:

22 *Sample characteristics*

23
24 The final analytical sample consisted of 14,568 dentate patients, following the exclusion of 52
25 patients (0.003%) with missing data. The mean age of patients in this cohort was 55.5 years (SD 15.7,
26 range 17-106), 6,280 (43%) were male, 830 (5.7%) reported being diagnosed with diabetes and
27 9,146 (63%) were never smokers. The mean number of teeth present, not including wisdom teeth,
28 was 25 (SD 4, range 1-28). Of these, a mean of 11 (SD 5.7, range 0-28) were restored and 0.3 (SD 1,
29 range 0-26) needed restorations. 696 (5%) did not attend the dentist as regularly as recommended,
30 6,469 (44%) had inadequate oral hygiene and 1,656 (11%) reported a high frequency of sugar intake
31 (Table 1).

32
33
34 With regards to their periodontal parameters, 2,693 (18.5%) were in the healthiest group, 3,081
35 (21.2%) had BoP with no or limited periodontal tissue loss, with the remainder exhibiting various
36 levels of periodontal tissue loss evidenced by PPD 5+mm and/or ABL 2+mm with or without bleeding
37 on probing (Table 1).

38 *Unadjusted prevalence of PROs*

39
40
41 2.7% of patients reported experiencing oral pain or discomfort and 12.3% reported experiencing
42 some oral pain or discomfort; 3.5% of patients reported a restricted diet and 7.8% reported a
43 somewhat restricted diet; 2.7% of patients reported being unhappy with the appearance of their
44 teeth and 23.5% had some concerns with the appearance of their teeth. In addition, 40% of the
45 cohort reported at least one of these concerns and 11% reported more than one of these concerns
46 (Table 1). Patients' demographics, lifestyle factors and number of teeth with restorations, number of
47 teeth needing restorations, oral hygiene, dental attendance, abrasion, grinding habits and frequency
48 of sugar intake, were very similar regardless of whether they reported oral pain, diet restrictions or
49 being unhappy with the appearance of their teeth (data not shown).

50
51
52 There were statistically significant correlations between the PROs with tetrachoric correlation
53 coefficients being 0.3514 ($p < 0.0001$), 0.3110 ($p < 0.0001$) and 0.2940 ($p < 0.0001$) for the correlations
54 between reporting pain and restricted diet, restricted diet and unhappiness with appearance and
55 unhappiness with appearance and pain, respectively.

Adjusted prevalence of PROs

Overall, the prevalence of all reported PROs tended to increase with worsening periodontal parameters (Tables 1&2, Figures 1-5).

Specifically, the adjusted prevalence estimates, i.e., predicted probabilities from multivariable logistic models, for pain ranged from 13.8% (95% CI 12.4-15.2%) in the periodontally healthiest group to 20.7% (95% CI 17.2-24.2%) in patients with the worst periodontal parameters (PPD >7mm and ABL >4mm, Table 2 and Figure 1). Similarly, the periodontally healthiest patients had a 10.8% (95% CI 9.5-12.0%) probability of reporting a restricted diet, compared to 19.2% (95% CI 15.9-22.5%) for patients with PPD >7mm and ABL >4mm (Table 2 and Figure 2). For appearance, 22.2% (95% CI 20.6-23.8%) of the periodontally healthiest patients compared to 34.3% (95% CI 30.3-38.4%) of patients with the worst periodontal parameters reported an impact (Table 2 and Figure 3).

In addition, a similar trend was found in the adjusted probability of any impact, with 35.6% (95% CI 33.7-37.4%) and 9.2% (95% CI 8.1-10.4%) of the periodontally healthiest patients compared to 52.4% (95% CI 48.2-56.7%) and 17.5% (95% CI 14.3-20.8%) of patients with the worst periodontal parameters reporting at least one or more than one impact, respectively (Table 2 and Figures 4, 5).

Results from sensitivity analysis

A similar trend to the whole-group analyses, in regard to all PROs, was observed in analyses restricted to the 11,744 (81%) patients not requiring any restorations at the time of assessment. Similarly, omission of oral hygiene, dental attendance and salivary flow from the model did not yield appreciably different results.

Discussion

In the present cross-sectional study of a large, general dental practice-based population, we found a dose-dependent association between worsening periodontal status and the probability of reporting pain or restrictions in diet or unhappiness with dental appearance.

The large sample studied here allows for some meaningful and interesting comparisons between categories of different periodontal parameters. For example, a large proportion of patients (44.2%) studied here had evidence of radiographic bone loss of 2+mm, but no periodontal pockets deeper than 4mm, with approximately 20% exhibiting no BoP. While several causes of alveolar bone loss other than periodontitis must be considered, the absence of deep periodontal pockets in the presence of alveolar bone loss is consistent with the periodontal parameters one would expect to see following successful periodontal therapy, or resolution of active disease. Hence, our results are at least consistent with a beneficial effect of successful periodontal therapy and maintenance on the patient reported outcomes evaluated here, in particular given the specifics of this population, i.e., patients enrolled in a quality-assured prevention-oriented capitation-based payment plan scheme.

These results are in line with previous research on the association between periodontal disease and its treatment on oral health related quality of life. A recent systematic review summarising data from 14,087 patients in 37 studies, with sample sizes ranging from 21 to 3122 patients, found that patients with periodontal disease had poorer oral health related quality of life compared to periodontally healthy patients, and that the impact on quality of life was greater with increasing extent and severity of periodontal disease (Buset et al., 2016). All but one (Andersson et al., 2010) of the included 37 studies were based in hospital or specialist practices. The effect of treatment of periodontal disease on quality of life was reviewed by Shanbhag et al (2012), who reviewed 11 studies with sample sizes ranging from 32 to 183 patients, and concluded that periodontal treatment

1
2
3 can improve quality of life (Shanbhag et al., 2012). The authors reported an improvement in all
4 domains of oral health related quality of life following periodontal therapy. All of the 11 trials (7
5 prospective case-series or uncontrolled studies, one controlled study and three RCTs with a total of
6 639 participants) reviewed in this article were conducted in hospital or specialist practices.
7

8 There are several important strengths of this study. Firstly, we were able to analyse data from a
9 large, general practice based sample, resulting in precise estimates generalizable to a setting in
10 which a large number of patients receive care. Secondly, we were able to adjust for several
11 important potential confounders.
12

13 Several limitations of this study need to be considered. Firstly, the study used data from practices
14 that were part of a dental payment capitation scheme. As a result, this study likely included patients
15 with above average oral health motivation and compliance. For example, only 5% of patients were
16 classed as irregular attenders (Table 1). Therefore, results may not be generalizable to the entire UK
17 population. Secondly, as this is a pragmatic study using data collected as part of routine clinical care,
18 some measurement error and misclassification may arise due to lack of calibration, use of different
19 periodontal probes, use of routine radiographs. In this respect, assessment of oral hygiene, dental
20 attendance and salivary flow may be particularly subjective. These were included in the statistical
21 model as they are important confounders of the relationship between periodontitis and PROs
22 studied here. However, a sensitivity analysis conducted, omitting these three variables from the
23 model, resulted in findings that were not appreciably different. Thirdly, there is potential bias due to
24 unmeasured and residual confounding. This is likely to be most relevant for common risk factors for
25 caries and periodontal disease. For example, we had no data on some potential confounders, such as
26 socio-economic status. However, the fact that this is a somewhat more homogeneous population in
27 terms of oral health behaviours and socio-economic status reduces the risk of confounding relative
28 to the general population. Our sensitivity analysis, limited to patients who needed no restorations,
29 showed similar results.
30

31
32 In conclusion, our results demonstrate that in a large, non-specialist, general practice-based
33 population, worse periodontal health as measured by increased probing depth, alveolar bone loss
34 and bleeding on probing is associated with adverse patient reported outcomes including pain,
35 dietary restriction and unhappiness with appearance in a dose-dependent fashion. Hence,
36 prevention and successful management of periodontitis may have direct benefits on patient
37 reported outcomes.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References:

- 1
2
3
4
5
6 Andersson, P., Hakeberg, M., Karlberg, G. & Ostberg, A. L. (2010) Clinical correlates of oral impacts
7 on daily performances. *International Journal of Dental Hygiene* **8**, 219-226.
- 8 Busby, M., Chapple, E., Matthews, R. & Chapple, I. L. C. (2013) Practitioner evaluation of a novel
9 online integrated oral health and risk assessment tool: a practice pilot. *British Dental Journal*
10 **215**, 115-120.
- 11 Busby, M., Chapple, L., Matthews, R., Burke, F. J. T. & Chapple, I. (2014a) Continuing development of
12 an oral health score for clinical audit. *Br Dent J* **216**, E20-E20.
- 13 Busby, M., Martin, J. A., Matthews, R., Burke, F. J. T. & Chapple, I. (2014b) The relationship between
14 oral health risk and disease status and age, and the significance for general dental practice
15 funding by capitation. *British Dental Journal* **217**.
- 16 Buset, S. L., Walter, C., Friedmann, A., Weiger, R., Borgnakke, W. S. & Zitzmann, N. U. (2016) Are
17 periodontal diseases really silent? A systematic review of their effect on quality of life.
18 *Journal of Clinical Periodontology* **43**, 333-344.
- 19 Newton, J. T. & Asimakopoulou, K. (2017) The perceived acceptability of the DEPPA patient
20 assessment tool: A questionnaire survey of Denplan Excel patients. *British Dental Journal*
21 **222**, 767-770.
- 22 Shanbhag, S., Dahiya, M. & Croucher, R. (2012) The impact of periodontal therapy on oral health-
23 related quality of life in adults: a systematic review. *Journal of Clinical Periodontology* **39**,
24 725-735.
- 25 Sharma, P., Busby, M., Chapple, L., Matthews, R. & Chapple, I. (2016) The relationship between
26 general health and lifestyle factors and oral health outcomes. *British Dental Journal* **221**, 66-
27 70.
- 28 U. S. Department of Health and Human Services, F. a. D. A. (2009) Guidance for industry: patient-
29 reported outcome measures: use in medical product development to support labeling
30 claims. *Health and Quality of Life Outcomes* **4**, 79-79.
- 31 U.S. Department of Health and Human Services, F. a. D. A., Center for Drug Evaluation and Research,
32 Center for Biologics Evaluation and Research, Center for Devices and Radiological Health.
33 Guidance for Industry Patient-reported outcome measures: Use in medical product
34 development to support labeling claims., (ed.) M. Silver Spring. 2009 Food and Drug
35 Administration.
- 36 World Health Organization, W.H.O. (2012) Oral Health (Fact sheet N°318).
- 37
38 World Health Organization, W.H.O. Constitution of WHO: Principles.
39 <http://www.who.int/about/mission/en>. Accessed March 26, 2018
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Demographics of the cohort. Data are unadjusted and are expressed mean (SD) unless otherwise stated

	Whole cohort (n=14,568)	Cohort categorised by periodontal parameters							
		I (PPD <5 mm, ABL < 2mm, BoP -) (n=2,693; 18.5%)	II (PPD <5mm, ABL < 2mm, BoP +) (n=3,081; 21.2%)	III (PPD < 5mm, ABL 2-4mm, BoP -) (n=2,330; 18.0%)	IV (PPD < 5mm, ABL 2-4mm, BoP +) (n=2,898; 19.9%)	V (PPD 5-7mm, ABL 2-4mm, BoP +/-) (n=1,225; 8.6%)	VI (PPD < 5mm, ABL >4mm, BoP +/-) (n=911; 6.3%)	VII (PPD 5-7mm, ABL >4mm, BoP +/-) (n=868; 6.0%)	VIII (PPD >7mm; ABL >4mm, BoP +/-) (n=532; 3.7%)
Age (years)	55 (16)	48 (16)	46 (16)	62 (12)	59 (13)	59 (13)	66 (11)	64 (11)	63 (12)
Male (%)	43	40	43	43	43	48	44	46	49
Diabetic (%)	5.7	3.8	3.4	5.5	6.3	6.6	9.1	9.6	11.5
Never smoker (%)	63	69	70	62	62	58	51	48	48
Teeth present*	25 (4)	26 (4)	26 (3)	25 (4)	25 (4)	25 (4)	23 (5)	23 (5)	23 (5)
Restored teeth	11 (6)	9 (6)	9 (6)	13 (5)	13 (5)	12 (5)	12 (5)	12 (5)	11 (5)
Teeth needing restoration	0.3 (1.0)	0.2 (0.9)	0.3 (1.1)	0.3 (0.7)	0.4 (1.0)	0.3 (1.0)	0.4 (1.1)	0.4 (1.1)	0.4 (1.1)
Frequency of dental visits less than recommended (%)	5	3	6	4	4	7	3	7	8
Improvement in oral hygiene needed (%)	44	18	54	27	58	52	58	56	64
High frequency of sugar intake (%)	11	12	14	10	10	11	11	10	14
Patient reported outcomes (%)									
Pain	15	14	15	14	15	17	14	18	21
Diet restrictions	11	11	9	10	11	13	13	16	20
Unhappiness with appearance	26	23	26	26	26	27	30	31	33
Any patient reported concerns	40	36	38	39	39	42	43	46	52
More than one patient reported concerns	11	9	10	10	11	12	11	15	18

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

*not including wisdom teeth

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2: Probabilities (and 95% CI) of reporting various outcomes by periodontal parameters. Adjusted for age, sex, smoking status, diabetic status, number of teeth with restorations, number of teeth needing restorations, oral hygiene, dental attendance, abrasion, grinding habits, frequency of sugar intake and salivary flow rate.

Adjusted probability (%) of reporting	Periodontal Parameters							
	I (PPD < 5mm, ABL < 2mm, BoP -)	II (PPD < 5mm, ABL < 2mm, BoP +)	III (PPD < 5mm, ABL 2-4mm, BoP -)	IV (PPD < 5mm, ABL 2-4mm, BoP +)	V (PPD 5-7mm, ABL 2-4mm, BoP +/-)	VI (PPD < 5mm, ABL >4mm, BoP +/-)	VII (PPD 5-7mm, ABL >4mm, BoP +/-)	VIII (PPD >7mm; ABL >4mm, BoP +/-)
Oral pain	13.8 (12.4-15.2)	13.9 (12.7-15.2)	14.9 (13.4-16.4)	15.0 (13.7-16.3)	17.3 (15.2- 19.3)	13.7 (11.5-16.0)	17.7 (15.1- 20.3)	20.7 (17.2- 24.2)
Restricted diet	10.8 (9.5- 12.0)	9.7 (8.5- 10.8)	9.5 (8.3-10.7)	11.2 (10.1-12.4)	12.5 (10.7- 14.3)	12.1 (10.0-14.2)	15.5 (13.1- 17.9)	19.2 (15.9- 22.5)
Unhappiness with appearance	22.2 (20.6- 23.8)	24.5 (23.0- 26.1)	26.6 (24.8- 28.4)	26.0 (24.4- 27.6)	27.3 (24.8-29.8)	31.7 (28.6-34.8)	32.5 (29.4- 35.6)	34.3 (30.3- 38.4)
Reporting any concerns	35.6 (33.7-37.4)	37.2 (35.5-39.0)	39.1 (37.1-41.4)	39.1 (37.3-40.9)	42.3 (39.6-45.0)	43.8 (40.5-47.1)	46.7 (43.4-50.0)	52.4 (48.2-56.7)
Reporting more than one concern	9.2 (8.1-10.4)	9.6 (8.5-10.7)	9.9 (8.7-11.2)	10.8 (9.6-11.9)	12.2 (10.4-14.0)	11.4 (9.3-13.5)	15.3 (12.8-17.7)	17.5 (14.3-20.8)

ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1) Probability of reporting oral pain Vs Periodontal parameters
 ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

101x73mm (300 x 300 DPI)




Figure 2) Probability of reporting restricted diet Vs Periodontal parameters
 ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

101x73mm (300 x 300 DPI)




Figure 3) Probability of reporting unhappiness with appearance Vs Periodontal parameters
 ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

101x73mm (300 x 300 DPI)




Figure 4) Probability of reporting oral pain or restricted diet or unhappiness with appearance Vs Periodontal parameters

ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

101x73mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5) Probability of reporting more than one of oral pain or restricted diet or unhappiness with appearance Vs Periodontal parameters
 ABL: Alveolar bone loss ; Bleeding on Probing: BoP, either present (+) or absent (-) ; PPD: periodontal probing depth

101x73mm (300 x 300 DPI)