

## Global burden of postoperative death

National Institute for Health Research Global Health Research Unit on Global Surgery

DOI:

[10.1016/S0140-6736\(18\)33139-8](https://doi.org/10.1016/S0140-6736(18)33139-8)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

National Institute for Health Research Global Health Research Unit on Global Surgery 2019, 'Global burden of postoperative death', *The Lancet*, vol. 393, no. 10170, pp. 401-401. [https://doi.org/10.1016/S0140-6736\(18\)33139-8](https://doi.org/10.1016/S0140-6736(18)33139-8)

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for eligibility: 09/05/2019

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## **Deaths within 30 days of surgery in low-, middle-, and high-income countries**

*Dmitri Nepogodiev MBChB, Janet Martin PharmD, Prof Bruce Biccard PhD, Alex Makupe MMed Surgery, Aneel Bhangu PhD on behalf of the NIHR Global Health Research Unit on Global Surgery\**

\*Collaborating co-authors listed in appendix

**Corresponding author:** Mr Aneel Bhangu, MBChB PhD FRCS, NIHR Global Health Research Unit on Global Surgery, 2nd Floor, Institute of Translational Medicine, Heritage Building, University of Birmingham, Mindelsohn Way, Birmingham, B15 2TH. Correspondence to: [a.a.bhangu@bham.ac.uk](mailto:a.a.bhangu@bham.ac.uk)

**Category:** Correspondence

**Word count:** 395

**Acknowledgements:** None.

**Declaration of interests:** The authors have no competing interests to declare.

**Funding:** This report was funded by a National Institute for Health Research (NIHR) Global Health Research Unit Grant (NIHR 17-0799). The funder had no role in study design, data collection, analysis and interpretation, or writing of this report. The funder has approved the submission of this report for publication. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the UK Department of Health and Social Care.

The Lancet Commission on Global Surgery identified that 313 million surgical procedures are performed each year<sup>1</sup>. Little is known about the quality of surgery globally, as robust postoperative death rates are available for only 29 countries<sup>2</sup>. The rate of postoperative death captures the success of the system of surgical care and improvement in this metric is a priority worldwide.


We aimed to estimate how many people around the world die within 30 days of surgery, based on surgical volume, case-mix, and postoperative death rates adjusted for country income. England's Hospital Episode Statistics linked Office of National Statistics dataset (HES-ONS) is one of the world's most comprehensive procedure-specific mortality resources, with national coverage from a universal health care system. We used this dataset as the baseline for our estimations for high income settings, adjusting case-mix and mortality to estimate total postoperative deaths in low- and middle-income countries (LMICs). We estimated likely additional postoperative deaths if surgical volume was expanded to address the annual unmet need for 143 million operations in LMICs<sup>3</sup> (S1 Appendix).

Overall, our analysis suggests that currently at least 4.2 million people around the world die within 30 days of surgery each year, with half of these deaths occurring in LMICs. We project that expanding surgical services to address unmet need would increase total deaths to 6.1 million annually, with an additional 1.9 million occurring in LMICs each year. Based on 4.2 million deaths, 7.7% of all deaths globally occur within 30 days of surgery<sup>4</sup>. This figure is greater than that attributed to any other cause of death globally except ischaemic heart disease and stroke (Figure 1). More people die within 30 days of surgery than the combined 2.97 million deaths attributed to HIV, malaria, and tuberculosis annually<sup>4</sup>.

Our approach is limited by a number of necessary assumptions (S2 Appendix). For example, HES-ONS reports amongst the lowest postoperative death rates in the world. Basing our calculation on higher baseline postoperative death rates from other HIC sources substantially increases our projections of total postoperative deaths.

Whilst there is a pressing need to expand surgical services to currently underserved populations, this must be done in tandem with initiatives to reduce postoperative deaths. Funders and policy makers should prioritise research that aims to make surgery safer, particularly in LMICs. Routine measurement of surgical outcomes is essential to monitoring global progress in addressing the burden of postoperative death.

**Figure 1: Top ten global causes of death in the 2016 Global Burden of Death study, with addition of postoperative death\***



\*Percentages give proportion of total global deaths attributable to that cause based on Global Burden of Disease 2016 data<sup>4</sup>

IHD: ischaemic heart disease; COPD: chronic obstructive pulmonary disease; LRTI: lower respiratory tract infections

## Authors' contributions

DN and AB conducted data analysis and interpretation and had access to all data. DN, JM, BB, AM, AB drafted the manuscript. Collaborators listed in the Appendix critically revised the manuscript and have approved the final draft and the decision to submit the manuscript. AB is the guarantor for this report.

## References

1. Meara JG, Leather AJ, Hagander L, et al. Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *Lancet* 2015; **386**(9993): 569-624.
2. Marks IH, Kamali P, Khan MA, et al. Data for the sustainable development of surgical systems: A global collaboration, 2016.
3. Rose J, Weiser TG, Hider P, Wilson L, Gruen RL, Bickler SW. Estimated need for surgery worldwide based on prevalence of diseases: a modelling strategy for the WHO Global Health Estimate. *Lancet Glob Health* 2015; **3 Suppl 2**: S13-20.
4. Collaborators GBDCoD. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**(10100): 1151-210.