

Magnetic resonance imaging quantification of gastrointestinal liquid volumes and distribution in the gastrointestinal tract of children

Papadatou-Soulou, Eleni; Mason, Julie; Parsons, Caron; Oates, Adam; Thyagarajan, Manigandan; Batchelor, Hannah Katharine

DOI:

[10.1021/acs.molpharmaceut.9b00510](https://doi.org/10.1021/acs.molpharmaceut.9b00510)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Papadatou-Soulou, E, Mason, J, Parsons, C, Oates, A, Thyagarajan, M & Batchelor, HK 2019, 'Magnetic resonance imaging quantification of gastrointestinal liquid volumes and distribution in the gastrointestinal tract of children', *Molecular Pharmaceutics*, vol. 16, no. 9, pp. 3896-3903.
<https://doi.org/10.1021/acs.molpharmaceut.9b00510>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Molecular Pharmaceutics*, copyright © American Chemical Society after peer review and technical editing by the publisher.

To access the final edited and published work see <https://doi.org/10.1021/acs.molpharmaceut.9b00510>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Magnetic resonance imaging quantification of gastrointestinal liquid volumes and distribution in the gastro-intestinal tract of children

Authors

Eleni Papadatou-Soulou¹, Julie Mason¹, Caron Parsons², Adam Oates³, Manigandan Thyagarajan³,

5 *Hannah Katharine Batchelor^{1*}*

¹ *School of Pharmacy, Institute of Clinical Sciences, University of Birmingham, Edgbaston, B15 2TT, UK*

² *University Hospitals Coventry and Warwickshire NHS Trust, Clifford Bridge Rd, Coventry CV2 2DX, UK*



³ *Birmingham Children's Hospital NHS Trust, Steelhouse Lane, Birmingham B4 6NH, UK*

Corresponding author: Hannah Batchelor h.k.batchelor@bham.ac.uk

10 Dr Hannah Batchelor
Senior Lecturer in Pharmaceutics, Formulation and Drug Delivery
School of Pharmacy
Institute of Clinical Sciences
Robert Aitken Building
15 University of Birmingham
Edgbaston, B15 2TT
Tel: +44 (0)121 414 3717

Table of Contents Graphic

20



25 **Abstract**

The volume and localisation of fluid in the paediatric gastro-intestinal tract is crucial to inform the design of *in vitro* and *in silico* models that predict the absorption of oral drugs administered to children. Previous studies have used magnetic resonance imaging (MRI) to quantify fluid volumes and localisation in the intestines of adults; this study is the first to undertake similar analysis of paediatric participants.

30 This study quantified the amount and distribution of fluid in fasted and fluid-fed children using MRI data captured during routine clinical assessment.

Data from 32 fasted children (aged 0-16 years) and 23 fluid-fed (aged 8-16 years) was evaluated. The gastric volume ranged from 0-9mL in the fasted and 19-423mL in the fluid-fed state. The small intestinal volume was recorded to be 0-51mL in the fasted and 6-91mL in the fluid-fed state with an average
35 number of 7.7 and 22.4 fluid pockets respectively. The data showed significant differences in gastric volumes and the number of fluid pockets in the small intestine for age-matched fasted and fluid-fed children ($p < 0.05$). Both the number and the volume of pockets reported in children are much lower than those previously reported in adults.

This study is the first to report intestinal volumes and localisation in children and provides new
40 information to inform the design of biorelevant *in vitro* models and real values to update *in silico* models. The availability of data from both fluid-fed and fasted children show the extremes of fluid volumes that are present in the gastro-intestinal tract which is useful to understand the variability associated with drug absorption in children.

Keywords: gastric fluid; intestinal fluid; MRI; children; biorelevant dissolution;

45

Introduction

Oral drugs are often administered as solid dosage forms (e.g. suspensions; tablets or capsules) where disintegration and dissolution are essential prior to absorption. Variable absorption profiles are often observed that are attributed to individual inconsistencies within the gastro-intestinal environment ¹.

50 Significant efforts have been made to develop *in vitro* and *in silico* models that replicate the GI environment to better understand the processes of disintegration and dissolution with a view to minimising the variability associated with absorption. Typically pharmacokinetic profiles in children are predicted based on extrapolation from adult data based on body mass or surface area algorithms rather than using physiological data. However, for oral absorption knowledge of the GI fluid is important to
55 predict performance in children.

Prior to absorption of orally administered drugs, medicines often have to undergo disintegration and dissolution within the gastro-intestinal tract. The impact of the volume and composition of gastro-intestinal (GI) fluids will influence the processes of disintegration and dissolution and can affect the subsequent absorption profile. The volume of fluid used within *in vitro* testing systems varies from low
60 values (<50mL) in biorelevant testing systems to 900mL for pharmacopoeial methods as limited data exist on relevant values to use. The biopharmaceutics classification system uses fluid volume as a key parameter to classify drugs where the dose number (D_0) describes the volume in which the maximum dose strength is soluble; this is used to classify medicines as either highly or poorly soluble. The D_0 used in the calculations for determining BCS criteria is determined by the formula:

$$D_0 = \frac{M}{(V)C}$$

65 where M is the dose strength of the oral dosage form (mg), V is the volume administered (mL), and C is the drug's solubility (mg/mL).

A dose number value greater than 1 indicates a poorly soluble drug whereas less than 1 is highly soluble. A highly soluble drug is likely to be well absorbed as solubility within the GI environment is not a rate limiting step for absorption. In the USA and Europe a volume of 250mL (representing a glass of water) is
70 used to calculate the dose number, whereas in Japan the volume is lower at 150mL to account for reduced volumes of water co-administered in clinical testing protocols ². Extrapolation of adult dose numbers to paediatric populations has been undertaken to obtain age-appropriate fluid volumes

administered using body surface area ^{3,4} giving values of 33.6mL for a neonate; 66mL for an infant and 127.6mL for a 7 year old child. Other literature reported a “worst-case” scenario of 25 mL as an administration volume for all paediatric subgroups ⁵. Normalisation of the residual gastric fluid volume using extrapolation from adults on a mL per Kg basis gave gastric fluid volume values of 14.9mL in neonates; 29.8mL in a 6 month old infant and 86.8mL in a 7 year old child ⁶. A recent study on minipigs showed that gastric fluid volume was linearly related to the weight of the pig which suggests that weight is a suitable measure to use to extrapolate gastric fluid volumes ⁷.

Knowledge of the volume and composition of fluids in the gastro-intestinal environment are critical to prediction of absorption as they provide an understanding of the media for disintegration and dissolution which can be replicated in *in vitro* and *in silico* models. Significant efforts have been made to measure both volume ⁸⁻¹⁰ and composition of gastro-intestinal fluids in adults ¹¹⁻¹⁴ and to a more limited extent children ¹⁵. Fluid within the adult small intestine has been reported to be present in pockets rather than homogeneously distributed along the GI tract. This distribution pattern will further complicate the absorption profile and adds to variability observed in pharmacokinetic profiles from orally absorbed medicines ¹⁶. The comparative gastric and small intestinal volume values in children are likely to have a substantial impact on the rate and extent of drug absorption following oral administration which provides a more physiological approach to extrapolation of data.

The impact of gastro-intestinal fluid volume on predictions of pharmacokinetic profiles has been shown to be significant for poorly soluble drugs in adult populations ¹⁷. Therefore, it is critical that representative fluid volumes are used to predict absorption for all populations. The range of fluid volumes from the fasted to fed state is also of interest as the variation can impact upon the absorption of medicines. Therefore, knowledge of the fluid volumes in the extremes of fasted and fed states is of interest.

Previous studies conducted in adults demonstrated the value of magnetic resonance imaging (MRI) to accurately measure the volume and localisation of fluid within the gastro-intestinal tract ^{8-10, 16}. Many children undergo MRI as part of their clinical care where their abdomen is visible. This study sought ethical approval to use data from clinical MRI procedures in children to measure fluid volumes and localisation in the stomach and small intestine of these paediatric patients. The primary outcome measure was to quantify the mean and range of fluid volumes within the stomach and small intestine of children. Secondary outcome measures were to quantify the mean (and range) number of fluid pockets

distributed within the GI tract of children and to record their location (duodenum, jejunum and ileum). The study also explored the difference in these outcomes from fluid-fed and fasted participants.

105

Materials and Methods

This study was an observational, retrospective study with non- probability sampling. The project sites were two major teaching Hospitals in the West Midlands (University Hospitals Coventry and Warwickshire NHS Trust and Birmingham Children’s Hospital). The project was ethically approved REC reference: 18/EM/0251 (IRAS 237159 MRI: Fluid volumes and localisation in paediatric GI tract).

110

Study Participants

Anonymised MRI data sets were provided to the researchers from children (from 0-16 years) who required MRI for clinical purposes where the abdomen (stomach and small intestine) were clearly visible. The two sites involved had different protocols for the MRI procedure: one required children to ingest 500mL Oral Klean Prep (a macrogol solution) in the 60 minutes prior to their MRI (hereon referred to as fluid-fed children); the other site required children to fast overnight prior to the MRI.

115

In order to ensure that the patient population was as close to “normal-healthy” as possible, the following exclusion criteria were used, patients with: acute abdomen (appendicitis or perforated viscous); malignant bowel disease; surgery (bowel section, excluding appendectomy); bowels wall thickening/ stricture/ fistula/ abscess.

120

Magnetic resonance Imaging (MRI)

MRI scanning was performed at two sites using different apparatus, the methods of subsequent analysis accounted for these differences in equipment used to acquire images. This imaging process provided clear images where fluids can be clearly distinguished.

125

The fluid fed participants were scanned at University Hospital Coventry and Warwickshire using either a 1.5T MR imaging unit (Optima MR450w, GE Healthcare, Chicago, USA) with a 48-channel body coil where the MRI protocol used was a coronal balanced steady-state gradient echo sequence (FIESTA) (slice thickness= 4.0 mm; echo train length = 1; intersection gap = 5.0mm; matrix size, 512 x 512; field of view, a x b cm; TR/TE, 5.7/1.9 ms). Or the scan was conducted using a 1.5T MR imaging unit (Aera,

130

Siemens Healthcare, Erlingen, Germany) using a body coil and the MRI protocol used was a coronal balanced steady-state gradient echo sequence (True FISP) (slice thickness = 6.0 mm; echo train length = 1; intersection gap = 3.0mm; matrix size, 256 x 256; field of view, a x b cm; TR/TE, 652.8/2.1 ms).

135 The fasted participants were scanned at Birmingham Children's Hospital using either a 1.5T MR imaging unit (Siemens MAGNETOM Avanto 1.5T MRI System, USA) with a 16-element parallel imaging receiver coil where the MRI protocol used was a T2 SPACE coronal sequence (no gap between slices; slice thickness= 0.90 mm; matrix size = 0.8 x 0.8mm; field of view = 250; TR/TE = 1700/98 ms). Alternatively a 1.5T MR imaging unit (Aera, Siemens Healthcare, Erlingen, Germany) using a 16-element parallel imaging receiver coil was used where the MRI protocol used was a coronal T2 SPACE sequence (no gap
140 between slices; slice thickness, 0.9 mm; matrix size = 0.8 x 0.8 mm; field of view = 400; TR/TE = 2000/241 ms).

T2 SPACE is a variant of a three dimensional (3D) turbo spin echo sequence. Compared to a conventional turbo spin echo sequence, T2 SPACE uses non-selective, short refocusing pulse trains that consist of radiofrequency pulses with variable flip angles. This allows for very high turbo factors (> 100)
145 and high sampling efficiency. A SPACE sequence produces high resolution isotropic images which can be reconstructed in multiple planes. SPACE sequences are less sensitive to susceptibility, flow and chemical shift artifacts which make it superior over the conventional turbo spin echo. By using a SPACE sequence, it was possible to produce isotropic 0.9mm 3D T2 contrast images in less than 7 minutes.


150 **Image analysis**

Horos software was used to visualise the MRI data sets. Horos is a free and open source code software (FOSS) program that is distributed free of charge under the LGPL license at Horosproject.org and sponsored by Nimble Co LLC d/b/a Purview in Annapolis, MD USA. The fluid within images was identified using the cerebrospinal fluid as a reference point for each data set; this approach acts as an internal
155 control for each data set and represents the "free" water present (^{18,19}). This value was used to provide a threshold to enable the fluid in the MRI data set to be clearly visualised. In brief, the image slice that most clearly showed the CSF within the spinal canal was selected to generate the threshold value. A rectangle was drawn entirely within the CRF and HOROS software was used to calculate the signal intensity of this area to set the limit for free fluid within the MRI data set. In this study, signals from the
160 stomach and the small intestine were manually identified whereas the kidneys, gallbladder, bladder and

visible vessels were manually excluded from subsequent analysis. Representative images are shown in Figure 1. The gastro-intestinal fluid was identified in each slice manually and the area of each fluid pocket was calculated following transfer of the image into imageJ²⁰. The volume of each pocket was determined by multiplication of the slice thickness by the area measured as previously described¹⁶.

165 Using each patient's individual data, it was possible to create 3D MRI images (using Horos software) that assisted in mapping the GI anatomy which was particularly useful in the determination of the location of the identified fluid pockets. The 3D images were used to identify fluid pockets that overlapped into adjacent slices to better calculate the volume of each continuous pocket. For each participant, the total number of pockets was recorded as well as the volume and location of each pocket. Location of pockets
170 within the small intestine was based on the visual distinction of small and large bowel, based on anatomical knowledge and observed differences after thresholding the slices using HOROS software. It was challenging to identify where the duodenum, jejunum and ileum start and finish, as there is no specific small bowel length for all subjects. Approximation of the location of fluid pockets was undertaken by dividing the total small bowel into two parts; the upper left hand corner in an X,Y coronal
175 plane was used as the starting point and moving in a diagonal line the upper left 2/5 were designated to be the duodenum and jejunum; the lower right hand 3/5 of the image represented the ileum; this was used to then determine where the fluid pockets were located.

All analysis was undertaken by a single operator initially with a sub-sample of 10% of all participants being checked by a second operator to ensure that there was consistency ($\pm 10\%$) in the measurements
180 obtained.



185 **Figure 1.** Example MRI data from two participants; the images show a sequence of slices in the coronal plane moving from the front to the back (spine) of the subject; the red areas are the fluid areas thresholded and included in further analysis. The upper image set is from a fluid-fed 16 year old and the lower imaged from a fasted 8 year old.

190 **Statistical analysis**

The individual volumes and number of fluid pockets were plotted to provide data on individual variability. The data are also shown as the median, interquartile values and full range.

Statistical tests were performed using SPSS. An independent samples t-test was used to compare the volume of fluid and number of fluid pockets between age-matched fluid-fed and fasted children. Where multiple age-groups were compared ANOVA tests were undertaken.

195 **Results**

MRI data sets were available from 32 fasted children and 23 fluid-fed children. The demographics of the participants are shown in Table 1. The youngest child was one week old and the oldest 16 years old. The gender split was approximately even with 52% (12/23) of the fluid-fed participants and 59% (19/32) of the fasted participants being female. All data sets were of sufficient quality to allow subsequent analysis.

Number of data sets available

Age range	Fasted Children	Fluid-fed Children
<2 years (newborn/infant/toddler)	10	
2-5 years (pre-school children)	13	
6-11 years (school age children)	8	6
12-16 years (adolescents)	1	17

200 **Table 1.** Demographics of the participants included in the study.

Gastric fluid volume

205 The gastric fluid was either present as a single pocket or absent. Residual gastric fluid was absent in 65.6% (21/32) of fasted children and present in all fluid-fed children. A comparison of the total volume of gastric fluid present in fasted vs fluid-fed children is shown in Figure 2. The difference in gastric fluid present in fluid-fed versus fasted children was statistically significant ($p < 0.05$).





Figure 2. Comparison of gastric volumes in fluid-fed (n=23) and fasted (n=32) children. Data shows median; Q1; Q3 and range.

210 The age of the child had no impact on the volume of gastric fluid present in either fasted or fluid-fed children ($p > 0.05$); this data is shown in the supporting information (Figure S1).

Number of pockets in the small intestine

The total number of fluid pockets identified in the small intestine of fasted patients ranged from 0-16; whereas for fluid-fed patients it ranged from 11-40, as shown in Figure 3.



220 **Figure 3.** Comparison of the number of fluid pockets identified in fluid-fed (n=23) (shown as black circles) and fasted (n=32) (shown as white circles) children. Data is stratified by age group as well as the fluid-fed and fasted state.

As the fluid-fed children were typically older than the fasted children the impact of age on the number of fluid pockets was explored. Participants were stratified into groups based on those <2 years; 2-5
225 years; 6-11 years and 12-16 years to determine whether there was a difference in the number of fluid pockets identified. The number of pockets in aged-matched (school age children) was significantly different ($p = 0.011$) for fluid-fed (n=6) and fasted (n=8) children. Pooled analysis of all participants showed that there were significant differences in the number of pockets identified based on age for the


overall population ($p < 0.05$); see supplementary material. However, there was no statistically significant
230 difference ($p = 0.420$) for the number of pockets in fasted children based on their age.

Location of fluid pockets

In both the fluid-fed and fasted populations the majority of fluid pockets were located in the jejunum;
86% in fluid-fed children and 93% in fasted children, all remaining pockets were located in the ileum.

235 Small intestinal fluid volume

The overall volume of fluid in the small intestine was compared by both age range and the fluid-
fed/fasted state. The results are shown in Figure 4. The mean total small intestinal volumes were 30 ± 24 mL and 7 ± 10 mL in the fluid-fed and fasted state respectively.




240 **Figure 4.** Total small intestinal volume presented as a function of age range and the fluid-fed/fasted state.

245 Statistical analysis showed that there is not a statistically significant difference in the total volume of
 fluid in the small intestine in school age children between the fluid-fed and fasted state ($p=0.101$);
 although there is an outlier with one fasted participant with a small intestinal fluid volume of 50mL
 which has skewed the results; elimination of this outlier resulted in a significant difference in the values
 ($p<0.05$). Age had no significant effect on the total volume for those in the fluid-fed or the fasted state
 250 sub-populations.

The proportion of pockets that were less than 1mL was 66.0% (306/464) in fluid-fed children and 76.5%
 (202/264) in fasted children. These small pockets contributed to 21% of the total fluid found in the small
 intestine of the fluid-fed participants and 27% for the fasted participants. Figure 5 shows (a) the relative
 255 number and (b) proportion of fluid present based on pocket size in the intestine of the fluid-fed and
 fasted participants; this are shown within age sub-populations although no differences were seen based
 on age.

(a)



260 (b)




Figure 5. Comparison of (a) the percentage of fluid pockets by each volume per sub-population and (b) the total volume occupied by these pockets by each sub-population.

Discussion

265 Knowledge of the volume and distribution of gastric and small intestinal fluid in children in both the fluid-fed and fasted state is of great interest to the pharmaceutical industry. This information will inform the design of biorelevant *in vitro* and *in silico* methods that will predict the performance of oral solid dosage forms which can, in turn, aid in prediction of the pharmacokinetics of a drug. In the current study two populations were available; those who were fasted and those who were fluid-fed having ingested up to 500mL of KleanPrep in the 60 minutes prior to data collection. These two populations enable the impact of fluid on overall intestinal volumes to be explored although the data does not exactly replicate clinical testing as has been undertaken in adult studies due to ethical constraints surrounding paediatric clinical testing¹⁰.

275 The MRI protocol for fluid-fed participants required children to consume 500mL Oral Klean Prep (a macrogol solution) in the 60 minutes prior to their scan yet this was not monitored thus there is likely to be variability in the total amount consumed and the time frame for consumption prior to data capture. Data is available that details the time that the fluid-fed participants started the study and the time the MRI was conducted which ranged from 14 minutes to 2 hours 35 minutes with a mean value of 68 minutes. There was no correlation with time compared to volume in the small intestine (correlation showed $R^2=0.0006$) (Shown in Figure S2 of supporting information).

280

Our results showed that fasted children had gastric volumes ranging from 0 to 8mL with a mean volume of 1.3 mL. The fasted gastric volume in children is very much lower than that reported in adults (see Figure 6) which may have consequences on the disintegration and dissolution of medicines given to children in a fasted state. The inter-individual variability in adult fasted gastric fluid volume has previously been reported as having a relative standard deviation of 31% with intra-individual variability relative standard deviation of 23%²¹. A comparison of the gastric volume reported in this study compared to previous values from adults is shown in Figure 6.




Figure 6. Comparison of the mean (\pm standard deviation) reported for fasted gastric volumes in adult studies compared to the current paediatric study. [Steingoetter study only reported a range not a mean and standard deviation]. Note that several studies included sub-populations hence multiple data from the same source.


The fasted gastric volume reported here is very much lower than that previously shown in adults. Fasting gastric volumes previously reported in children were 0.4-0.56 ml/Kg^{22,23}. The data in this study would provide an estimated value would be 0.08mL/Kg (based on a mean age of 4 years with an approximate weight of 16Kg). This much lower fasting gastric volume has implications in dosing to children where it is important to highlight the need to dose with water. Oral suspensions are a

commonly used dosage form in children where drug dissolution is a rate limiting step for absorption. Previous work has demonstrated that larger gastric fluid volumes enhance dissolution of poorly soluble drugs²⁴, therefore low gastric volumes may impact the onset of drug action. Medicines administration in children is often not accompanied by an excess of water, particularly for liquid medications²⁵. Furthermore, there is evidence that children are often on fluid restricted diets prior to night-time dosing which may lead to slow onset of action for medicines given at night²⁶. The lowest fasted gastric volume previously reported in adults was 1mL and the highest was 95mL⁸; our results showed that almost two-thirds of fasted children had no measureable gastric fluid volume.

Previous work that has looked at the impact of gastric volumes following ingestion of water have shown increased gastric volumes of 296mL immediately after ingestion of 300mL water²⁷ and 242mL following ingestion of 240mL water¹⁰. Our study collected data from children who had ingested up to 500mL Kleanprep in the 60 minutes prior to the MRI; the data showed increased gastric volume yet the data is not directly comparable to that found in adults due to the differences in protocols used. Ethical restrictions limit the extent of experimentation that can be conducted using MRI with paediatric populations.

This study has confirmed that, as previously seen in adults, fluid is present in discontinuous pockets in the small intestine of children in both the fluid-fed and fasted state. Discontinuous pockets of fluid in the small intestine of adults was first reported in 2005 by Schiller et al. The total number of pockets identified in our study was 8 ± 4 and 22 ± 4 in the fasted and fluid-fed state respectively. This data is similar to that reported by Mudie et al where 8 pockets were reported in the fasted state in adults and 16 in the fed state (in adults)¹⁰.

A total mean small intestinal volume of 105 mL (range 45-319mL) was reported in the fasted state in a study conducted by Schiller et al in 2005. A lower total volume of 43mL (range 5-158mL) was reported by Mudie et al (2014)¹⁰ which was similar to a value of 54 ± 37 mL reported by Grimm et al (2018)⁹. Our study reported a fasted mean small intestinal volume of 7.4mL (range 0-50.58 mL) in children which is much lower than values previously reported in adults. The forest plot in Figure 7 compares the volumes of fasted small intestinal fluid reported in the literature to the values found in our study.



325

Figure 7. Comparison of reported volumes of fasted small intestinal fluid in adult studies compared to the current paediatric study (data shows mean and range except for Grimm (2018b) study where mean \pm standard deviation is shown as range data was not available).

330

This lower volume in children is even smaller than values previously used in representative paediatric dissolution testing. Conventional dissolution testing of pharmaceutical dosage forms typically involves large volumes of fluid (500-900mL). The appropriate volume to be used to represent the paediatric intestinal media has previously been suggested to be 50mL for neonates; 100mL for infants and 200mL for pre-school children²⁸. This study suggests that volumes of 7-40mL may be more appropriate in pre-school and school aged children. However, it is important to consider that this volume is not present as a single homogenous pocket but is distributed into several small pockets along the intestinal tract.

335

Figure 8 shows a cartoon bubble plot that represents the distribution and relative size of fluid pockets in fluid-fed and fasted children.

340




Figure 8. Representation of fluid pockets in fluid-fed children (blue) and fasted children (red); each bubble represents one pocket and the bubble size the relative volume. The data is presented from left to right in order of age for each sub-population.

Limitations of our data set need to be considered. The presented data were all collected from children undergoing MRI for clinical reasons, therefore the protocol for ingestion of fluid was not tightly controlled and the actual volume and timing of ingestion of the recommended 500mL of Oral Klean Prep was not accurately recorded for the fluid-fed participants. Adherence to the protocol for fasted children was also not checked where children were asked to fast overnight prior to the MRI. It is recognised that using Klean Prep (an osmotic laxative) to represent fluid fed children is not fully representative of a typical fluid fed child. The mechanism of action of this product is to draw water into the intestines which maximises the fluid content increasing the water content and volume of stools in the bowel. The Klean prep was administered within 60 minutes of imaging thus the fluid present in the intestine was observed prior to defecation. This provided a fluid fed scenario to allow comparisons to the fluid fasted children. The age ranges of the children who were fasted and fluid-fed are not matched and additional efforts are ongoing to try to match the demographics of fluid-fed and fasted children.

360 **Conclusion**

This study is the first to quantify the total volume and distribution of fluid in the stomach and small intestine of children in the fasted and in a fluid-fed state. Fluid in the small intestine is present in discontinuous pockets with a mean volume of 7.4mL in the fasted state and 30.4mL in the fluid-fed state. There are a significantly higher number of pockets in the fluid-fed state with a significantly higher overall volume. The information presented has implications to inform the design of biorelevant *in vitro* and *in silico* methods to predict drug absorption in children.

365

The very low volumes observed in childrens' intestines compared to previous studies in adults raises concerns about the direct extrapolation of *in vitro* dissolution data based on adult physiology to predict performance of medicines in paediatric populations.

370

Author Information

Corresponding Author: *(H.K.B) Tel: +44 (0)121 4143717 E-mail: h.k.batchelor@bham.ac.uk

375 Author Contributions: J.M, C.P, A.O, M.T and E.P-S and H.K.B. designed the study. E.P-S. analyzed the
image data. The manuscript was written through contributions of all authors. All authors have given
approval to the final version of the manuscript.

The authors declare no competing financial interest.

Acknowledgments

380 The research described in this paper was sponsored by a Ph.D. grant from Janssen Research &
Development, A Division of Janssen Pharmaceutica NV.

Abbreviations

GI, gastrointestinal; MRI, magnetic resonance imaging; CSF, cerebrospinal fluid; PBPK models,
Physiologically based Pharmacokinetic models; UHCW, University Hospitals Coventry and Warwickshire;
BCH; Birmingham Children's Hospital.

385

Supporting Information is Available

- Comparison of gastric volume vs age for fluid-fed and fasted participants
- Relationship between the total small intestinal volume measured and the time lapsed from instruction to consume fluid to the time of MRI data collection for the fluid-fed participants

1. Kimura, T.; Higaki, K., Gastrointestinal transit and drug absorption. *Biological & pharmaceutical bulletin* **2002**, *25* (2), 149-64.
2. Kuribayashi, R.; Takishita, T.; Mikami, K., Regulatory Considerations of Bioequivalence Studies for Oral Solid Dosage Forms in Japan. *Journal of Pharmaceutical Sciences* **2016**, *105* (8), 2270-2277.
3. del Moral Sanchez, J. M.; Gonzalez-Alvarez, I.; Cerda-Revert, A.; Gonzalez-Alvarez, M.; Navarro-Ruiz, A.; Amidon, G. L.; Bermejo, M., Biopharmaceutical optimization in neglected diseases for paediatric patients by applying the provisional paediatric biopharmaceutical classification system. *British Journal of Clinical Pharmacology* **2018**, *84* (10), 2231-2241.
4. Gandhi, S. V.; Rodriguez, W.; Khan, M.; Polli, J. E., Considerations for a pediatric Biopharmaceutics Classification System (BCS): Application to five drugs. *AAPS PharmSciTech* **2014**, *15* (3), 601-611.
5. Abdel-Rahman, S. M.; Amidon, G. L.; Kaul, A.; Lukacova, V.; Vinks, A. A.; Knipp, G. T., Summary of the National Institute of Child Health and Human Development-Best Pharmaceuticals for Children Act Pediatric Formulation Initiatives Workshop-Pediatric Biopharmaceutics Classification System Working Group. *Clinical Therapeutics* **2012**, *34* (11), S11-S24.
6. Shawahna, R., Pediatric Biopharmaceutical Classification System: Using Age-Appropriate Initial Gastric Volume. *The AAPS Journal* **2016**, *18* (3), 728-736.
7. Guo, H.; Wang, C.; Liu, Z.; Gu, H.; Li, Y.; Zhao, L.; Hou, R.; Zhu, J.; Ho, H.; Hao, Z., Quantification of in vivo gastric fluid volume in Bama miniature pigs in fasted state. *Biopharmaceutics and Drug Disposition* **2018**, *39* (8), 403-407.
8. Grimm, M.; Koziolk, M.; Kühn, J. P.; Weitschies, W., Interindividual and intraindividual variability of fasted state gastric fluid volume and gastric emptying of water. *European Journal of Pharmaceutics and Biopharmaceutics* **2018**, *127*, 309-317.
9. Grimm, M.; Koziolk, M.; Saleh, M.; Schneider, F.; Garbacz, G.; Kühn, J. P.; Weitschies, W., Gastric Emptying and Small Bowel Water Content after Administration of Grapefruit Juice Compared to Water and Isocaloric Solutions of Glucose and Fructose: A Four-Way Crossover MRI Pilot Study in Healthy Subjects. *Molecular Pharmaceutics* **2018**, *15* (2), 548-559.
10. Mudie, D. M.; Murray, K.; Hoad, C. L.; Pritchard, S. E.; Garnett, M. C.; Amidon, G. L.; Gowland, P. A.; Spiller, R. C.; Amidon, G. E.; Marciani, L., Quantification of gastrointestinal liquid volumes and distribution following a 240 mL dose of water in the fasted state. *Molecular Pharmaceutics* **2014**, *11* (9), 3039-3047.
11. Reppas, C.; Karatza, E.; Goumas, C.; Markopoulos, C.; Vertzoni, M., Characterization of Contents of Distal Ileum and Cecum to Which Drugs/Drug Products are Exposed During Bioavailability/Bioequivalence Studies in Healthy Adults. *Pharmaceutical Research* **2015**, *32* (10), 3338-3349.
12. Riethorst, D.; Mols, R.; Duchateau, G.; Tack, J.; Brouwers, J.; Augustijns, P., Characterization of Human Duodenal Fluids in Fasted and Fed State Conditions. *Journal of Pharmaceutical Sciences* **2016**, *105* (2), 673-681.
13. Clarysse, S.; Tack, J.; Lammert, F.; Duchateau, G.; Reppas, C.; Augustijns, P., Postprandial evolution in composition and characteristics of human duodenal fluids in different nutritional states. *Journal of Pharmaceutical Sciences* **2009**, *98* (3), 1177-1192.
14. Fuchs, A.; Dressman, J. B., Composition and physicochemical properties of fasted-state human duodenal and jejunal fluid: A critical evaluation of the available data. *Journal of Pharmaceutical Sciences* **2014**, *103* (11), 3398-3411.

15. Van Den Abeele, J.; Rayyan, M.; Hoffman, I.; Van de Vijver, E.; Zhu, W.; Augustijns, P., Gastric fluid composition in a paediatric population: Age-dependent changes relevant for gastrointestinal drug disposition. *European Journal of Pharmaceutical Sciences* **2018**, *123*, 301-311.
16. Schiller, C.; Frohlich, C. P.; Giessmann, T.; Siegmund, W.; Monnikes, H.; Hosten, N.;
440 Weitschies, W., Intestinal fluid volumes and transit of dosage forms as assessed by magnetic resonance imaging. *Alimentary pharmacology & therapeutics* **2005**, *22* (10), 971-9.
17. Sutton, S. C., Role of physiological intestinal water in oral absorption. *The AAPS journal* **2009**, *11* (2), 277-285.
18. Luoma, E. K.; Raininko, R.; Nummi, P. J.; Luukkonen, R.; Manninen, H. I.; Riihimaki, H. A.,
445 Suitability of cerebrospinal fluid as a signal-intensity reference on MRI: evaluation of signal-intensity variations in the lumbosacral dural sac. *Neuroradiology* **1997**, *39* (10), 728-32.
19. Hoad, C. L.; Marciani, L.; Foley, S.; Totman, J. J.; Wright, J.; Bush, D.; Cox, E. F.; Campbell, E.; Spiller, R. C.; Gowland, P. A., Non-invasive quantification of small bowel water content by MRI: a validation study. *Physics in Medicine and Biology* **2007**, *52* (23), 6909-6922.
- 450 20. Abràmoff, M. D.; Magalhães, P. J.; Ram, S. J., Image processing with imageJ. *Biophotonics International* **2004**, *11* (7), 36-41.
21. Fruehauf, H.; Goetze, O.; Steingoetter, A.; Kwiatek, M.; Boesiger, P.; Thumshirn, M.; Schwizer, W.; Fried, M., Intersubject and intrasubject variability of gastric volumes in response to isocaloric liquid meals in functional dyspepsia and health2. *Neurogastroenterology & Motility* **2007**, *19*
455 (7), 553-561.
22. Meakin, G.; Dingwall, A. E.; Addison, G. M., EFFECTS OF FASTING AND ORAL PREMEDICATION ON THE pH AND VOLUME OF GASTRIC ASPIRATE IN CHILDREN. *British Journal of Anaesthesia* **1987**, *59* (6), 678-682.
23. Crawford, M.; Lerman, J.; Christensen, S.; Farrow-Gillespie, A., Effects of duration of fasting on gastric fluid pH and volume in healthy children. *Anesthesia and Analgesia* **1990**, *71* (4), 400-403.
- 460 24. Nader, A. M.; Quinney, S. K.; Fadda, H. M.; Foster, D. R., Effect of Gastric Fluid Volume on the In Vitro Dissolution and In Vivo Absorption of BCS Class II Drugs: a Case Study with Nifedipine. *The AAPS journal* **2016**, *18* (4), 981-988.
25. Hens, B.; Van Den Abeele, J.; Rubbens, J.; Keirsebilck, M.; Roelens, J.; Schreurs, C.; Verheyen, K.; Casteels, M.; Laekeman, G.; Augustijns, P., Evaluation of real-life dosing of oral medicines with respect to fluid and food intake in a Dutch-speaking population. *Journal of Clinical Pharmacy and Therapeutics* **2017**, *42* (4), 467-474.
- 465 26. National Clinical Guideline Centre (UK). Nocturnal Enuresis: The Management of Bedwetting in Children and Young People. The Management of Bedwetting in Children and Young People. London: Royal College of Physicians (UK); 2010. NICE Clinical Guidelines, No. 111.) 7, Fluid and diet restriction for the management of bedwetting. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK62715/>.
- 470 27. Steingoetter, A.; Fox, M.; Treier, R.; Weishaupt, D.; Marincek, B.; Boesiger, P.; Fried, M.; Schwizer, W., Effects of posture on the physiology of gastric emptying: a magnetic resonance imaging study. *Scandinavian journal of gastroenterology* **2006**, *41* (10), 1155-64.
- 475 28. Wollmer, E.; Neal, G.; Whitaker, M. J.; Margetson, D.; Klein, S., Biorelevant in vitro assessment of dissolution and compatibility properties of a novel paediatric hydrocortisone drug product following exposure of the drug product to child-appropriate administration fluids. *European Journal of Pharmaceutics and Biopharmaceutics* **2018**, *133*, 277-284.