

Identifying the at risk smokers

Madan, A ; Turner, Alice

DOI:

[10.1183/13993003.01613-2019](https://doi.org/10.1183/13993003.01613-2019)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Madan, A & Turner, A 2019, 'Identifying the at risk smokers: who goes on to get COPD?', *The European respiratory journal*, vol. 54, no. 4, 1901613. <https://doi.org/10.1183/13993003.01613-2019>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is an author-submitted, peer-reviewed version of a manuscript that has been accepted for publication in the European Respiratory Journal, prior to copy-editing, formatting and typesetting. This version of the manuscript may not be duplicated or reproduced without prior permission from the copyright owner, the European Respiratory Society. The publisher is not responsible or liable for any errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final, copy-edited, published article, which is the version of record, is available without a subscription 18 months after the date of issue publication.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Identifying the at risk smokers: who goes on to get COPD?

A Madan¹ & Alice M Turner^{1,2}

1 University Hospitals Birmingham, Heartlands hospital, B9 5SS, UK

2 Institute of Applied Health Research, University of Birmingham, Birmingham, B15 2TT, UK

Corresponding author: Alice M Turner, University of Birmingham, address as above. Email:
a.m.turner@bham.ac.uk

Keywords: Chronic obstructive pulmonary disease; emphysema; cigarette smoking; tomography, X-ray computed; prognosis; screening

Word count: 1496

Chronic obstructive Pulmonary Disease (COPD) is a common disease that predominantly affects smokers and is characterised by persistent and usually progressive airflow limitation(1). Whilst international bodies such as the Global Obstructive Lung Disease (GOLD) initiative have varied in their classification system for COPD severity over the years, there has long been recognition that pathologically(2) and clinically(3) relevant disease may be present before spirometry becomes abnormal – indeed GOLD included ‘at risk’ as a disease stage in early iterations of their documents on COPD. In addition there is debate on what constitutes early disease, in terms of chronology, versus what constitutes mild disease, in terms of either physiology or impact on the patient – these may not always be the same. For example, a patient who has developed COPD at a late age, through the aetiology of poor lung growth in early life(4) could have mild disease if symptom burden and physiology were not markedly impaired, but this is not early, since the processes driving the diagnosis have been present lifelong. Identifying those patients who have early disease which is likely to progress, particularly as we move toward an era of potentially disease modifying therapies is an area of great clinical need.

The research reported in this issue of the journal by Arjomandi et al [editorial team to insert reference for paper ERJ-02214-2018.R1] is important as it contributes to this debate, and may change the way clinicians prognosticate and predict progression to COPD. The study was conducted in part because inflammatory signals in symptomatic smokers’ airway secretions have been reported whose pattern was consistent with COPD(5); this suggests that early pathological changes are present in these patients which are not detected by spirometry. Since only 20% of smokers are classified as having COPD, yet many are symptomatic, measurements other than spirometry may be required to describe their disease, and determine risk of progression to COPD. Several studies have suggested that small airways disease (SAD) is the earliest abnormality to occur in COPD (6, 7), perhaps because of altered tissue repair mechanisms (8). However emphysema can also occur prior to spirometric abnormalities, and when this occurs the extent appears to be a risk factor for respiratory death, even at low smoke exposures, or without a diagnosis of COPD(9). Consequently any study of patients at risk of progression to COPD needs to include a reliable method of assessing both SAD and emphysema. CT scanning is attractive in this regard since it can be used to quantify emphysema, typically using lung density from an inspiratory scan at total lung capacity, and also to measure SAD – a number of techniques have been used for this, of which parametric response mapping (PRM) (10) involving use of paired inspiratory and expiratory scans is now well known.

SPIROMICS is a multi-centre observational study designed to guide further development of treatment for COPD through characterising sub-populations over time(11). The paper in this issue [ERJ-02214-2018.R1] reports a post-hoc analysis of 849 current and ex-smokers (≥ 20 year pack history) with preserved spirometry within the SPIROMICS cohort, and specifically asks whether CT guided lung volumes defined by the ratio of residual volume to total lung capacity (RV_{CT}/TLC_{CT}) could predict future lung function decline and progression to COPD. The authors found that increased air trapping based on these CT-derived volumes predicts faster spirometric decline and progression to COPD, such that over 2.7 years of follow up, subjects with RV_{CT}/TLC_{CT} in the upper tertile had a faster rate of decline in FEV1/FEVC (0.66% per year (95%CI=0.06%-1.27%) , $p=0.015$) when compared to the lower tertile. They were also more likely develop spirometric COPD (odds ratio = 5.7, $p < 0.001$) compared to the lower tertile, regardless of baseline characteristics such as baseline spirometry and smoking status. Whilst they were not able to report physiology confirming the RV/TLC ratio by plethysmography, other studies have shown a reasonable correlation between gas trapping on CT and this more traditional volume measure(12). When all the CT indices of air trapping were included in the model, including PRM of functional SAD and percent voxel on expiratory CT (as a measure of emphysema), only RV_{CT}/TLC_{CT}

remained significant, suggesting that the critical feature is SAD and not emphysema. Whilst SAD certainly appeared more important in these multivariate models the tables of characteristics show that at baseline emphysema was more severe as the RV_{CT}/TLC_{CT} tertile rose, such that both components probably play a role. Irrespective of the cause of gas trapping (due to SAD or emphysema) it appeared clinically important, in that those with the highest tertile of RV_{CT}/TLC_{CT} had a lower exercise capacity, as measured by 6 minute walk test distance, which exceeded at least one definition of the minimum clinical important difference for this measure(13). Interestingly no association was seen with exacerbations of COPD requiring hospital admission for any of the three tertiles; this may be explicable because lower FEV1 is a major factor in admission(14) and any patients who moved from normal spirometry to COPD within the short follow up period then had mild disease. Consistent with this interpretation only 36 subjects had severe exacerbations, and follow up FEV1 in the highest risk group was 94% predicted. Furthermore symptomatic deterioration was found using only one of the three SPIROMICS study questionnaires, with the other two questionnaires showing non- significant trends. Again this finding probably reflects the mild disease present even at follow up, alongside the fact that by the time symptoms develop, disease may no longer be early, even if it is mild.

The study has several strengths. Firstly the authors carefully considered the validity of their CT scan derived lung volumes by selecting those whose CT derived vital capacity matched the slow FVC obtained at spirometry for their primary analysis; this resulted in 618/814 subjects with high quality complete data being analysed. Whilst this process excluded a significant proportion of patients subsequent sensitivity analysis demonstrated that this had no impact on the direction of association of RV_{CT}/TLC_{CT} with subsequent decline. They also considered the impact of a number of known risk factors for lung function decline, such as active smoking, amount of smoke exposure and bronchodilator reversibility, by including these features in either multivariable or sensitivity analyses. These factors are known to influence decline not only in COPD(15) but in other 'at risk' groups, such as those with a genetic susceptibility to COPD, like alpha 1 antitrypsin deficiency(16) or who are exposed to biomass smoke(17). These analyses demonstrated that smoking status and pack years smoked did not influence decline; considering that the SPIROMICS cohort all have significant smoke exposure the authors concluded that after 20 pack years exposure there are some subjects who are predisposed to decline ('susceptible smokers'), and that they might be identified by their higher RV_{CT}/TLC_{CT} . None of the other confounders markedly influenced the primary reported association of gas trapping with decline.

However, as with many cohort studies, follow up data appeared more difficult to collect than baseline; consequently only 496 out of 618 subjects had a follow up spirometry result. This equates to nearly 20% of subjects lacking follow up spirometry. Furthermore, of the 496 subjects, 44 only had one follow-up spirometry result yet two or more spirometry results are generally required over this time period to see a reliable trend in lung function decline. This meant that only 157 of 618 subjects in the highest RV_{CT}/TLC_{CT} tertile had three follow up spirometry results over which to calculate decline, with 122 having no follow up spirometry results. The reduction in numbers naturally results in loss of power, and whilst it does not diminish the significance of their findings, it suggests some caution in interpretation is required, and that replication will be important. In addition the median follow-up for the reported data was only 2.7 years; it will be important to see outcome data after a longer period of follow-up, given the chronicity and slow progression generally seen in COPD. The authors recognized this when discussing the limitations of their work.

Finally, it is worth considering the messages this paper brings to clinical practice. Clinicians, particularly in primary care, would welcome a way to identify which smokers should be more closely followed up so that treatment for significant COPD can be instituted quickly and through so doing

deterioration due to deconditioning and exacerbations reduced. Whilst the authors have demonstrated that CT guided lung volumes can accurately predict progression to COPD, the feasibility of using this technique in clinical practice due to cost and radiation exposure is questionable. Additional research into techniques to identify gas trapping using measures possible in primary care will therefore be important. However, in the era of CT screening for lung cancer, which generally occurs in smoke exposed people, due to their higher risk of malignancy relative to never smokers, many countries will have imaging data available on large numbers of subjects at risk of progression to COPD; using these visits (and CT images) for more than just lung cancer screening(18) could enhance existing cost-effectiveness(19) whilst also being acceptable to patients.

References

1. Global Initiative for Obstructive Lung Disease [Available from: www.goldcopd.com.
2. Higham A, Quinn AM, Cancado JED, Singh D. The pathology of small airways disease in COPD: historical aspects and future directions. *Respir Res*. 2019;20(1):49.
3. Woodruff PG, Barr RG, Bleecker E, Christenson SA, Couper D, Curtis JL, et al. Clinical Significance of Symptoms in Smokers with Preserved Pulmonary Function. *N Engl J Med*. 2016;374(19):1811-21.
4. Savran O, Ulrik CS. Early life insults as determinants of chronic obstructive pulmonary disease in adult life. *Int J Chron Obstruct Pulmon Dis*. 2018;13:683-93.
5. Garudadri S, Woodruff PG, Han MK, Curtis JL, Barr RG, Bleecker ER, et al. Systemic Markers of Inflammation in Smokers With Symptoms Despite Preserved Spirometry in SPIROMICS. *Chest*. 2019;155(5):908-17.
6. Hogg JC, Chu F, Utokaparch S, Woods R, Elliott WM, Buzatu L, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *The New England journal of medicine*. 2004;350(26):2645-53.
7. Koo HK, Vasilescu DM, Booth S, Hsieh A, Katsamenis OL, Fishbane N, et al. Small airways disease in mild and moderate chronic obstructive pulmonary disease: a cross-sectional study. *The Lancet Respiratory medicine*. 2018;6(8):591-602.
8. Gosselink JV, Hayashi S, Elliott WM, Xing L, Chan B, Yang L, et al. Differential expression of tissue repair genes in the pathogenesis of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2010;181(12):1329-35.
9. Oelsner EC, Carr JJ, Enright PL, Hoffman EA, Folsom AR, Kawut SM, et al. Per cent emphysema is associated with respiratory and lung cancer mortality in the general population: a cohort study. *Thorax*. 2016;71(7):624-32.
10. Galban CJ, Han MK, Boes JL, Chughtai KA, Meyer CR, Johnson TD, et al. Computed tomography-based biomarker provides unique signature for diagnosis of COPD phenotypes and disease progression. *Nat Med*. 2012;18(11):1711-5.
11. Couper D, LaVange LM, Han M, Barr RG, Bleecker E, Hoffman EA, et al. Design of the Subpopulations and Intermediate Outcomes in COPD Study (SPIROMICS). *Thorax*. 2014;69(5):491-4.
12. Barjaktarevic I, Springmeyer S, Gonzalez X, Sirokman W, Coxson HO, Cooper CB. Diffusing capacity for carbon monoxide correlates best with tissue volume from quantitative CT scanning analysis. *Chest*. 2015;147(6):1485-93.
13. Polkey MI, Spruit MA, Edwards LD, Watkins ML, Pinto-Plata V, Vestbo J, et al. Six-minute-walk test in chronic obstructive pulmonary disease: minimal clinically important difference for death or hospitalization. *Am J Respir Crit Care Med*. 2013;187(4):382-6.
14. Mullerova H, Maselli DJ, Locantore N, Vestbo J, Hurst JR, Wedzicha J, et al. Hospitalized Exacerbations of Chronic Obstructive Pulmonary Disease: Risk Factors and Outcomes in the ECLIPSE Cohort. *Chest*. 2014.
15. Vestbo J, Edwards LD, Scanlon PD, Yates JC, Agusti A, Bakke P, et al. Changes in forced expiratory volume in 1 second over time in COPD. *N Engl J Med*. 2011;365(13):1184-92.

16. Dawkins PA, Dawkins CL, Wood AM, Nightingale PG, Stockley JA, Stockley RA. Rate of progression of lung function impairment in alpha1-antitrypsin deficiency. *Eur Respir J*. 2009;33(6):1338-44.
17. Ramirez-Venegas A, Sansores RH, Quintana-Carrillo RH, Velazquez-Uncal M, Hernandez-Zenteno RJ, Sanchez-Romero C, et al. FEV1 decline in patients with chronic obstructive pulmonary disease associated with biomass exposure. *Am J Respir Crit Care Med*. 2014;190(9):996-1002.
18. Balata H, Blandin Knight S, Barber P, Colligan D, Crosbie EJ, Duerden R, et al. Targeted lung cancer screening selects individuals at high risk of cardiovascular disease. *Lung Cancer*. 2018;124:148-53.
19. Hinde S, Crilly T, Balata H, Bartlett R, Crilly J, Barber P, et al. The cost-effectiveness of the Manchester 'lung health checks', a community-based lung cancer low-dose CT screening pilot. *Lung Cancer*. 2018;126:119-24.