

## High frequency DOM dynamics in urban rivers revealed by in-situ fluorometry

Bradley, Chris; Croghan, Danny; Khamis, Kieran; Van Loon, Anne; Sadler, Jon; Hannah, David

*License:*  
Creative Commons: Attribution (CC BY)

*Document Version*  
Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*  
Bradley, C, Croghan, D, Khamis, K, Van Loon, A, Sadler, J & Hannah, D 2018, High frequency DOM dynamics in urban rivers revealed by in-situ fluorometry. in *Geophysical Research Abstracts*. vol. 20, 10263, European Geosciences Union, European Geosciences Union General Assembly 2018, Vienna, Austria, 8/04/18.  
<<https://meetingorganizer.copernicus.org/EGU2018/EGU2018-10263.pdf>>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**  
Checked for eligibility: 14/10/2019

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



## **High frequency DOM dynamics in urban rivers revealed by in-situ fluorometry.**

Chris Bradley, Danny Croghan, Kieran Khamis, Anne van Loon, Jon Sadler, and David Hannah  
University of Birmingham, School of Geography, Earth and Environmental Sciences, Birmingham, United Kingdom  
(c.bradley@bham.ac.uk)

To-date Dissolved Organic Matter (DOM) dynamics have been quantified poorly in urban rivers despite substantial water quality problems. Research has been hindered by the low temporal resolution of observations and over-reliance on manual sampling which often fail to capture precipitation events and diurnal dynamics. High frequency data are essential to quantify DOM fluxes/loads and understand DOM furnishing and transport processes. While recent advances in optical sensor technology are yielding new high resolution DOM information, there is a lack of consensus regarding the monitoring resolution required for urban systems, with no studies routinely monitoring at <15 min time steps.

High-frequency monitoring (5 min resolution; 4 week duration) was conducted on a headwater urban stream in Birmingham, UK to determine the optimum temporal resolution to characterize DOM event dynamics. A through-flow GGNU-30 monitored wavelengths corresponding to tryptophan-like fluorescence (TLF; Peak T1) (Ex 285 nm/ Em 345 nm) and humic-like fluorescence (HLF; Peak C) (Ex 365 nm/Em 490 nm).

Five minute temporal resolution reveals distinct peaks in TLF and HLF indicating separate DOM pathways are apparent during storms. Initial pulses in TLF and HLF appear to relate to scouring of streambed biofilms, whilst secondary TLF and HLF peaks may relate to terrestrial OM inputs from the wider catchment. High temporal variation occurs during storm events in TLF and HLF intensity: TLF intensity is highest during the rising limb of the hydrograph and can rapidly decline thereafter, indicating the importance of fast flow-path and close proximity sources. HLF also peaks on the rising limb, but exhibits little decline as flow recedes, with a hysteresis effect apparent. Furthermore, the ratio of TLF:HLF, which aids in identifying inputs from Combined Sewage Overflow suggests smaller events (< 3mm precipitation) tend to be more Peak C dominated, whereas large events (> 3mm precipitation) tend to be more Peak T dominated, indicating that sewage discharges are more prominent during larger events.

Our work highlights the need for future studies to utilise shorter temporal scales than previously used to monitor urban DOM dynamics. The application of higher frequency monitoring enables the identification of finer-scale patterns and subsequently aids in deciphering the sources and pathways controlling urban DOM dynamics.