

Minitablets may be an acceptable alternative to liquid in infants and young children

Batchelor, Hannah

DOI:

[10.1016/j.jpeds.2018.12.075](https://doi.org/10.1016/j.jpeds.2018.12.075)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Batchelor, H 2019, 'Minitablets may be an acceptable alternative to liquid in infants and young children', *Journal of Pediatrics*, vol. 206, pp. 298-301. <https://doi.org/10.1016/j.jpeds.2018.12.075>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Batchelor, H. (2019) Minitablets may be an acceptable alternative to liquid in infants and young children, *The Journal of Pediatrics*, 206:298-301; <https://doi.org/10.1016/j.jpeds.2018.12.075>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Acceptability of minitablets in children is important to those working in pharmaceutical development to ensure medicines meet the needs of patients. Previous work showed that tablets are superior to alternative oral formulations in children aged 1-4 years¹. Klingmann et al provide evidence that multiple minitablets (up to 400) are accepted by children aged 2-5 years. The study is limited by ethical requirements to include only children who are (i) able to swallow and (ii) are compliant and willing to accept the study procedures both of which may introduce bias in study outcomes. Reluctance to use solid dosage forms is often attributed to risks associated with choking yet no deglutition issues were detected with minitablets in a total of 372 participants. Taste is not an issue for placebo minitablets which may not be valid for a commercial product. Future work should focus on the acceptability of commercial minitablets with the associated hurdles associated with a real paediatric population and a real drug product.

1. Van Riet-Nales, D.A., De Neef, B.J., Schobben, A.F.A.M., Ferreira, J.A., Egberts, T.C.G., Rademaker, C.M.A. Acceptability of different oral formulations in infants and preschool children (2013) *Archives of Disease in Childhood*, 98 (9), pp. 725-731.