

$\gamma\delta$ TCR Recognition of MR1

Willcox, Benjamin E; Mohammed, Fiyaz; Willcox, Carrie R

DOI:

[10.1016/j.tibs.2020.03.012](https://doi.org/10.1016/j.tibs.2020.03.012)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Willcox, BE, Mohammed, F & Willcox, CR 2020, ' $\gamma\delta$ TCR Recognition of MR1: Adapting to Life on the Flip Side', *Trends in Biochemical Sciences*, vol. 45, no. 7, pp. 551-553. <https://doi.org/10.1016/j.tibs.2020.03.012>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Acknowledgments

This work was funded by a Medical Research Council (<https://mrc.ukri.org/>) grant, MR/P019374/1, to M.B.

¹Department of Life Sciences, Imperial College London, London, SW7 2AZ, UK

*Correspondence:

mbeeb@imperial.ac.uk (M. Beeby).

®Twitter: @BeebyLab (M. Beeby).

<https://doi.org/10.1016/j.tibs.2020.03.006>

© 2020 Elsevier Ltd. All rights reserved.

References

1. von Kügelgen, A. *et al.* (2020) In situ structure of an intact lipopolysaccharide-bound bacterial surface layer. *Cell* 180, 348–358
2. Fagan, R.P. and Fairweather, N.F. (2014) Biogenesis and functions of bacterial S-layers. *Nat. Rev. Microbiol.* 12, 211–222
3. Kalynych, S. *et al.* (2014) Progress in understanding the assembly process of bacterial O-antigen. *FEMS Microbiol. Rev.* 38, 1048–1065
4. Neidhardt, F.C. *et al.*, eds (1987) *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*, American Society for Microbiology
5. Cohen, E.J. *et al.* (2017) Nanoscale-length control of the flagellar driveshaft requires hitting the tethered outer membrane. *Science* 356, 197–200
6. Mühlradt, P.F. *et al.* (1973) Outer membrane of *Salmonella*. *Eur. J. Biochem.* 35, 471–481
7. Ursell, T.S. *et al.* (2012) Analysis of surface protein expression reveals the growth pattern of the Gram-negative outer membrane. *PLoS Comput. Biol.* 8, e1002680
8. Kleanthous, C. *et al.* (2015) Protein–protein interactions and the spatiotemporal dynamics of bacterial outer membrane proteins. *Curr. Opin. Struct. Biol.* 35, 109–115
9. Comerci, C.J. *et al.* (2019) Topologically-guided continuous protein crystallization controls bacterial surface layer self-assembly. *Nat. Commun.* 10, 1–10
10. Delarue, M. *et al.* (2018) mTORC1 controls phase separation and the biophysical properties of the cytoplasm by tuning crowding. *Cell* 174, 338–349

Spotlight

$\gamma\delta$ TCR Recognition of MR1: Adapting to Life on the Flip Side

Benjamin E. Willcox,^{1,2,*}
Fiyaz Mohammed,^{1,2} and
Carrie R. Willcox^{1,2}



Nonclassical class I MHC-like molecules are ligands for several unconventional T cell populations.

Recently, Le Nours *et al.* identified human $\gamma\delta$ T cells recognising MHC-related protein-1 (MR1) via their T cell receptor (TCR). Also recognised by the $\alpha\beta$ -TCR of mucosal associated invariant T cells, MR1 interacts with specific $\gamma\delta$ -TCRs using strikingly diverse binding modes, suggesting fundamental differences in $\gamma\delta$ T cell recognition.

$\gamma\delta$ T cells, defined by their surface expression of paired γ and δ T cell receptor (TCR) chain heterodimers, have been retained throughout vertebrate evolution and play critical roles in host immunity in diverse settings, including infection, antitumour immunity, and immune regulation [1,2]. They are also of increasing therapeutic interest. Although it is widely accepted that unlike $\alpha\beta$ T cells, they do not recognise peptide-MHC molecules, the question of what antigens they do recognise via their TCR still remains substantially unresolved.

Since the discovery of $\gamma\delta$ T cells, diverse molecules have been proposed as candidate $\gamma\delta$ -TCR ligands [3]. While recent evidence has confirmed butyrophilin/butyrophilin-like (BTN/BTNL) family molecules as direct TCR ligands for $\gamma\delta$ T cell populations bearing specific TCR γ chain variable regions (either V γ 4/V γ 7 [4] or V γ 9 chains [5,6]), the mouse nonclassical class I MHC molecules T10 and T22 were the first $\gamma\delta$ -TCR ligands to be confirmed biochemically [7]. Since then, $\gamma\delta$ T cells capable of interacting via their TCR with the nonclassical class I MHC molecule CD1d have also been defined [8].

Recently, Le Nours and colleagues have made an important step forward by demonstrating a third category of nonclassical class I MHC molecule, MHC-related protein 1 (MR1), is also a target for $\gamma\delta$ -TCR binding [9]. Their study combines use of MR1-tetramer staining to identify MR1-binding $\gamma\delta$ T cell populations, surface

plasmon resonance (SPR) to assess direct $\gamma\delta$ -TCR/MR1 binding, and structural techniques to establish relevant binding modes. Their findings significantly advance our understanding of MR1 and may hold some fundamental lessons regarding $\gamma\delta$ -TCR recognition itself.

Adaptive $\gamma\delta$ T Cell Recognition of a Monomorphic Ligand

CD1d and MR1 are established recognition targets for defined $\alpha\beta$ T cell populations, namely invariant natural killer T cells (iNKTs) and mucosa associated invariant T cells (MAITs), respectively. Aligning with the monomorphic nature of these ligands, both iNKTs and MAITs express a highly restricted TCR repertoire and also exhibit distinct innate-like phenotypes relative to the bulk $\alpha\beta$ T cell compartment. Using MR1 tetramers, Le Nours *et al.* showed the situation is very different for MR1-binding $\gamma\delta$ T cells.

In most people, MR1-specific $\gamma\delta$ T cells comprised a low percentage (~0.1%) of $\gamma\delta$ T cells. Strikingly, their TCR repertoire was diverse, reflecting the TCR-diverse adaptive-like V δ 2^{neg} repertoire as a whole and chiefly focussed on the prevalent V δ 1 and V δ 3 subsets, combined with a broad range of V γ chains. Also, MR1-specific $\gamma\delta$ T cells phenotypically resembled the entire $\gamma\delta$ T cell pool. By contrast, the semi-invariant, innate-like V γ 9V δ 2 T cell subset, which bears a highly restricted TCR repertoire, was not a source of MR1-specific $\gamma\delta$ T cells. These features closely mirror those of CD1d-specific $\gamma\delta$ T cells (and $\gamma\delta$ T cells specific for the exogenous model antigen phycoerythrin), but contrast with properties of iNKTs and MAITs. Relative to $\alpha\beta$ T cells, $\gamma\delta$ T cell recognition of nonclassical class I MHC molecules may therefore be fundamentally skewed towards highly TCR-diverse, adaptive-like $\gamma\delta$ subsets, which are thought to bind a diverse array of ligands.


Whilst limited phenotypic analysis of MR1-specific $\gamma\delta$ T cells was carried out, their differentiation status was not defined. Addressing this question, highly relevant for adaptive compartments, would clarify if MR1-specific $\gamma\delta$ T cells reside within the T_{effector} subpopulation, consistent with *bona fide* MR1-directed adaptive T_{effector} responses, or alternatively within the T_{naive} subpopulation, which lacks effector capability and would be more suggestive of potential adaptive reactivities [10] yet to encounter MR1 *in vivo*. *In vitro* assays involving transduction of MR1-binding TCRs into Jurkat T cells showed that although CD69 upregulation was not always observed, MAP kinase/ERK kinase activation was universal, confirming a potential to support TCR triggering. Although

low levels of MR1-specific T cells were detected in most individuals, MR1-tetramer-positive cells were enriched in some individual samples, including in newly diagnosed coeliac disease and Merkel cell carcinoma. This finding suggests both TCR-diverse T_{naive} and clonally focused T_{effector} subpopulations may contribute to the MR1-specific $\gamma\delta$ T cell pool; the latter could contribute to physiological adaptive $\gamma\delta$ T_{effector} responses in some individuals. Future studies will no doubt shed light on these questions.

Diverse Modes of Antigen-Agnostic $\gamma\delta$ TCR Binding to MR1

Le Nours and colleagues also outlined the molecular basis of $\gamma\delta$ -TCR/MR1

interaction. SPR binding studies revealed MR1-binding $\gamma\delta$ -TCRs tested were largely ‘antigen agnostic’ and either entirely unaffected or only slightly impacted by the presence/absence of MR1-bound antigen, suggesting potential ‘inherent autoreactivity’ to MR1-expressing cells even in the absence of antigenic challenge. Although iNKT and MAIT TCR/ligand recognition has also been linked to ‘inherent autoreactivity’, this operates via binding modes apparently exclusively involving interaction of $\alpha\beta$ -TCR CDR loops with the $\alpha 1\alpha 2$ platform (Figure 1A). Moreover, conserved iNKT and MAIT TCR V-region usage and respective germline-encoded CDR1/2 loops provide a clear basis for such semi-invariant interactions, which appear literally and immunologically ‘restricted’



Trends in Biochemical Sciences

Figure 1. Overview of the MAIT $\alpha\beta$ TCR-MR1-5-OP-RU, G7 $\gamma\delta$ TCR-MR1-5-OP-RU, and CD8 α -HLA-A2 Complexes. (A) Cartoon representation of the MAIT $\alpha\beta$ TCR-MR1-5-OP-RU complex (PDB ID: 4NQC): MR1, brown; β_2 -microglobulin (β_2M), blue; 5-OP-RU, green; α -chain, light green; β -chain, cyan. (B) Cartoon representation of the G7 $\gamma\delta$ TCR-MR1-5-OP-RU complex (PDB ID: 6MWR): MR1, brown; β_2M , blue; 5-OP-RU, green; V γ 9 chain, pale cyan; V δ 1 chain, salmon. (C) Cartoon representation of the CD8 α -HLA-A2 complex (PDB ID: 1AKJ): HLA-A2, orange; β_2M , blue; peptide, hot pink; CD8 α , red and yellow. Ig-like variable and constant domains for α , β , γ , and δ chains are indicated by V α , C α , V β , C β , V γ , C γ , and V δ , C δ respectively. Abbreviations: MAIT, mucosa associated invariant T cell; MR1, MHC-related protein 1; TCR, T cell receptor.

to the $\alpha 1\alpha 2$ platform of CD1d or MR1, allowing potential for discriminating presence/absence and nature of bound antigen.

By contrast, mutational analyses suggested that collectively, the MR1-binding $\gamma\delta$ -TCR pool was not limited to interaction with the upper-face of the $\alpha 1\alpha 2$ platform, but also contained TCR specificities recognising the membrane-proximal 'flip-side' of MR1, predominantly to the $\alpha 3$ domain. X-ray crystallographic analysis confirmed this highly novel binding mode. Importantly, 'flip-side' interaction was consistent with antigen 'agnosticism' and involved no contacts to upper-facing $\alpha 1\alpha 2$ helical platform residues, instead predominantly featuring $\alpha 3$ domain contacts, with additional interactions to the platform's underside (Figure 1B). Consistent with diverse V γ usage in the MR1-binding $\gamma\delta$ -TCR pool, interaction was dominated by V δ -mediated contacts. Moreover, while some CDR1 δ -mediated involvement was evident, V δ interactions involved critical hydrophobic contacts formed by CDR3 δ residues, consistent with only a small proportion of the extremely diverse V $\delta 1$ TCR repertoire satisfying the molecular criteria for MR1 recognition. This mode resembled CD8 $\alpha\alpha$ /class I MHC recognition (Figure 1C), which itself was likened to antibody/antigen interaction [11]. These observations confirm that $\gamma\delta$ T cell recognition of MR1 is indeed fundamentally different to CD1d/MR1-restricted recognition by semi-invariant iNKTs and MAITs.

In summary, the identification of MR1-binding $\gamma\delta$ T cells is a significant advance for both MR1 and $\gamma\delta$ T cell biology and

should be applauded. By contrast to iNKTs and MAITs that now have established contributions to immune regulation, including in diverse models of infection/disease, the physiological role and importance of $\gamma\delta$ T cells that recognise nonclassical class I MHC molecules has remained largely unclear since their initial identification 20 years ago. In this context, the immunobiological meaning and relevance of antigen-agnostic recognition of MR1 by $\gamma\delta$ T cells is currently unclear. Moreover, future studies should consider the parallel and nonmutually exclusive possibilities that MR1 interactions with $\gamma\delta$ -TCRs either contribute to physiological adaptive $\gamma\delta$ T cell effector immune responses, or alternatively in some cases largely represent potential autoreactivities. In this second scenario, the presence of MR1-specific cells may reflect the fundamental potential of the adaptive $\gamma\delta$ -TCR repertoire to recognise diverse self-antigens, from which particular autoreactive TCR specificities may be selected to differentiate into T_{effector} cells to support adaptive $\gamma\delta$ T cell immunosurveillance following relevant immune challenges. Given the recent finding that MR1-directed $\alpha\beta$ -TCR alloreactive recognition of an antigenically altered form of MR1 can mediate broad antitumour responses [12], it is tempting to speculate on the potential relevance of $\gamma\delta$ -TCR/MR1 interactions in such settings, particularly given established *in vitro* antitumour capabilities of $\gamma\delta$ T cells. The study by Le Nours and colleagues is a fundamental step forward that should pave the way for future studies to address such fascinating questions.

Acknowledgments

This work was supported by the Wellcome Trust, United Kingdom (grants 099266/Z/12/Z and 099266/Z/12/A to B.E.W. supporting C.R.W. and F.M.).

¹Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK

²Cancer Immunology and Immunotherapy Centre, University of Birmingham, Birmingham, UK

*Correspondence:

B.Willcox@bham.ac.uk (B.E. Willcox).

<https://doi.org/10.1016/j.tibs.2020.03.012>

© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

References

1. Silva-Santos, B. *et al.* (2019) Gammadelta T cells: pleiotropic immune effectors with therapeutic potential in cancer. *Nat. Rev. Cancer* 19, 392–404
2. Vantourout, P. and Hayday, A. (2013) Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nat. Rev. Immunol.* 13, 88–100
3. Willcox, B.E. and Willcox, C.R. (2019) Gammadelta TCR ligands: the quest to solve a 500-million-year-old mystery. *Nat. Immunol.* 20, 121–128
4. Willcox, C.R. *et al.* (2019) Butyrophilin-like 3 directly binds a human Vgamma4(+) T cell receptor using a modality distinct from clonally-restricted antigen. *Immunity* 51, 813–825
5. Rigau, M. *et al.* (2020) Butyrophilin 2A1 is essential for phosphoantigen reactivity by gammadelta T cells. *Science* 367, eaay5516
6. Karunakaran, M.M. *et al.* (2020) Essential role of Butyrophilin-2A1 in a composite ligand model of Vgamma9Vdelta2-mediated phosphoantigen sensing. *Immunity* 52, 487–498
7. Crowley, M.P. *et al.* (2000) A population of murine gammadelta T cells that recognize an inducible MHC class Ib molecule. *Science* 287, 314–316
8. Bai, L. *et al.* (2012) The majority of CD1d-sulfatide-specific T cells in human blood use a semiinvariant Vdelta1 TCR. *Eur. J. Immunol.* 42, 2505–2510
9. Le Nours, J. *et al.* (2019) A class of gammadelta T cell receptors recognize the underside of the antigen-presenting molecule MR1. *Science* 366, 1522–1527
10. Davey, M.S. *et al.* (2017) Clonal selection in the human Vdelta1 T cell repertoire indicates gammadelta TCR-dependent adaptive immune surveillance. *Nat. Commun.* 8, 14760
11. Gao, G.F. *et al.* (1997) Crystal structure of the complex between human CD8alpha(alpha) and HLA-A2. *Nature* 387, 630–634
12. Crowther, M.D. *et al.* (2020) Genome-wide CRISPR-Cas9 screening reveals ubiquitous T cell cancer targeting via the monomorphic MHC class I-related protein MR1. *Nat. Immunol.* 21, 178–185